

Midazolam

Statut de médicament essentiel

Description générale

Midazolam is a benzodiazepine with a wide range of effects including anxiolytic, sedative, muscle relaxant, antiepileptic and anterograde amnesic.

INN

Midazolam

Codes ATC

[N05CD08](#)

Type de médicament

Chemical agent

Historique des statuts LME

Ajouté pour la première fois en 2009 ([TRS 958](#)) pour [Palliative care](#)

Ajouté en 2011 ([TRS 965](#)) pour [Sedative-hypnotic and anxiolytic drugs](#)

Ajouté en 2015 ([TRS 994](#)) pour [Status epilepticus](#)

Modifié en 2021 ([TRS 1035](#)) pour [Status epilepticus](#)

Équivalent thérapeutique pour

[lorazepam](#) pour [Status epilepticus](#)

Wikipédia

[Midazolam](#)

DrugBank

[Midazolam](#)

Recommandations

Section

Preoperative medication and sedation for short-term procedures

- Parenteral > General injections > IV: 1 mg per mL in 5 vial; 5 mg per mL in 1 mL vial; 5 mg per mL in 3 mL vial
- Oral > Liquid: 2 mg per mL
- Oral > Solid > dosage form: 15 mg; 7.5 mg

Indications

[Sedative-hypnotic and anxiolytic drugs](#)

Section

Medicines for other common symptoms in palliative care

- Oral > Liquid: 2 mg per mL
- Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial; 5 mg per mL in 1 mL vial; 5 mg per mL in 3 mL vial
- Local > Buccal > Solution: ; 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)
- Oral > Solid > dosage form: 7.5 mg; 15 mg

Indications

[Palliative care](#)

Section

Antiseizure medicines

- Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration); 5 mg per mL in 1 mL vial (for buccal administration); 5 mg per mL in 3 mL vial (for buccal administration)
- Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)

Indications

[Status epilepticus](#)