



General description

Acetylsalicylic acid is used for the management of mild to moderate pain such as headache, acute migraine, transient musculoskeletal pain, dysmenorrhoea and for reducing fever. It is also used for its antiplatelet properties.

Medicine type	Chemical agent
EML status history	First added in 1977 (TRS 615) for Pain Added in 1987 (TRS 770) for Migraine Added in 1989 (TRS 796) for Acute myocardial infarction Added in 1989 (TRS 796) for Cerebral ischaemic stroke Added in 2007 (TRS 950) for Mucocutaneous lymph node syndrome Added in 2007 (TRS 950) for Juvenile idiopathic arthritis Added in 2007 (TRS 950) for Acute rheumatic fever without mention of heart involvement
Therapeutic equivalent for	
Wikipedia	Acetylsalicylic acid ↗
DrugBank	Acetylsalicylic acid ↗

Recommendations

Section Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Solid: 100 to 500 mg

Local > Rectal > Suppository: 50 to 150 mg

Indications

Pain

Section Antimigraine medicines > For treatment of acute attack

Oral > Solid: 300 to 500 mg

Indications

Migraine

Section Anti-platelet medicines

Oral > Solid: 100 mg

Indications

Acute myocardial infarction

Cerebral ischaemic stroke

Section Juvenile joint diseases

Oral > Solid: 100 to 500 mg

Local > Rectal > Suppository: 50 to 150 mg

Indications

Acute rheumatic fever without mention of heart involvement

Juvenile idiopathic arthritis

Mucocutaneous lymph node syndrome

