



## Description générale

Acetylsalicylic acid is used for the management of mild to moderate pain such as headache, acute migraine, transient musculoskeletal pain, dysmenorrhoea and for reducing fever. It is also used for its antiplatelet properties.

<b>Codes ATC</b>	N02BA01 B01AC06
<b>Type de médicament</b>	Chemical agent
<b>Historique des statuts LME</b>	Ajouté pour la première fois en 1977 (TRS 615) pour Pain Ajouté en 1987 (TRS 770) pour Migraine Ajouté en 1989 (TRS 796) pour Acute myocardial infarction Ajouté en 1989 (TRS 796) pour Cerebral ischaemic stroke Ajouté en 2007 (TRS 950) pour Mucocutaneous lymph node syndrome Ajouté en 2007 (TRS 950) pour Juvenile idiopathic arthritis Ajouté en 2007 (TRS 950) pour Acute rheumatic fever without mention of heart involvement
<b>Wikipédia</b>	<a href="#">Acetylsalicylic acid</a> ↗
<b>DrugBank</b>	<a href="#">Acetylsalicylic acid</a> ↗

## Recommandations

### Section Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs)

Oral > Solid: 100 to 500 mg  
Local > Rectal > Suppository: 50 to 150 mg

### Indications

**Pain**

### Section Antimigraine medicines > For treatment of acute attack

Oral > Solid: 300 to 500 mg

### Indications

**Migraine**

Section      Anti-platelet medicines

Oral > Solid: 100 mg

Indications

Acute myocardial infarction

Cerebral ischaemic stroke

Section      Medicines for juvenile joint diseases

Oral > Solid: 100 to 500 mg

Local > Rectal > Suppository: 50 to 150 mg

Indications

Acute rheumatic fever without mention of heart involvement

Juvenile idiopathic arthritis

Mucocutaneous lymph node syndrome

