Ciprofloxacin

General description

Ciprofloxacin is a fluoroquinolone antibiotic

INN	Ciprofloxacin
ATC codes	S02AA15 J01MA02
Medicine type	Chemical agent
Antibiotic groups	W WATCH
EML status history	First added in 1991 (TRS 825) for Bacterial infection of unspecified site Changed in 2009 (TRS 958) for Infectious diseases of external ear Changed in 2017 (TRS 1006) for Infectious gastroenteritis or colitis without specification of infectious agent Changed in 2017 (TRS 1006) for Cholera Changed in 2017 (TRS 1006) for Neutropaenia Changed in 2017 (TRS 1006) for Peritoneal abscess Changed in 2017 (TRS 1006) for Peritonitis Changed in 2017 (TRS 1006) for Acute pyelonephritis Changed in 2017 (TRS 1006) for Inflammatory or other diseases of prostate Removed in 2017 (TRS 1006) for Bacterial infection of unspecified site Added in 2019 (TRS 1021) for Typhoid fever Changed in 2019 (TRS 1021) for Paratyphoid fever
Wikipedia	Ciprofloxacin
DrugBank	Ciprofloxacin

Recommendations

Section Ear, nose and throat medicines [c]

- Local > Otological > drops: 0.3% (as hydrochloride)

Indications

Infectious diseases of external ear

Section Watch group antibiotics

- Parenteral > General injections > IV: 2 mg per mL (as hyclate)
- Oral > Liquid: 250 mg per 5 mL (anhydrous)
- Oral > Solid: 250 mg (as hydrochloride)
- Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride)
- Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride)

Indications

First choice

Infectious gastroenteritis or colitis without specification of infectious agent

Neutropaenia (low-risk)

Acute pyelonephritis (mild to moderate)

Inflammatory or other diseases of prostate (mild to moderate)

Typhoid fever

Paratyphoid fever

Second choice

Cholera

Cholera [children]

Peritonitis (mild-moderate)

Peritoneal abscess (mild-moderate)

Other indications

Bacterial infection of unspecified site