




Description générale

Ciprofloxacin is a fluoroquinolone antibiotic

INN	Ciprofloxacin
Codes ATC	S02AA15 J01MA02
Type de médicament	Chemical agent
Groupes d'antibiotiques	 WATCH
Historique des statuts LME	<p>Ajouté pour la première fois en 1991 (TRS 825) pour Bacterial infection of unspecified site</p> <p>Ajouté en 2009 (TRS 958) pour Infectious diseases of external ear</p> <p>Ajouté en 2017 (TRS 1006) pour Infectious gastroenteritis or colitis without specification of infectious agent</p> <p>Ajouté en 2017 (TRS 1006) pour Cholera</p> <p>Ajouté en 2017 (TRS 1006) pour Neutropenia</p> <p>Ajouté en 2017 (TRS 1006) pour Peritoneal abscess</p> <p>Ajouté en 2017 (TRS 1006) pour Peritonitis</p> <p>Ajouté en 2017 (TRS 1006) pour Acute pyelonephritis</p> <p>Ajouté en 2017 (TRS 1006) pour Inflammatory and other diseases of prostate</p> <p>Retiré en 2017 (TRS 1006) pour Bacterial infection of unspecified site</p> <p>Ajouté en 2019 (TRS 1021) pour Typhoid fever</p> <p>Ajouté en 2019 (TRS 1021) pour Paratyphoid fever</p>
Wikipédia	Ciprofloxacin 
DrugBank	Ciprofloxacin 

Recommandations

Section Ear, nose and throat medicines [c]

Local > Otological > drops: 0.3% (as hydrochloride)

Indications

Infectious diseases of external ear

Section Watch group antibiotics

Parenteral > General injections > IV: 2 mg per mL (as hyclate)

Oral > Liquid: 250 mg per 5 mL (anhydrous)

Oral > Solid: 250 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride)

Indications

Premier choix

Typhoid fever

Paratyphoid fever

Inflammatory and other diseases of prostate (mild to moderate)

Acute pyelonephritis (mild to moderate)

Neutropenia (low-risk)

Infectious gastroenteritis or colitis without specification of infectious agent

Second choix

Cholera

Cholera [children]

co-prescrire avec [metronidazole](#)

Peritoneal abscess (mild-moderate)

co-prescrire avec [metronidazole](#)

Peritonitis (mild-moderate)

Autres indications

Bacterial infection of unspecified site

