Amoxicillin

Statut de médicament

Description générale

Amoxicillin is an extended-spectrum beta-lactam antibiotic.

INN	Amoxicillin
Codes ATC	J01CA04
Type de médicament	Chemical agent
Groupes d'antibiotiques	A ACCESS
Historique des statuts LME	Ajouté pour la première fois en 1989 (TRS 796) pour Bacterial infection of unspecified site Ajouté en 2017 (TRS 1006) pour Acute otitis media Ajouté en 2017 (TRS 1006) pour Acute pharyngitis Ajouté en 2017 (TRS 1006) pour Infectious cystitis Retiré en 2017 (TRS 1006) pour Bacterial infection of unspecified site Ajouté en 2017 (TRS 1006) pour Bacterial meningitis Ajouté en 2017 (TRS 1006) pour Acute malnutrition in infants, children or adolescents Ajouté en 2017 (TRS 1006) pour Chronic obstructive pulmonary disease with acute exacerbation Ajouté en 2017 (TRS 1006) pour Acute sinusitis Ajouté en 2017 (TRS 1006) pour Bacterial pneumonia Ajouté en 2019 (TRS 1021) pour Periapical abscess without sinus Retiré en 2021 (TRS 1035) pour Infectious cystitis Ajouté en 2023 (TRS 1049) pour Sepsis without septic shock Ajouté en 2025 (TRS 1064) pour Bacterial meningitis
Wikipédia	Amoxicillin
DrugBank	Amoxicillin

Recommandations

- Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for ora liquid
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)
- Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid
- Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (a sodium) powder for injection; 1 g in vial (as sodium) powder for injection
- Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate); 1 g (as trihydrate)

Indications

Premier choix

Acute otitis media

Acute pharyngitis

Acute malnutrition in infants, children or adolescents (uncomplicated) [children]

Acute malnutrition in infants, children or adolescents (complicated) [children]

Chronic obstructive pulmonary disease with acute exacerbation

Acute sinusitis

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Periapical abscess without sinus

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Infectious cystitis

- Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid
- Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (a sodium) powder for injection; 1 g in vial (as sodium) powder for injection
- Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)

Indications

Second choix

Bacterial meningitis