

Amoxicillin

Statut de médicament essentiel

Description générale

Amoxicillin is an extended-spectrum beta-lactam antibiotic.

INN

Amoxicillin

Codes ATC

[J01CA04](#)

Type de médicament

Chemical agent

Groupes d'antibiotiques

[ACCESS](#)

Historique des statuts LME

Ajouté pour la première fois en 1989 ([TRS 796](#)) pour [Bacterial infection of unspecified site](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Acute otitis media](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Acute pharyngitis](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Infectious cystitis](#)

Retiré en 2017 ([TRS 1006](#)) pour [Bacterial infection of unspecified site](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Bacterial meningitis](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Acute malnutrition in infants, children or adolescents](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Chronic obstructive pulmonary disease with acute exacerbation](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Acute sinusitis](#)

Ajouté en 2017 ([TRS 1006](#)) pour [Bacterial pneumonia](#)

Ajouté en 2019 ([TRS 1021](#)) pour [Periapical abscess without sinus](#)

Retiré en 2021 ([TRS 1035](#)) pour [Infectious cystitis](#)

Ajouté en 2023 ([TRS 1049](#)) pour [Sepsis without septic shock](#)

Ajouté en 2025 ([TRS 1064](#)) pour [Bacterial meningitis](#)

Wikipédia

[Amoxicillin](#)

DrugBank

[Amoxicillin](#)

Recommandations

Section

Access group antibiotics

- Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)
- Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid
- Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection
- Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate); 1 g (as trihydrate)

Indications

Premier choix

[Acute otitis media](#) [Acute pharyngitis](#) [Acute malnutrition in infants, children or adolescents \(uncomplicated\) \[children\]](#) [Acute malnutrition in infants, children or adolescents \(complicated\) \[children\]](#) [Chronic obstructive pulmonary disease with acute exacerbation](#) [Acute sinusitis](#) [Bacterial pneumonia \(Community-acquired pneumonia - mild to moderate\)](#) [Bacterial pneumonia \(Community-acquired pneumonia - severe\) \[children\]](#) [Periapical abscess without sinus](#)

Second choix

[Bacterial meningitis](#)

Autres indications

[Bacterial infection of unspecified site](#) [Infectious cystitis](#)

Section

Medicines for bacterial central nervous system infections

- Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid
- Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection
- Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)
- Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)

Indications

Second choix

[Bacterial meningitis](#)

