



## Description générale

Amoxicillin is an extended spectrum penicillin antibiotic. Clavulanic acid is a beta-lactamase inhibitor.

|                            |   |
|----------------------------|---|
| INN                        | Amoxicillin + clavulanic acid   |
| Codes ATC                  | J01CR02   |
| Type de médicament         | Chemical agent  |
| Groupes d'antibiotiques    | <span>A ACCESS</span>   |
| Historique des statuts LME | Ajouté pour la première fois en 1997 (TRS 882) pour Bacterial infection of unspecified site<br>Ajouté en 2017 (TRS 1006) pour Acute otitis media<br>Ajouté en 2017 (TRS 1006) pour Infectious cystitis<br>Ajouté en 2017 (TRS 1006) pour Chronic obstructive pulmonary disease with acute exacerbation<br>Ajouté en 2017 (TRS 1006) pour Acute sinusitis<br>Ajouté en 2017 (TRS 1006) pour Bacterial cellulitis, erysipelas or lymphangitis<br>Ajouté en 2017 (TRS 1006) pour Osteomyelitis or osteitis<br>Ajouté en 2017 (TRS 1006) pour Bacterial infection of joint<br>Retiré en 2017 (TRS 1006) pour Bacterial infection of unspecified site<br>Ajouté en 2017 (TRS 1006) pour Neutropenia<br>Ajouté en 2017 (TRS 1006) pour Peritoneal abscess<br>Ajouté en 2017 (TRS 1006) pour Peritonitis<br>Ajouté en 2017 (TRS 1006) pour Bacterial pneumonia<br>Ajouté en 2017 (TRS 1006) pour Other specified pneumonia<br>Ajouté en 2019 (TRS 1021) pour Other specified prophylactic measures<br>Ajouté en 2019 (TRS 1021) pour Multi-drug resistant Mycobacterium tuberculosis |
| Wikipédia                  | Amoxicillin + clavulanic acid   |
| DrugBank                   | Amoxicillin , Clavulanic acid (Clavulanate)   |

## Recommandations

Section Access group antibiotics

Parenteral > General injections > IV: 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection  
Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)  
Oral > Liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt) / 5 mL powder for oral liquid; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid  
Oral > Solid > dispersible tablet: 200 mg (as trihydrate) + 28.5 mg (as potassium salt); 250 mg (as trihydrate) + 62.5 mg (as potassium salt)  
Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt)

Indications

Premier choix

Peritonitis (mild-moderate)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Other specified pneumonia (Hospital-acquired pneumonia)

Peritoneal abscess (mild-moderate)

Neutropenia (low-risk)

Infectious cystitis

Acute sinusitis

Bacterial cellulitis, erysipelas or lymphangitis

Chronic obstructive pulmonary disease with acute exacerbation

Second choix

Other specified prophylactic measures

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Acute otitis media

Osteomyelitis or osteitis

Bacterial infection of joint

co-prescrite avec [clarithromycin](#)

Bacterial pneumonia (Community-acquired pneumonia - severe)

Autres indications

Bacterial infection of unspecified site

Section      **Antituberculosis medicines**

Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid

Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)

**Indications**

Multi-drug resistant *Mycobacterium tuberculosis*

