Doxycycline

Description générale

INN	Doxycycline
Codes ATC	J01AA02
Type de médicament	Chemical agent
Groupes d'antibiotiques	A ACCESS
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) pour Bacterial infection of unspecified site Modifié en 1995 (TRS 867) pour Malaria due to Plasmodium falciparum Modifié en 1999 (TRS 895) pour Malaria due to Plasmodium falciparum Modifié en 2017 (TRS 1006) pour Chronic obstructive pulmonary disease with acute exacerbation Modifié en 2017 (TRS 1006) pour Cholera Modifié en 2017 (TRS 1006) pour Bacterial pneumonia Modifié en 2017 (TRS 1006) pour Chlamydia trachomatis Retiré en 2017 (TRS 1006) pour Bacterial infection of unspecified site Modifié en 2025 (TRS 1064) pour Malaria due to Plasmodium falciparum Ajouté en 2025 (TRS 1064) pour Malaria due to Plasmodium falciparum Retiré en 2025 (TRS 1064) pour Malaria due to Plasmodium falciparum
Wikipédia	Doxycycline
DrugBank	Doxycycline

Recommandations

Section Access group antibiotics

- Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium)
- Parenteral > General injections > unspecified: 100 mg in vial powder for injection
- Oral > Solid > dispersible tablet: 100 mg (as monohydrate)
- Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)
- Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous)
- Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate)
- Oral > Solid > dosage form: 100 mg (as hyclate)

Indications

Premier choix

Cholera | Chlamydia trachomatis

Second choix

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Chronic obstructive pulmonary disease with acute exacerbation | Cholera [children]

Autres indications

Bacterial infection of unspecified site

Section Antimalarial medicines > Medicines for curative treatment

- Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Indications

Malaria due to Plasmodium falciparum

Section Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 100 mg (hydrochloride or hyclate)

Indications

Malaria due to Plasmodium falciparum

Section Antimalarial medicines > Medicines for chemoprophylaxis in travellers

- Oral > Liquid: 50 mg per 5 mL (calcium); 25 mg per 5 mL (monohydrate) powder for oral liquid
- Oral > Solid > dispersible tablet: 100 mg (as monohydrate)
- Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)

Indications

Malaria due to Plasmodium falciparum