



## Description générale

INN	Vancomycin
Codes ATC	A07AA09 J01XA01
Type de médicament	Chemical agent
Groupes d'antibiotiques	WATCH
Historique des statuts LME	Ajouté pour la première fois en 1995 (TRS 867) pour <b>Methicillin resistant Staphylococcus aureus</b> Ajouté en 2017 (TRS 1006) pour <b>Neutropenia</b> Ajouté en 2017 (TRS 1006) pour <b>Intestinal infections due to Clostridioides difficile</b> Retiré en 2017 (TRS 1006) pour <b>Methicillin resistant Staphylococcus aureus</b> Ajouté en 2021 (TRS 1035) pour <b>Endophthalmitis</b> Ajouté en 2021 (TRS 1035) pour <b>Necrotising fasciitis</b>
Wikipédia	<a href="#">Vancomycin</a> ↗
DrugBank	<a href="#">Vancomycin</a> ↗

## Recommandations

Section Watch group antibiotics

Oral > Solid > capsule: 125 mg (as hydrochloride); 250 mg (as hydrochloride)  
Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection  
Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection  
Parenteral > Locoregional injections > Intravitreal: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection

Indications

Premier choix

Necrotising fasciitis

co-prescrite avec [ceftazidime](#)

Endophthalmitis

co-prescrite avec [ceftriaxone](#)

Endophthalmitis

Second choix

Neutropenia (high-risk)

Intestinal infections due to Clostridioides difficile

Autres indications

Methicillin resistant Staphylococcus aureus

