



## General description

Calcium folinate (also known as leucovorin, folic acid) is a folic acid derivative that acts as a modulating and rescue agent in cancer chemotherapy protocols.

<b>INN</b>	Calcium folinate
<b>ATC codes</b>	V03AF03
<b>Medicine type</b>	Chemical agent
<b>EML status history</b>	<p>First added in 1979 (TRS 641) for <b>Malignant neoplasms of colon</b>            Added in 1979 (TRS 641) for <b>Malignant neoplasm metastasis in large intestine</b>            Added in 1979 (TRS 641) for <b>Malignant trophoblastic neoplasms of placenta</b>            Added in 1979 (TRS 641) for <b>Burkitt lymphoma including Burkitt leukaemia</b>            Added in 1979 (TRS 641) for <b>Osteosarcoma of bone and articular cartilage of unspecified sites</b>            Added in 1979 (TRS 641) for <b>Malignant neoplasms of rectum</b>            Added in 1979 (TRS 641) for <b>Unspecified malignant neoplasms of ill-defined or unspecified sites</b>            Removed in 2015 (TRS 994) for <b>Unspecified malignant neoplasms of ill-defined or unspecified sites</b></p>
<b>Wikipedia</b>	<a href="#">Calcium folinate (leucovorin calcium)</a>
<b>DrugBank</b>	<a href="#">Calcium folinate (Leucovorin)</a>

## Recommendations

### Section Cytotoxic medicines

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Solid > tablet: 15 mg; 5 mg; 25 mg

Oral > Solid: 15 mg tablet

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule

### Indications

Malignant neoplasms of colon

Osteosarcoma of bone and articular cartilage of unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of rectum

Malignant trophoblastic neoplasms of placenta

Unspecified malignant neoplasms of ill-defined or unspecified sites

