



## Description générale

Lidocaine is used in the management of cardiac arrhythmias. Lidocaine is also a local anaesthetic used for surface anaesthesia of mucous membranes, infiltration anaesthesia, peripheral and sympathetic nerve block, dental anaesthesia, spinal anaesthesia, and intravenous regional anaesthesia.

INN	Lidocaine
Codes ATC	N01BB02 C01BB01
Type de médicament	Chemical agent
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) pour Local anaesthetics Ajouté en 1977 (TRS 615) pour Ventricular tachyarrhythmia
Wikipédia	<a href="#">Lidocaine</a> ↗
DrugBank	<a href="#">Lidocaine</a> ↗

## Recommandations

### Section [Local anaesthetics](#)

Parenteral > Locoregional injections > Spinal anaesthesia: 5% in 2 mL ampoule (hydrochloride) + 7.5% glucose solution  
Parenteral > Locoregional injections > Other: 1% in vial (hydrochloride); 2% in vial (hydrochloride)  
Local > Topical > unspecified: 2 to 4% (hydrochloride)

### Indications

[Local anaesthetics](#)

### Section [Antiarrhythmic medicines](#)

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)

### Indications

[Ventricular tachyarrhythmia](#)

