Liste modèle des médicaments essentiels

BCG vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against tuberculosis

Japanese encephalitis vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Medicines for COVID-19

Section	Indications
Medicines for COVID-19 Refer to WHO living guidelines	COVID-19

Abacavir

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 300 mg tablet (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abacavir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abemaciclib

Section	Indications
Targeted therapies	Therapeutic equivalent to palbociclib pour Other specified malignant neoplasms of breast

Abiraterone

Section	Indications
Hormones and antihormones Oral > Solid: 250 mg; 500 mg	Malignant neoplasms of prostate

Acamprosate calcium

Section	Indications
Medicines for alcohol use disorders Oral > Solid > tablet: 333 mg	Disorders due to use of alcohol

Acetazolamide

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Oral > Solid: 250 mg	Primary open-angle glaucoma

Acetylcysteine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule Oral > Liquid: 10%; 20%	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs Acute or subacute hepatic failure

Acetylsalicylic acid

Section	Indications
Antimigraine medicines > For treatment of acute attack Oral > Solid: 300 to 500 mg	Migraine
Anti-platelet medicines Oral > Solid: 100 mg	Acute myocardial infarction Cerebral ischaemic stroke
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Pain

Acetylsalicylic acid + atorvastatin + ramipril

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Aciclovir

Section	Indications
Antiherpes medicines Parenteral > General injections > IV: 250 mg in vial powder for injection (as sodium salt) Oral > Liquid: 200 mg per 5 mL Oral > Solid: 200 mg tablet	Zoster Varicella Herpes simplex infections
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 3% w/w	Herpes simplex keratitis

Aclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Activated charcoal

Section	Indications
Antidotes and other substances used in poisonings > Non-specific Oral > Other: powder	Unspecified injury, poisoning or certain other consequences of external causes

Adalimumab

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Axial spondyloarthritis Crohn disease site Rheumatoid arthritis
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Afatinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib pour Other specified malignant neoplasms of bronchus or lung

Albendazole

Section	Indications
Intestinal anthelminthics Oral > Solid > tablet: 400 mg (chewable, scored)	Echinococcosis Enterobiasis Trichuriasis Hookworm diseases Cysticercosis Ascariasis
Cysticidal medicines Oral > Solid > tablet: 400 mg (chewable, scored); 200 mg (chewable)	Other specified echinococcosis Cysticercosis of central nervous system
Antifilarials Oral > Solid > tablet: 400 mg (chewable, scored)	Lymphatic filariasis

Alcohol based hand rub

Section	Indications
Disinfectants Local > Topical > Solution: 80% v/v ethanol; 75% v/v isopropyl alcohol	Denatured alcohol

Alcuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 5 mg per mL in 2 mL ampoule (alcuronium chloride)	Muscle relaxants

Alfacalcidol

Section	Indications
Medicines for endocrine disorders Oral > Liquid: 2 µg per mL Oral > Solid > capsule: 0.25 µg; 1 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

All-trans retinoic acid

Section	Indications
Targeted therapies Oral > Solid: 10 mg capsule	Acute myeloid leukaemia with recurrent genetic abnormalities

Allopurinol

Section	Indications
Medicines used to treat gout Oral > Solid: 100 mg	Gout

Alteplase

Section	Indications
Thrombolytic medicines Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection	Cerebral ischaemic stroke

Amidotrizoate

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)	Amidotrizoate

Amikacin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis Therapeutic equivalent to gentamicin pour Infectious blepharitis
Antituberculosis medicines Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)	Multi-drug resistant Mycobacterium tuberculosis
Access group antibiotics Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate); 50 mg per mL in 2 mL vial (as sulfate) Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)	Premier choix Neutropenia (high-risk) co-prescrite avec cefotaxime Inflammatory and other diseases of prostate (severe) co-prescrite avec cefotaxime Acute pyelonephritis (severe) co-prescrite avec ceftriaxone Inflammatory and other diseases of prostate (severe) co-prescrite avec ceftriaxone Acute pyelonephritis (severe)

Amiloride

Section	Indications
Diuretics Oral > Solid: 5 mg (hydrochloride)	Oedema Ascites

Amiodarone

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride) Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)	Cardiac arrhythmia

Amitriptyline

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 25 mg; 10 mg; 75 mg	Palliative care
Medicines for depressive disorders Oral > Solid: 25 mg; 75 mg	Depressive disorders

Amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 5 mg (as maleate, mesylate or besylate)	Essential hypertension

Amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Amoxicillin

Section	Indications
Access group antibiotics Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate) Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate) Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate); 1 g (as trihydrate)	Premier choix Chronic obstructive pulmonary disease with acute exacerbation Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) Acute pharyngitis Acute otitis media Acute sinusitis Periapical abscess without sinus co-prescrite avec gentamicin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] Second choix Bacterial meningitis Autres indications Infectious cystitis Bacterial infection of unspecified site

Amoxicillin + clavulanic acid

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection Oral > Liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt) / 5 mL powder for oral liquid; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid Oral > Solid > dispersible tablet: 200 mg (as trihydrate) + 28.5 mg (as potassium salt); 250 mg (as trihydrate) + 62.5 mg (as potassium salt) Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt) Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	Premier choix Peritoneal abscess (mild-moderate) Neutropenia (low-risk) Other specified pneumonia (Hospital-acquired pneumonia) Infectious cystitis Chronic obstructive pulmonary disease with acute exacerbation Peritonitis (mild-moderate) Acute sinusitis Bacterial cellulitis, erysipelas or lymphangitis Bacterial pneumonia (Community-acquired pneumonia - severe) [children] Second choix Osteomyelitis or osteitis Acute otitis media Other specified prophylactic measures Bacterial infection of joint Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) co-prescrite avec clarithromycin Bacterial pneumonia (Community-acquired pneumonia - severe) Autres indications Bacterial infection of unspecified site
Antituberculosis medicines Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	Multi-drug resistant Mycobacterium tuberculosis

Amphotericin B

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	Blastomycosis Cryptococcosis Candidosis Coccidioidomycosis Histoplasmosis Sporotrichosis Aspergillosis
Antileishmaniasis medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	Visceral leishmaniasis

Ampicillin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 1 g in vial (as sodium salt) powder for injection Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral suspension Oral > Solid: 250 mg (anhydrous); 500 mg (anhydrous)	Second choix Bacterial meningitis Autres indications Bacterial infection of unspecified site

Anakinra

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

Anastrozole

Section	Indications
Hormones and antihormones Oral > Solid: 1 mg tablet	Other specified malignant neoplasms of breast Malignant neoplasms of breast

Anidulafungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin pour Systemic or invasive candidosis

Ansuvimab

Section	Indications
Medicines for Ebola virus disease Parenteral > General injections > IV: 400 mg powder for injection	Ebola virus disease

Anti-D immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IV: 250 µg in single-dose vial Parenteral > General injections > IM: 250 µg in single-dose vial	Rh incompatibility reaction Maternal care for red cell antibodies

Anti-rabies immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 150 IU per mL in vial Parenteral > Locoregional injections > Other: 150 IU per mL in vial	Rabies

Anti-rabies virus monoclonal antibodies

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > Locoregional injections > Intradermal: 40 IU per mL in 1.25 mL vial (human); 40 IU per mL in 2.5 mL vial (human); 100 IU per mL in 2.5 mL vial (human); 300 IU per mL in 10 mL vial (murine); 600 IU per mL in 1 mL vial (murine); 600 IU per mL in 2.5 mL vial (murine); 600 IU per mL in 5 mL vial (murine)	Rabies

Anti-tetanus immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 500 IU in vial	Tetanus

Antirabies hyperimmune serum

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > unspecified: 1000 IU in 5 mL ampoule	Rabies

Antivenom immunoglobulin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IV:	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Antivenom sera

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IV:	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Apixaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Aprepitant

Section	Indications
Antiemetic medicines Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid: 80 mg; 125 mg; 165 mg	Nausea or vomiting

Aripiprazole

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders

Arsenic trioxide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion	Acute myeloid leukaemia with recurrent genetic abnormalities

Artemether

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artemether + lumefantrine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution Oral > Solid: 50 mg tablet Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + mefloquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + pyronaridine tetraphosphate

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Ascorbic acid

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg	Scurvy

Asparaginase

Section	Indications
Cytotoxic medicines Powder for injection: 10 000 IU in vial. Parenteral > General injections > unspecified: 10000 IU in vial powder for injection	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified

Atazanavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atazanavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atenolol

Section	Indications
Antianginal medicines Oral > Solid: 50 mg; 100 mg	Angina pectoris
Antihypertensive medicines Oral > Solid: 50 mg; 100 mg	Essential hypertension Therapeutic equivalent to bisoprolol pour Essential hypertension

Atezolizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion	Other specified malignant neoplasms of bronchus or lung Therapeutic equivalent to pembrolizumab pour Other specified malignant neoplasms of bronchus or lung

Atoltivimab + maftivimab + odesivimab

Section	Indications
Medicines for Ebola virus disease Parenteral > General injections > IV: 241.7 mg + 241.7 mg + 241.7 mg in 14.5 mL vial	Ebola virus disease

Atorvastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg	Chronic ischaemic heart disease Atherosclerotic chronic arterial occlusive disease Cerebral ischaemic stroke

Atracurium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg per mL (besylate)	Muscle relaxants Therapeutic equivalent to vecuronium pour Muscle relaxants

Atropine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	Exposure to or harmful effects of undetermined intent of pesticides
Preoperative medication and sedation for short-term procedures Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	Parasympatholytics [anticholinergics and antimuscarinics] and spasmolytics
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour Cycloplegic drug
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 0.1% (sulfate); 0.5% (sulfate); 1% (sulfate)	Anterior uveitis

Axicabtagene ciloleucel

Section	Indications
Antineoplastics and supportive medicines Cell suspension for infusion: 2 × 10*6 anti-CD19 CAR-positive viable T-cells per kg of body weight.	Diffuse large B-cell lymphomas

Azacitidine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection	Myeloid leukaemia

Azathioprine

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg in vial powder for injection (as sodium salt); 100 mg in vial powder for injection (as sodium salt) Oral > Liquid: 10 mg per mL Oral > Solid > tablet: 25 mg; 50 mg (scored) Oral > Solid > tablet: 50 mg (scored)	Failure or rejection of transplanted organs or tissues Relapsing-remitting multiple sclerosis
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 50 mg (scored)	Rheumatoid arthritis, serology unspecified Psoriatic arthritis

Azithromycin

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 1.5% eye drops	Trachoma
Watch group antibiotics Oral > Liquid: 200 mg per 5 mL (anhydrous) powder for oral liquid Oral > Solid > dosage form: 250 mg (anhydrous); 500 mg (anhydrous)	Premier choix Chlamydia trachomatis Typhoid fever Yaws Paratyphoid fever Cholera Cholera [children] Trachoma co-prescrite avec ceftriaxone Gonococcal infection Second choix Infectious gastroenteritis or colitis without specification of infectious agent co-prescrite avec cefixime Gonococcal infection

Aztreonam

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified bacterial diseases

Barium sulfate

Section	Indications
Diagnostic agents > Radiocontrast media Oral > Liquid: Aqueous suspension	Barium sulfate with suspending agents

Beclometasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 50 µg per dose (dipropionate); 250 µg per dose (dipropionatae)	Therapeutic equivalent to budesonide pour Asthma Asthma

Beclometasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Bedaquiline

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg tablet; 20 mg tablet	Multi-drug resistant Mycobacterium tuberculosis

Bendamustine

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL	Chronic lymphocytic leukaemia or small lymphocytic lymphoma Follicular lymphoma

Benzathine benzylpenicillin

Section	Indications
Access group antibiotics Parenteral > General injections > IM: 1.2 million IU in vial (≈ 900 mg) powder for injection; 2.4 million IU in vial (≈ 1.8 g) powder for injection	Premier choix Syphilis Autres indications Bacterial infection of unspecified site

Benznidazole

Section	Indications
American trypanosomiasis Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg	Chagas disease

Benzoic acid + salicylic acid

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Cream: 6% + 3% Local > Topical > Ointment: 6% + 3%	Fungal infection of the skin

Benzoyl peroxide

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Cream: 5% Local > Topical > Lotion: 5%	Acne

Benzyl benzoate

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Lotion: 25%	Pthiriasis Pediculosis corporis Pediculosis capitis Scabies

Benzylpenicillin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 600 mg in vial powder for injection (=1 million IU as sodium or potassium salt); 3 g in vial powder for injection (=5 million IU as sodium or potassium salt) Parenteral > General injections > unspecified: 600 mg in vial powder for injection (= 1 million IU as sodium or potassium salt); 3 g in vial powder for injection (= 5 million IU as sodium or potassium salt)	Premier choix Syphilis Second choix Bacterial meningitis Autres indications Bacterial infection of unspecified site

Betamethasone

Section	Indications
Dermatological medicines > Anti-inflammatory and antipruritic medicines Local > Topical > Cream: 0.1% (as valerate) Local > Topical > Ointment: 0.1% (as valerate)	Pruritus due to skin disorder Psoriasis of unspecified type Dermatitis or eczema Lichen planus

Bevacizumab

Section	Indications
Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations Parenteral > Locoregional injections > Intravitreal: 25 mg per mL	Age related macular degeneration

Bicalutamide

Section	Indications
Hormones and antihormones Oral > Solid: 50 mg	Malignant neoplasms of prostate

Binimetinib

Section	Indications
Targeted therapies	Therapeutic equivalent to trametinib pour Other specified melanoma of skin

Biperiden

Section	Indications
Medicines for parkinsonism Parenteral > General injections > IV: 5 mg per mL in 1 mL ampoule (lactate) Oral > Solid: 2 mg tablet (hydrochloride)	Parkinson disease

Bisacodyl

Section	Indications
Laxatives	Therapeutic equivalent to senna pour Constipation

Bisoprolol

Section	Indications
Antianginal medicines Oral > Solid: 1.25 mg; 5 mg	Angina pectoris
Medicines used in heart failure Oral > Solid: 1.25 mg; 5 mg	Heart failure
Antihypertensive medicines Oral > Solid: 1.25 mg; 5 mg	Essential hypertension
Antiarrhythmic medicines Oral > Solid: 1.25 mg; 5 mg	Cardiac arrhythmia

Bleomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)	Germ cell tumour of testis Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Unspecified malignant neoplasms of ill-defined or unspecified sites

Bortezomib

Section	Indications
Targeted therapies Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection	Plasma cell myeloma

Bromocriptine

Section	Indications
Antiseizure medicines Oral > Solid: 2.5 mg; 5 mg	Parkinson disease
Medicines for disorders of the pituitary hormone system	Therapeutic equivalent to cabergoline pour Other specified benign neoplasm of endocrine glands

Budesonide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 100 µg per dose; 200 µg per dose	Asthma

Budesonide + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > dry powder: 100 µg + 6 µg per dose; 200 µg + 6 µg per dose	Asthma

Budesonide + salmeterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Bumetanide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure
Diuretics	Therapeutic equivalent to furosemide pour Anuria or oliguria Therapeutic equivalent to furosemide pour Oedema

Bupivacaine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 0.5% in 4 mL ampoule (hydrochloride) + 7.5% glucose solution Parenteral > General injections > unspecified: 0.25% in vial (hydrochloride); 0.5% in vial (hydrochloride)	Local anaesthetics

Buprenorphine

Section	Indications
Medicines for opioid use disorders	Therapeutic equivalent to methadone pour Opioid dependence

Bupropion

Section	Indications
Medicines for nicotine use disorders Oral > Solid: 150 mg sustained-release (hydrochloride)	Nicotine dependence

C1 esterase inhibitor

Section	Indications
Plasma-derived medicines Parenteral > General injections > IV: 500 IU in vial powder for injection	Hereditary angioedema

Cabergoline

Section	Indications
Medicines for disorders of the pituitary hormone system Oral > Solid > tablet: 0.5 mg; 1 mg	Other specified benign neoplasm of endocrine glands
Antiseizure medicines Oral > Solid > tablet: 0.5 mg	Parkinson disease

Calamine

Section	Indications
Dermatological medicines > Anti-inflammatory and antipruritic medicines Local > Topical > Lotion:	Pruritus

Calcipotriol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Cream: 50 µg per mL (0.005%) Local > Topical > Lotion: 50 µg per mL (0.005%) Local > Topical > Ointment: 50 µg per mL (0.005%)	Psoriasis of unspecified type

Calcitriol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to calcipotriol pour Psoriasis of unspecified type
Medicines for endocrine disorders Oral > Solid > capsule: 0.25 µg; 0.5 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

Calcium

Section	Indications
Vitamins and minerals Oral > Solid: 500 mg (elemental calcium)	Calcium deficiency

Calcium folinate (leucovorin calcium)

Section	Indications
Cytotoxic medicines Oral > Solid: 15 mg tablet Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Solid > tablet: 15 mg; 5 mg; 25 mg	Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasm metastasis in large intestine Malignant trophoblastic neoplasms of placenta Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of rectum Malignant neoplasms of colon

Calcium gluconate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Hypermagnesaemia
Vitamins and minerals Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Tetany due to acute calcium deficiency

Canagliflozin

Section	Indications
Oral hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Capecitabine

Section	Indications
Cytotoxic medicines Oral > Solid > tablet: 150 mg; 500 mg	Other specified malignant neoplasms of breast Malignant neoplasm metastasis in large intestine Malignant neoplasms of rectum Malignant neoplasms of colon

Capreomycin

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

Captopril

Section	Indications
Medicines used in heart failure Oral > Solid: 25 mg	Heart failure

Carbachol

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines	Therapeutic equivalent to pilocarpine pour Acute angle closure with pupillary block Therapeutic equivalent to pilocarpine pour Ocular hypertension Therapeutic equivalent to pilocarpine pour Primary open-angle glaucoma

Carbamazepine

Section	Indications
Medicines for bipolar disorders Oral > Solid > tablet: 100 mg (scored); 200 mg (scored); 400 mg	Bipolar or related disorders
Antiseizure medicines Oral > Liquid: 100 mg per 5 mL Oral > Solid > tablet: 100 mg (scored); 200 mg (scored); 100 mg (chewable); 200 mg (chewable); 400 mg (scored)	Epilepsy or seizures

Carbetocin

Section	Indications
Uterotonics Parenteral > General injections > IV: 100 µg per mL (heat stable)	Postpartum haemorrhage

Carbimazole

Section	Indications
Thyroid hormones and antithyroid medicines	Therapeutic equivalent to methimazole pour Thyrotoxicosis

Carboplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL	Other specified gliomas of brain Other specified carcinomas of ovary Retinoblastoma Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Malignant neoplasms of kidney, except renal pelvis Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of lip, oral cavity or pharynx Malignant neoplasms of cervix uteri Other specified malignant neoplasms of the ovary Germ cell tumour of testis

Carmellose

Section	Indications
Ophthalmological preparations	Therapeutic equivalent to hypromellose pour Keratoconjunctivitis sicca

Carvedilol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Caspofungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin pour Systemic or invasive candidosis

Cefalexin

Section	Indications
Access group antibiotics Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid Oral > Solid: 250 mg (as monohydrate) Oral > Solid: 250 mg (as monohydrate); 500 mg (as monohydrate) Oral > Solid > dispersible tablet: 125 mg; 250 mg	Premier choix Bacterial cellulitis, erysipelas or lymphangitis Second choix Acute pharyngitis Chronic obstructive pulmonary disease with acute exacerbation Autres indications Bacterial infection of unspecified site

Cefazolin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 1 g in vial (as sodium salt) powder for injection	Premier choix Other specified prophylactic measures co-prescrite avec metronidazole Other specified prophylactic measures Second choix Osteomyelitis or osteitis Bacterial infection of joint **Autres indications** Surgical site infection

Cefepime

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection; 2 g in vial (as hydrochloride) powder for injection	Other specified bacterial diseases

Cefiderocol

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 1 g in vial (as sulfate tosylate) powder for injection	Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Cefixime

Section	Indications
Watch group antibiotics Oral > Liquid: 100 mg per 5 mL powder for oral liquid Oral > Solid: 200 mg (as trihydrate); 400 mg (as trihydrate)	Second choix Infectious gastroenteritis or colitis without specification of infectious agent co-prescrite avec azithromycin Gonococcal infection

Cefotaxime

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)	Premier choix Other specified pneumonia (Hospital-acquired pneumonia) Bacterial meningitis Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec amikacin Inflammatory and other diseases of prostate (severe) co-prescrite avec amikacin Acute pyelonephritis (severe) co-prescrite avec clarithromycin Bacterial pneumonia (Community-acquired pneumonia - severe) co-prescrite avec metronidazole Peritonitis (severe) co-prescrite avec metronidazole Peritonitis (mild-moderate) co-prescrite avec metronidazole Peritoneal abscess (severe) co-prescrite avec metronidazole Peritoneal abscess (mild-moderate) Second choix Osteomyelitis or osteitis Bacterial infection of joint Inflammatory and other diseases of prostate (mild to moderate) Acute pyelonephritis (mild to moderate)

Ceftaroline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 400 mg in vial (as fosamil) powder for injection; 600 mg in vial (as fosamil) powder for injection	Other specified bacterial diseases

Ceftazidime

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial (as pentahydrate) powder for injection; 1 g in vial (as pentahydrate) powder for injection	Premier choix co-prescrite avec vancomycin Endophthalmitis **Autres indications** Pseudomonas aeruginosa resistant to other antibiotic

Ceftazidime + avibactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 2 g in vial + 0.5 g in vial powder for injection	Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Ceftolozane + tazobactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 1 g + 0.5 g powder for injection	Carbapenem-resistant Pseudomonas aeruginosa

Ceftriaxone

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt) Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt) Parenteral > Locoregional injections > Intravitreal: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt) Parenteral > General injections > unspecified: 2 g in vial; 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)	Premier choix Typhoid fever Bacterial pneumonia (Community-acquired pneumonia - severe) [children] Paratyphoid fever Other specified pneumonia (Hospital-acquired pneumonia) Bacterial meningitis co-prescrite avec amikacin Acute pyelonephritis (severe) co-prescrite avec amikacin Inflammatory and other diseases of prostate (severe) co-prescrite avec azithromycin Gonococcal infection co-prescrite avec clarithromycin Bacterial pneumonia (Community-acquired pneumonia - severe)

co-prescrite avec metronidazole

Necrotising fasciitis

co-prescrite avec metronidazole

Peritonitis (mild-moderate)

co-prescrite avec metronidazole

Peritonitis (severe)

co-prescrite avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrite avec metronidazole

Peritoneal abscess (severe)

co-prescrite avec vancomycin

Endophthalmitis

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Infectious gastroenteritis or colitis without specification of infectious agent

Inflammatory and other diseases of prostate (mild to moderate)

Acute pyelonephritis (mild to moderate)

Autres indications

Bacterial infection of unspecified site

Cefuroxime

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 250 mg in vial (as sodium salt) powder for injection; 750 mg in vial (as sodium salt) powder for injection; 1.5 g in vial (as sodium salt) powder for injection	Second choix Other specified prophylactic measures

Cemiplimab

Section	Indications
Immunomodulators	Therapeutic equivalent to pembrolizumab pour Other specified malignant neoplasms of bronchus or lung

Certolizumab pegol

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis Therapeutic equivalent to adalimumab pour Crohn disease site Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Cetirizine

Section	Indications
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to loratadine pour Allergic or hypersensitivity conditions of unspecified type

Chlorambucil

Section	Indications
Cytotoxic medicines Oral > Solid: 2 mg tablet	Unspecified malignant neoplasms of ill-defined or unspecified sites Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Chloramphenicol

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 1 g in vial powder for injection (sodium succinate) Parenteral > General injections > IM: 500 mg in 2 mL ampoule oily suspension for injection Oral > Liquid: 150 mg per 5 mL (as palmitate) Oral > Solid: 250 mg capsule	Second choix Bacterial meningitis Autres indications Bacterial infection of unspecified site

Chlorhexidine

Section	Indications
Antiseptics Local > Topical > Solution: 5% (digluconate) for dilution	Chlorhexidine

Chlorine base compound

Section	Indications
Disinfectants Local > Topical > Powder: 0.1% available chlorine for solution Local > Topical > Liquid: 0.1% available chlorine for solution Local > Topical > Solid: 0.1% available chlorine for solution	Disinfectant, not elsewhere classified

Chloroquine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) Oral > Solid: 150 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax Malaria due to Plasmodium ovale Malaria due to Plasmodium malariae
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate) Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)	Rheumatoid arthritis

Chlorothiazide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chloroxylenol

Section	Indications
Disinfectants Local > Topical > Solution: 4.8%	Phenol disinfectant, not elsewhere classified

Chlorpromazine

Section	Indications
Medicines for psychotic disorders Oral > Liquid: 25 mg per 5 mL (hydrochloride) Oral > Solid: 100 mg (hydrochloride) Oral > Solid: 100 mg (hydrochloride); 10 mg (hydrochloride); 25 mg (hydrochloride); 50 mg (hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (hydrochloride)	Therapeutic equivalent to haloperidol pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Chlortalidone

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chlortetracycline

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to tetracycline pour Other specified conjunctivitis Therapeutic equivalent to tetracycline pour Infectious keratitis Therapeutic equivalent to tetracycline pour Infectious blepharitis Therapeutic equivalent to tetracycline pour Trachoma

Cholera vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against cholera alone

Ciclesonide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Ciclosporin

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection Oral > Liquid: 100 mg per mL Oral > Solid > capsule: 25 mg	Failure or rejection of transplanted organs or tissues

Cimetidine

Section	Indications
Antiulcer medicines Oral > Solid: 200 mg Parenteral > General injections > unspecified: 200 mg in 2 mL ampoule	Peptic ulcer, site unspecified Gastro-oesophageal reflux disease

Ciprofloxacin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 2 mg per mL (as hyclate) Oral > Liquid: 250 mg per 5 mL (anhydrous) Oral > Solid: 250 mg (as hydrochloride) Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride) Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride)	Premier choix Neutropenia (low-risk) Inflammatory and other diseases of prostate (mild to moderate) Typhoid fever Paratyphoid fever Infectious gastroenteritis or colitis without specification of infectious agent Acute pyelonephritis (mild to moderate) Second choix Cholera Cholera [children] co-prescrite avec metronidazole Peritoneal abscess (mild-moderate) co-prescrite avec metronidazole Peritonitis (mild-moderate) Autres indications Bacterial infection of unspecified site

Cisplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Germ cell tumour of testis Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of lip, oral cavity or pharynx Malignant neoplasms of cervix uteri

Cladribine

Section	Indications
Medicines for multiple sclerosis Oral > Solid > tablet: 10 mg	Multiple sclerosis
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial	Langerhans cell histiocytosis

Clarithromycin

Section	Indications
Watch group antibiotics Oral > Liquid: 125 mg per 5 mL powder for oral liquid; 250 mg per 5 mL powder for oral liquid Parenteral > General injections > unspecified: 500 mg in vial powder for injection Oral > Solid > dosage form: 500 mg; 250 mg Oral > Solid > dosage form: 500 mg	Premier choix co-prescrite avec cefotaxime Bacterial pneumonia (Community-acquired pneumonia - severe) co-prescrite avec ceftriaxone Bacterial pneumonia (Community-acquired pneumonia - severe) Second choix Acute pharyngitis co-prescrite avec amoxicillin + clavulanic acid Bacterial pneumonia (Community-acquired pneumonia - severe) Autres indications Helicobacter pylori associated gastric ulcer

Clindamycin

Section	Indications
Access group antibiotics Oral > Liquid: 75 mg per 5 mL (as palmitate hydrochloride) Oral > Solid: 150 mg (as hydrochloride) capsule Parenteral > General injections > unspecified: 150 mg per mL (as phosphate) injection Parenteral > General injections > IV: 150 mg per mL (as phosphate); 600 mg per 4 mL (as phosphate); 900 mg per 6 mL (as phosphate) Oral > Liquid: 75 mg per 5 mL powder for oral liquid (as palmitate hydrochloride) Oral > Solid: 150 mg (as hydrochloride)	Premier choix co-prescrite avec piperacillin + tazobactam Necrotising fasciitis Second choix Osteomyelitis or osteitis Bacterial infection of joint Autres indications Bacterial infection of unspecified site

Clofazimine

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg; 50 mg	Multi-drug resistant Mycobacterium tuberculosis
Antileprosy medicines Oral > Solid: 100 mg; 50 mg	Leprosy

Clomifene

Section	Indications
Ovulation inducers Oral > Solid: 50 mg (citrate)	Female infertility without specification whether primary or secondary

Clomipramine

Section	Indications
Medicines for obsessive compulsive disorders Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)	Obsessive-compulsive disorder

Clonazepam

Section	Indications
Antiseizure medicines Oral > Solid: 500 µg	Generalised myoclonic seizure

Clopidogrel

Section	Indications
Anti-platelet medicines Oral > Solid: 75 mg; 300 mg	Acute ischaemic heart disease Presence of coronary angioplasty implant or graft

Clotrimazole

Section	Indications
Antifungal medicines Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet	Vulvovaginal candidosis

Cloxacillin

Section	Indications
Access group antibiotics Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid Oral > Solid: 500 mg (as sodium salt) capsule; 1 g (as sodium salt) capsule Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid; 250 mg per 5 mL (as sodium salt) powder for oral liquid Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 250 mg in vial (as sodium salt) powder for injection Oral > Solid > capsule: 500 mg (as sodium salt); 1 g (as sodium salt); 250 mg (as sodium salt)	Premier choix Bacterial infection of joint Osteomyelitis or osteitis Bacterial cellulitis, erysipelas or lymphangitis Autres indications Bacterial infection of unspecified site

Clozapine

Section	Indications
Medicines for psychotic disorders Oral > Solid: 25 mg; 50 mg; 100 mg; 200 mg	Schizophrenia or other primary psychotic disorders

Coagulation factor IX

Section	Indications
Coagulation factors Parenteral > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection	Coagulation defects, purpura or other haemorrhagic or related conditions Haemophilia B

Coagulation factor IX complex

Section	Indications
Coagulation factors	Therapeutic equivalent to coagulation factor IX pour Coagulation defects, purpura or other haemorrhagic or related conditions Therapeutic equivalent to coagulation factor IX pour Haemophilia B

Coagulation factor VIII

Section	Indications
Coagulation factors Parenteral > General injections > IV: 250 IU in vial powder for injection; 500 IU in vial powder for injection; 1000 IU in vial powder for injection	Haemophilia A

Coal tar

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Solution: 5%	Psoriasis of unspecified type

Cobicistat + elvitegravir + emtricitabine + tenofovir disoproxil fumarate

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobicistat + elvitegravir + emtricitabine+ tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 10 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobimetinib

Section	Indications
Targeted therapies	Therapeutic equivalent to trametinib pour Other specified melanoma of skin

Codeine

Section	Indications
Opioid analgesics Oral > Solid: 30 mg tablet (codeine phosphate)	Pain
Antitussives Oral > Solid: 10 mg tablet (codeine phosphate)	Cough
Medicines for diarrhoea Oral > Solid: 30 mg tablet (codeine phosphate)	Diarrhoea

Colchicine

Section	Indications
Medicines used to treat gout Oral > Solid: 0.5 mg	Gout

Colecalciferol

Section	Indications
Vitamins and minerals	Therapeutic equivalent to ergocalciferol pour Vitamin D deficiency Therapeutic equivalent to ergocalciferol pour Calcium deficiency

Colistin (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium) Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium) (equivalent to 34 mg colistin base activity)	Other specified bacterial diseases Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Compound sodium lactate solution

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: injectable solution	Other specified disorders of fluid, electrolyte or acid-base balance Hypovolaemic shock

Condoms

Section	Indications
Barrier methods -	Contact with health services for contraceptive management

Copper-containing intrauterine device

Section	Indications
Intrauterine devices -	Contact with health services for insertion of contraceptive device

Crizotinib

Section	Indications
Targeted therapies Oral > Solid: 200 mg; 250 mg Oral > Solid > capsule: 200 mg; 250 mg	Other specified malignant neoplasms of bronchus or lung Anaplastic large cell lymphoma, ALK-positive

Cryoprecipitate (not pathogen-reduced)

Section	Indications
Blood and blood components	Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour Haemophilia A Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour Coagulation defects, purpura or other haemorrhagic or related conditions Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour Von Willebrand disease Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour Obstetric haemorrhage

Cryoprecipitate (pathogen-reduced)

Section	Indications
Blood and blood components Injection: frozen liquid in bag or lyophilized powder in vial containing: > 50 IU Factor VIII > 100 IU vWF > 140 mg clottable fibrinogen per unit	Haemophilia A Coagulation defects, purpura or other haemorrhagic or related conditions Von Willebrand disease Obstetric haemorrhage

Cyanocobalamin

Section	Indications
Antianaemia medicines .	Anaemias or other erythrocyte disorders

Cyclopentolate

Section	Indications
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour Cycloplegic drug
Ophthalmological preparations > Mydriatics	Therapeutic equivalent to atropine pour Anterior uveitis

Cyclophosphamide

Section	Indications
Cytotoxic medicines Oral > Solid: 25 mg tablet Parenteral > General injections > unspecified: 500 mg in vial powder for injection Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection Oral > Solid > dosage form: 25 mg; 50 mg	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified malignant neoplasms of breast Malignant trophoblastic neoplasms of placenta Follicular lymphoma Malignant neoplasms of breast Chronic lymphocytic leukaemia or small lymphocytic lymphoma Rhabdomyosarcoma primary site Hodgkin lymphoma Diffuse large B-cell lymphomas Plasma cell myeloma Other specified gliomas of brain Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Cycloserine

Section	Indications
Antituberculosis medicines Oral > Solid: 250 mg; 125 mg	Multi-drug resistant Mycobacterium tuberculosis

Cytarabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 100 mg in vial powder for injection Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Acute myeloid leukaemia with recurrent genetic abnormalities Burkitt lymphoma including Burkitt leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis

Dabigatran

Section	Indications
Medicines affecting coagulation Oral > Solid: 110 mg; 150 mg	Venous thromboembolism Other specified atrial fibrillation

Dabrafenib

Section	Indications
Targeted therapies Oral > Solid: 50 mg; 75 mg	Other specified melanoma of skin

Dacarbazine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection	Hodgkin lymphoma

Daclatasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 30 mg (as hydrochloride); 60 mg (as hydrochloride)	Chronic hepatitis C

Daclatasvir + sofosbuvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 60 mg + 400 mg	Chronic hepatitis C

Dactinomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 µg in vial powder for injection	Malignant trophoblastic neoplasms of placenta Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site Ewing sarcoma of bone and articular cartilage of unspecified sites

Dalteparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin pour Acute ischaemic heart disease Therapeutic equivalent to enoxaparin pour Venous thromboembolism

Dapagliflozin

Section	Indications
Oral hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Dapsone

Section	Indications
Antileprosy medicines Oral > Solid: 100 mg tablet; 50 mg tablet; 25 mg tablet	Leprosy

Daptomycin

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 350 mg in vial powder for injection; 500 mg in vial powder for injection	Other specified bacterial diseases

Daratumumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL	Plasma cell myeloma

Darbepoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Darunavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dasabuvir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 250 mg tablet	Chronic hepatitis C

Dasatinib

Section	Indications
Targeted therapies Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet	Chronic myeloid leukaemia, not elsewhere classified

Daunorubicin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)	Acute myeloid leukaemia with recurrent genetic abnormalities Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia

Deferasirox

Section	Indications
Other medicines for haemoglobinopathies Oral > Solid > dispersible tablet: 100 mg; 125 mg; 250 mg; 400 mg; 500 mg Oral > Solid > tablet: 90 mg; 180 mg; 360 mg	Other specified sickle cell disorders or other haemoglobinopathies Thalassaemias Therapeutic equivalent to deferoxamine pour Thalassaemias

Deferiprone

Section	Indications
Other medicines for haemoglobinopathies	Therapeutic equivalent to deferasirox pour Other specified sickle cell disorders or other haemoglobinopathies Therapeutic equivalent to deferasirox pour Thalassaemias

Deferoxamine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)	Harmful effects of drugs, medicaments or biological substances, not elsewhere classified
Other medicines for haemoglobinopathies Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)	Other specified sickle cell disorders or other haemoglobinopathies Thalassaemias

Delafloxacin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 300 mg lyophilized powder for injection Oral > Solid: 450 mg	Methicillin resistant Staphylococcus aureus

Delamanid

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 25 mg Oral > Solid > tablet: 50 mg	Multi-drug resistant Mycobacterium tuberculosis

Dengue vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Desmopressin

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 4 µg per mL in 1 mL ampoule (acetate) Local > Nasal > Spray: 10 µg per dose (acetate)	Haemophilia A Von Willebrand disease

Dexamethasone

Section	Indications
Other medicines administered to the mother Parenteral > General injections > unspecified: 4 mg per mL (dexamethasone phosphate (as sodium))	Respiratory distress syndrome of the newborn, altered by maternal corticosteroid therapy
Hormones and antihormones Oral > Solid: 4 mg Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosophate salt) Oral > Liquid: 2 mg per 5 mL Oral > Solid: 2 mg; 4 mg	Plasma cell myeloma Lymphoid leukaemia, not elsewhere classified Burkitt lymphoma including Burkitt leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive
Antiallergics and medicines used in anaphylaxis Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)	Allergic or hypersensitivity conditions of unspecified type
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per 5 mL Oral > Solid: 2 mg; 4 mg Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)	Palliative care
Antiemetic medicines Oral > Liquid: 0.5 mg per 5 mL; 2 mg per 5 mL Oral > Solid: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)	Nausea or vomiting

Dextran 40

Section	Indications
Plasma substitutes .	Hypovolaemia

Dextran 70

Section	Indications
Plasma substitutes Parenteral > General injections > IV: 6% injectable solution	Hypovolaemia

Diaphragms

Section	Indications
Barrier methods -	Contact with health services for contraceptive management

Diazepam

Section	Indications
Antiseizure medicines	Therapeutic equivalent to lorazepam pour
Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system	Status epilepticus
Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube	Status epilepticus
Medicines for other common symptoms in palliative care	Palliative care
Oral > Liquid: 2 mg per 5 mL	
Oral > Solid: 5 mg; 10 mg	
Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system	
Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube	
Parenteral > General injections > unspecified: 5 mg per mL	
Medicines for anxiety disorders	Anxiety
Oral > Solid: 5 mg (scored); 2 mg (scored)	

Didanosine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified
Oral > Liquid: 100 mg buffered powder for oral solution; 167 mg buffered powder for oral solution; 250 mg buffered powder for oral solution	
Oral > Solid: 25 mg buffered chewable dispersible tablet; 50 mg buffered chewable dispersible tablet; 100 mg buffered chewable dispersible tablet; 150 mg buffered chewable dispersible tablet; 200 mg buffered chewable dispersible tablet; 125 mg unbuffered enteric coated capsule; 200 mg unbuffered enteric coated capsule; 250 mg unbuffered enteric coated capsule; 400 mg unbuffered enteric coated capsule	

Diethylcarbamazine

Section	Indications
Antifilarials Oral > Solid: 50 mg tablet (dihydrogen citrate); 100 mg tablet (dihydrogen citrate)	Lymphatic filariasis

Digitoxin

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule Oral > Solid: 50 µg; 100 µg	Heart failure

Digoxin

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid: 62.5 µg; 250 µg	Cardiac arrhythmia
Medicines used in heart failure Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg	Heart failure

Dihydroartemisinin + piperaquine phosphate

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Dihydroergocryptine mesylate

Section	Indications
Antiseizure medicines Oral > Solid:	Parkinson disease

Diloxanide

Section	Indications
Antiamoebic and antigiardiasis medicines Oral > Solid: 500 mg (furoate)	Amoebiasis

Dimercaprol

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Diphtheria antitoxin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IM: 10000 IU in vial; 20000 IU in vial	Need for immunization against diphtheria alone

Diphtheria vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against diphtheria alone

Diphtheria-pertussis-tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against diphtheria-tetanus-pertussis, combined

Diphtheria-tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against combinations of infectious diseases

Docetaxel

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL	Other specified malignant neoplasms of breast Malignant neoplasms of breast Malignant neoplasms of prostate Unspecified malignant neoplasms of ill-defined or unspecified sites

Docusate sodium

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 50 mg per 5 mL oral liquid Oral > Solid: 100 mg capsule	Palliative care

Dolasetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Dolutegravir

Section	Indications
Antiretrovirals > Integrase inhibitors Oral > Solid: 50 mg tablet; 10 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dolutegravir + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Donepezil

Section	Indications
Medicines for diseases of the nervous system Oral > Liquid: 1 mg per mL (hydrochloride) Oral > Solid > tablet: 5 mg (hydrochloride); 10 mg (hydrochloride); 5 mg (hydrochloride) (orodispersible); 10 mg (hydrochloride) (orodispersible)	Dementia due to Alzheimer disease

Dopamine

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)	Heart failure Cardiogenic shock

Doxorubicin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride) Parenteral > General injections > IV: 10 mg in vial (hydrochloride) powder for injection; 50 mg in vial (hydrochloride) powder for injection; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)	Unspecified malignant neoplasms of ill-defined or unspecified sites Rhabdomyosarcoma primary site Malignant neoplasms of breast Hodgkin lymphoma Diffuse large B-cell lymphomas Plasma cell myeloma Follicular lymphoma Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Osteosarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Anaplastic large cell lymphoma, ALK-negative Other specified malignant neoplasms of breast Anaplastic large cell lymphoma, ALK-positive Ewing sarcoma of bone and articular cartilage of unspecified sites Kaposi sarcoma of unspecified primary site

Doxorubicin (as pegylated liposomal)

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)	Kaposi sarcoma of unspecified primary site

Doxycycline

Section	Indications
Access group antibiotics Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous) Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate) Parenteral > General injections > unspecified: 100 mg in vial powder for injection Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium) Oral > Solid > dispersible tablet: 100 mg (as monohydrate) Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate) Oral > Solid > dosage form: 100 mg (as hyclate)	Premier choix Chlamydia trachomatis Cholera Second choix Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) Chronic obstructive pulmonary disease with acute exacerbation Cholera [children] Autres indications Bacterial infection of unspecified site
Antimalarial medicines > For curative treatment Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)	Malaria due to Plasmodium falciparum
Antimalarial medicines > For chemoprevention Oral > Solid: 100 mg (hydrochloride or hyclate)	Malaria due to Plasmodium falciparum

Durvalumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 120 mg per 2.4 mL in vial; 500 mg per 10 mL in vial	Other specified malignant neoplasms of bronchus or lung

Edoxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Efavirenz

Section	Indications
Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors Oral > Solid: 600 mg tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Efavirenz + emtricitabine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Efavirenz + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Therapeutic equivalent to efavirenz + emtricitabine + tenofovir pour Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Eflornithine

Section	Indications
Medicines for the treatment of 2nd stage African trypanosomiasis Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)	African trypanosomiasis

Elbasvir + grazoprevir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 50 mg + 100 mg	Chronic hepatitis C

Empagliflozin

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 10 mg; 25 mg	Type 2 diabetes mellitus

Emtricitabine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 10 mg per mL Oral > Solid: 200 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 25 mg + 25 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir disoproxil fumarate

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 25 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Contact with or exposure to human immunodeficiency virus

Emtricitabine + tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 10 mg; 200 mg + 25 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Enalapril

Section	Indications
Antihypertensive medicines Oral > Liquid: 1 mg per mL (as hydrogen maleate) Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Essential hypertension
Medicines used in heart failure Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Heart failure

Encorafenib

Section	Indications
Targeted therapies	Therapeutic equivalent to dabrafenib pour Other specified melanoma of skin

Enoxaparin

Section	Indications
Medicines affecting coagulation Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule; 120 mg per 0.8 mL in ampoule	Acute ischaemic heart disease Venous thromboembolism

Entecavir

Section	Indications
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 0.05 mg per mL Oral > Solid: 0.5 mg; 1 mg	Chronic hepatitis B

Enzalutamide

Section	Indications
Hormones and antihormones Oral > Solid: 40 mg	Therapeutic equivalent to abiraterone pour Malignant neoplasms of prostate Malignant neoplasms of prostate

Ephedrine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 30 mg per mL in 1 mL ampoule (ephedrine hydrochloride)	Cardiac complications of anaesthesia during labour or delivery

Epinephrine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)	Cardiac arrest
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 2% (as hydrochloride)	Primary open-angle glaucoma
Antiasthmatic and medicines for chronic obstructive pulmonary disease Parenteral > General injections > unspecified: 1 mg per mL in ampoule (as hydrochloride or hydrogen tartrate)	Asthma Chronic obstructive pulmonary disease
Antiallergics and medicines used in anaphylaxis Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (as hydrochloride or hydrogen tartrate)	Anaphylaxis

Epoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Epoetin beta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Epoetin theta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Equine rabies immunoglobulin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > Locoregional injections > Intradermal: 150 IU per mL in vial; 200 IU per mL in vial; 300 IU per mL in vial; 400 IU per mL in vial	Rabies

Eravacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg lyophilized powder for injection	Carbapenem resistant Enterobacterales

Ergocalciferol

Section	Indications
Vitamins and minerals Oral > Liquid: 250 µg per mL (10 000 IU per mL) Oral > Solid: 1.25 mg (50 000 IU)	Vitamin D deficiency Calcium deficiency

Ergometrine

Section	Indications
Uterotonics Parenteral > General injections > unspecified: 200 µg in 1 mL ampoule (hydrogen maleate)	Postpartum haemorrhage

Erlotinib

Section	Indications
Targeted therapies Oral > Solid: 100 mg; 150 mg	Other specified malignant neoplasms of bronchus or lung

Erythromycin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 500 mg in vial powder for injection (as lactobionate) Oral > Liquid: 125 mg per 5 mL (as stearate or ethylsuccinate) Oral > Solid: 250 mg (as stearate or ethylsuccinate)	Second choix Therapeutic equivalent to clarithromycin pour Acute pharyngitis **Autres indications** Bacterial infection of unspecified site

Erythropoiesis-stimulating agents

Section	Indications
Antianaemia medicines Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe	Anaemia due to chronic disease

Estradiol

Section	Indications
Estrogens Oral > Solid > tablet: 0.5 mg; 1 mg; 2 mg	Gonadotropin deficiency Hypopituitarism Turner syndrome

Estradiol cypionate + medroxyprogesterone acetate

Section	Indications
Injectable hormonal contraceptives Parenteral > General injections > IM: 5 mg + 25 mg	Contact with health services for reasons associated with reproduction

Etanercept

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis Therapeutic equivalent to adalimumab pour Crohn disease site Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Ethambutol

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg per 10 mL; 2000 mg per 20 mL Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 100 mg (hydrochloride); 400 mg (hydrochloride)	Other specified tuberculosis Tuberculosis

Ethambutol + isoniazid

Section	Indications
Antituberculosis medicines Oral > Solid: 400 mg + 150 mg tablet	Tuberculosis

Ethambutol + isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet	Tuberculosis

Ethambutol + isoniazid + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid: 275 mg + 75 mg + 150 mg tablet	Tuberculosis

Ethanol

Section	Indications
Antiseptics Local > Topical > Solution: 70% (denatured)	Denatured alcohol

Ether

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > solution:	Anaesthetics and therapeutic gases

Ethinylestradiol + etonogestrel

Section	Indications
Intravaginal contraceptives Local > Vaginal > vaginal ring: 2.7 mg + 11.7 mg	Contact with health services for contraceptive management

Ethinylestradiol + levonorgestrel

Section	Indications
Oral hormonal contraceptives Oral > Solid: 50 µg + 250 µg [4] Oral > Solid: 30 µg + 150 µg	Contact with health services for postcoital contraception Contact with health services for contraceptive management

Ethinylestradiol + norethisterone

Section	Indications
Oral hormonal contraceptives Oral > Solid: 35 µg + 1 mg	Contact with health services for contraceptive management

Ethionamide

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 125 mg Oral > Solid > tablet: 250 mg	Multi-drug resistant Mycobacterium tuberculosis Tuberculous meningitis

Ethosuximide

Section	Indications
Antiseizure medicines Oral > Liquid: 250 mg per 5 mL Oral > Solid: 250 mg	Absence seizures, typical

Etonogestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 68 mg single rod	Contact with health services for insertion of contraceptive device

Etoposide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule Oral > Solid: 100 mg capsule Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate) Oral > Solid: 100 mg capsule; 50 mg capsule	Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Germ cell tumour of testis Malignant trophoblastic neoplasms of placenta Malignant neoplasms of kidney, except renal pelvis Osteosarcoma of bone and articular cartilage of unspecified sites Hodgkin lymphoma Retinoblastoma Anaplastic large cell lymphoma, ALK-positive Anaplastic large cell lymphoma, ALK-negative

Everolimus

Section	Indications
Targeted therapies Oral > Solid > dispersible tablet: 2 mg; 3 mg; 5 mg Oral > Solid > tablet: 2.5 mg; 5 mg; 7.5 mg; 10 mg	Other specified gliomas of brain

Fentanyl

Section	Indications
Opioid analgesics Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate) Oral > Other: 200 to 1200 µg buccal film (as citrate)	Chronic cancer pain Other specified chronic cancer related pain

Ferrous salt

Section	Indications
Antianaemia medicines Oral > Liquid: 25 mg per mL iron (as sulfate) Oral > Solid: 60 mg iron	Iron deficiency anaemia

Ferrous salt + folic acid

Section	Indications
Antianaemia medicines Oral > Solid: 60 mg iron + 2.8 mg Oral > Solid > tablet: 60 mg elemental iron + 400 µg tablet; 60 mg elemental iron + 2.8 mg	Iron deficiency anaemia Clefts of lip, alveolus or palate Iron deficiency anaemia Anencephaly or similar anomalies Spina bifida

Fexinidazole

Section	Indications
Medicines for the treatment of 1st stage African trypanosomiasis Oral > Solid: 600 mg	African trypanosomiasis
Medicines for the treatment of 2nd stage African trypanosomiasis Oral > Solid: 600 mg	African trypanosomiasis

Fexofenadine

Section	Indications
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to loratadine pour Allergic or hypersensitivity conditions of unspecified type

Fifth generation cephalosporins

Section	Indications
Reserve group antibiotics .	Other specified bacterial diseases

Filgrastim

Section	Indications
Immunomodulators Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial	Acquired neutropaenia

Fingolimod

Section	Indications
Immunomodulators for non-malignant disease Oral > Solid: 0.25 mg; 0.5 mg	Relapsing-remitting multiple sclerosis

Flomoxef

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 0.5 g in vial powder for injection; 1 g in vial powder for injection	Other specified other antibiotic resistant Enterobacterales

Fluconazole

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 2 mg per mL in vial Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid Oral > Solid > capsule: 50 mg	Cryptococcosis Candidosis

Flucytosine

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 2.5 g per 250 mL infusion Oral > Solid: 250 mg capsule	Cryptococcosis

Fludarabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection Oral > Solid: 10 mg tablet (fludarabine phosphate)	Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Fludrocortisone

Section	Indications
Adrenal hormones and synthetic substitutes Oral > Solid: 100 µg (acetate)	Adrenocortical insufficiency

Flunisolide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Fluorescein

Section	Indications
Diagnostic agents > Ophthalmic medicines Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)	Fluorescein

Fluoride

Section	Indications
Dental medicines and preparations Local > Dental > Gel: 2500 to 12500 ppm fluoride (any type) Local > Dental > Toothpaste, cream or gel: 1000 to 1500 ppm fluoride (any type) Local > Dental > Mouthrinse: 230 to 900 ppm fluoride (any type) Local > Dental > Varnish: 22500 ppm fluoride (any type)	Dental caries

Fluorouracil

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Ointment: 5%	Plantar warts
Cytotoxic medicines Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule Parenteral > General injections > IV: 50 mg per mL in vial	Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of nasopharynx Malignant neoplasm metastasis in large intestine Malignant neoplasms of breast Malignant neoplasms of rectum Malignant neoplasms of colon

Fluoxetine

Section	Indications
Medicines for depressive disorders Oral > Solid: 20 mg (as hydrochloride)	Depressive disorders
Medicines for other common symptoms in palliative care Oral > Solid: 20 mg (as hydrochloride)	Palliative care
Medicines for anxiety disorders Oral > Solid: 20 mg (as hydrochloride)	Social anxiety disorder Generalised anxiety disorder Panic disorder
Medicines for obsessive compulsive disorders Oral > Solid: 20 mg (as hydrochloride)	Obsessive-compulsive disorder

Fluphenazine

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 25 mg per mL in ampoule (decanoate); 25 mg per mL in ampoule (enanthate)	Schizophrenia or other primary psychotic disorders

Flutamide

Section	Indications
Hormones and antihormones	Therapeutic equivalent to bicalutamide pour Malignant neoplasms of prostate

Fluticasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Fluticasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Fluticasone furoate + vilanterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Fluvastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Folic acid

Section	Indications
Antianaemia medicines Oral > Solid: 1 mg tablet; 5 mg tablet Oral > Solid: 400 µg	Folate deficiency anaemia Anencephaly or similar anomalies Spina bifida Cephalocele

Fomepizole

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base)	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fosfomycin (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 2 g in vial (as sodium) powder for injection; 4 g in vial (as sodium) powder for injection	Other specified bacterial diseases Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Fourth generation cephalosporins

Section	Indications
Watch group antibiotics .	Other specified bacterial diseases

Fresh-frozen plasma

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Fulvestrant

Section	Indications
Hormones and antihormones Parenteral > General injections > IV: 250 mg per 5 mL	Malignant neoplasms of breast

Furosemide

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL Oral > Solid > tablet: 40 mg; 20 mg	Heart failure
Diuretics Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL Oral > Solid > tablet: 40 mg; 20 mg	Anuria or oliguria Oedema

Gabapentin

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg	Neuropathic pain

Gallamine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 40 mg per mL in 2 mL ampoule (gallamine triethiodide)	Muscle relaxants

Gatifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis

Gefitinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib pour Other specified malignant neoplasms of bronchus or lung

Gemcitabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection	Other specified carcinomas of ovary Other specified malignant neoplasms of bronchus or lung

Gentamicin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)	Premier choix co-prescrite avec amoxicillin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec ampicillin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec ampicillin Acute malnutrition in infants, children or adolescents (complicated) [children] co-prescrite avec benzylpenicillin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec benzylpenicillin Acute malnutrition in infants, children or adolescents (complicated) [children] Second choix Other specified prophylactic measures Gonococcal infection Autres indications Bacterial infection of unspecified site
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.3% (gentamicin sulfate)	Other specified conjunctivitis Infectious blepharitis

Glass ionomer cement

Section	Indications
Dental medicines and preparations Local > Dental > Powder + liquid: Single-use capsules: 0.4 g powder + 0.09 mL liquid; Multi-use bottle: powder + liquid	Dental caries

Glatiramer acetate

Section	Indications
Medicines for multiple sclerosis Parenteral > General injections > SC: 20 mg per mL; 40 mg per mL Parenteral > General injections > SC: 20 mg per mL in pre-filled syringe; 40 mg per mL in pre-filled syringe	Relapsing-remitting multiple sclerosis Multiple sclerosis

Glecaprevir + pibrentasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid: 50 mg + 20 mg granules Oral > Solid > tablet: 100 mg + 40 mg	Chronic hepatitis C

Glibenclamide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 5 mg; 2.5 mg	Type 2 diabetes mellitus

Gliclazide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release	Type 2 diabetes mellitus

Glucagon

Section	Indications
Medicines for hypoglycaemia Parenteral > General injections > unspecified: 1 mg per mL	Hypoglycaemia in the context of diabetes

Glucose

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 5% isotonic; 10% hypertonic Parenteral > General injections > IV: 50% hypertonic	Other specified disorders of fluid, electrolyte or acid-base balance Hypoglycaemia without associated diabetes

Glucose + sodium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 4% + 0.18% (equivalent to Na+ 30 mmol/L and Cl- 30 mmol/L; 5% + 0.9% (equivalent to Na+ 150 mmol/L and Cl- 150 mmol/L); 5% + 0.45% (equivalent to Na+ 75 mmol/L and Cl- 75 mmol/L)	Other specified disorders of fluid, electrolyte or acid-base balance

Glutaral

Section	Indications
Disinfectants Local > Topical > Solution: 2%	Glutaral nonmedicinal

Glyceryl trinitrate

Section	Indications
Antianginal medicines Oral > Solid: 500 µg sublingual tablet	Angina pectoris

Glycopyrronium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Golimumab

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis Therapeutic equivalent to adalimumab pour Crohn disease site Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Goserelin

Section	Indications
Hormones and antihormones	Therapeutic equivalent to leuprorelin pour Malignant neoplasms of breast Therapeutic equivalent to leuprorelin pour Malignant neoplasms of prostate

Granisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Griseofulvin

Section	Indications
Antifungal medicines Oral > Liquid: 125 mg per 5 mL Oral > Solid: 125 mg; 250 mg	Dermatophytosis

Haemophilus influenzae type b vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Haloperidol

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per mL Oral > Solid: 0.5 mg; 2 mg; 5 mg Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule	Palliative care
Medicines for psychotic disorders Parenteral > General injections > IM: 5 mg per mL in ampoule Oral > Solid: 2 mg; 5 mg	Schizophrenia or other primary psychotic disorders

Haloperidol decanoate

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to fluphenazine pour Schizophrenia or other primary psychotic disorders

Halothane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > liquid:	Anaesthetics and therapeutic gases

Heparin sodium

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 1000 IU per mL in 1 mL ampoule; 20000 IU per mL in 1 mL ampoule; 5000 IU per mL in 1 mL ampoule	Acute ischaemic heart disease Unstable angina Venous thromboembolism

Hepatitis A vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against viral hepatitis

Hepatitis B vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against viral hepatitis

Homatropine

Section	Indications
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 2% (hydrobromide)	Anterior uveitis Therapeutic equivalent to atropine pour Anterior uveitis

Human papilloma virus (HPV) vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Hydralazine

Section	Indications
Antihypertensive medicines Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride) Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride) Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)	Essential hypertension Gestational hypertension

Hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Liquid: 50 mg per 5 mL Oral > Solid: 25 mg; 12.5 mg	Essential hypertension
Medicines used in heart failure Oral > Liquid: 50 mg per 5 mL Oral > Solid: 25 mg	Heart failure
Diuretics Oral > Solid: 25 mg	Oedema

Hydrocortisone

Section	Indications
Hormones and antihormones Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)	Lymphoid leukaemia, not elsewhere classified Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia
Adrenal hormones and synthetic substitutes Oral > Solid: 5 mg; 10 mg; 20 mg	Adrenocortical insufficiency
Ophthalmological preparations > Anti-inflammatory agents Local > Ophthalmological > Ointment: 1% (acetate)	Scleritis Anterior uveitis
Antiallergics and medicines used in anaphylaxis Parenteral > General injections > unspecified: 100 mg in vial powder for injection (as sodium succinate)	Anaphylaxis
Anti-inflammatory medicines Local > Rectal > Suppository: 25 mg (acetate) Local > Rectal > Retention enema: 100 mg per 60 mL	Crohn disease site Ulcerative colitis
Dermatological medicines > Anti-inflammatory and antipruritic medicines Local > Topical > Cream: 1% (acetate) Local > Topical > Ointment: 1% (acetate)	Dermatitis or eczema Lichen planus Pruritus due to skin disorder Psoriasis of unspecified type

Hydromorphone

Section	Indications
Opioid analgesics	Therapeutic equivalent to morphine pour Pain

Hydroxocobalamin

Section	Indications
Antianaemia medicines Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)	Megaloblastic anaemia due to vitamin B12 deficiency

Hydroxycarbamide (hydroxyurea)

Section	Indications
Cytotoxic medicines Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg	Squamous cell carcinoma of oropharynx Chronic myeloid leukaemia, not elsewhere classified
Other medicines for haemoglobinopathies Oral > Solid > dosage form: 200 mg; 500 mg; 1 g; 100 mg	Other specified sickle cell disorders or other haemoglobinopathies

Hydroxychloroquine

Section	Indications
Antimalarial medicines > For chemoprevention	Therapeutic equivalent to chloroquine pour Malaria due to Plasmodium falciparum
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 200 mg (as sulfate)	Systemic lupus erythematosus Cutaneous lupus erythematosus of unspecified type

Hyoscine butylbromide

Section	Indications
Medicines for other common symptoms in palliative care Parenteral > General injections > unspecified: 20 mg per mL injection	Palliative care

Hypochlorous acid

Section	Indications
Antiseptics Local > Topical > Solution:	Disinfectant, not elsewhere classified

Hypromellose

Section	Indications
Ophthalmological preparations Local > Ophthalmological > Solution (eye drops): 0.3%	Keratoconjunctivitis sicca

Ibrutinib

Section	Indications
Targeted therapies Oral > Solid: 140 mg	Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Ibuprofen

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL Oral > Solid > tablet: 200 mg; 400 mg; 600 mg	Pain

Idoxuridine

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.1% Local > Ophthalmological > Ointment: 0.2%	Herpes simplex keratitis

Ifosfamide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified malignant neoplasms of the ovary Germ cell tumour of testis Ewing sarcoma of bone and articular cartilage of unspecified sites Osteosarcoma of bone and articular cartilage of unspecified sites Rhabdomyosarcoma primary site Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Imatinib

Section	Indications
Targeted therapies Oral > Solid: 100 mg; 400 mg	B lymphoblastic leukaemia or lymphoma with t(9:22) (q34;q11.2); BCR-ABL1 Gastrointestinal stromal tumour of unspecified gastrointestinal sites Chronic myeloid leukaemia, not elsewhere classified

Imipenem + cilastatin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 250 mg (as monohydrate) + 250 mg (as sodium salt) powder for injection; 500 mg (as monohydrate) + 500 mg (as sodium salt) powder for injection	Second choix Therapeutic equivalent to meropenem pour Neutropenia (high-risk) Therapeutic equivalent to meropenem pour Peritoneal abscess (severe) Therapeutic equivalent to meropenem pour Peritonitis (severe) **Autres indications** Pseudomonas aeruginosa resistant to other antibiotic Antibiotic resistant Acinetobacter baumannii
Antituberculosis medicines	Therapeutic equivalent to meropenem pour Multi-drug resistant Mycobacterium tuberculosis

Imipenem + cilastatin + relebactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500 mg + 500 mg + 250 mg powder for injection	Other specified other antibiotic resistant Enterobacterales Carbapenem resistant Enterobacterales Carbapenem-resistant Pseudomonas aeruginosa

Indapamide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Indinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 400 mg (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Infliximab

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis Therapeutic equivalent to adalimumab pour Crohn disease site Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Influenza vaccine (seasonal)

Section	Indications
Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against influenza

Insulin

Section	Indications
Insulins Parenteral > General injections > SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen (soluble)	Type 2 diabetes mellitus Type 1 diabetes mellitus

Insulin analogues

Section	Indications
Insulins Not specified	Type 1 diabetes mellitus Type 2 diabetes mellitus

Insulin degludec

Section	Indications
Insulins Parenteral > General injections > SC: 100 units per mL	Therapeutic equivalent to insulin glargine pour Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 2 diabetes mellitus

Insulin detemir

Section	Indications
Insulins	Therapeutic equivalent to insulin glargine pour Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 2 diabetes mellitus

Insulin glargine

Section	Indications
Insulins Parenteral > General injections > SC: 100 units per mL	Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues pour Type 2 diabetes mellitus

Intermediate-acting insulin

Section	Indications
Insulins Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin	Type 1 diabetes mellitus Type 2 diabetes mellitus

Intraperitoneal dialysis solution

Section	Indications
Peritoneal dialysis solution parenteral solution	Care involving peritoneal dialysis

Iodine

Section	Indications
Vitamins and minerals Parenteral > General injections > IM: 0.5 mL (240 mg iodine) iodized oil in ampoule; 1 mL (480 mg iodine) iodized oil in ampoule Oral > Liquid: 1 mL (480 mg iodine) iodized oil in ampoule; 0.5 mL (240 mg iodine) iodized oil in ampoule; 0.57 mL (308 mg iodine) iodized oil in dispenser bottle Oral > Solid: 190 mg	Iodine deficiency
Antiseptics	Therapeutic equivalent to povidone iodine pour Povidone iodine

Iohexol

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule	X-ray contrast media, iodinated

Ipecacuanha

Section	Indications
Antidotes and other substances used in poisonings > Non-specific Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine	Unspecified injury, poisoning or certain other consequences of external causes

Ipratropium bromide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 20 µg per dose	Asthma Chronic obstructive pulmonary disease

Irinotecan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site

Isoflurane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > liquid:	Anaesthetics and therapeutic gases

Isoniazid

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 100 mg per mL Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 100 mg; 300 mg	Other specified tuberculosis Tuberculosis

Isoniazid + pyridoxine + sulfamethoxazole + trimethoprim

Section	Indications
Medicines for prevention of HIV-related opportunistic infections Oral > Solid: 300 mg + 25 mg + 800 mg + 160 mg tablet (scored)	Other specified prophylactic measures

Isoniazid + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 50 mg + 75 mg Oral > Solid > tablet: 75 mg + 150 mg; 150 mg + 300 mg	Tuberculosis

Isoniazid + rifapentine

Section	Indications
Antituberculosis medicines Oral > Solid: 300 mg + 300 mg tablet (scored)	Latent tuberculosis

Isoprenaline

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride) Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)	Bradycardia Cardiogenic shock

Isosorbide dinitrate

Section	Indications
Antianginal medicines Oral > Solid: 5 mg sublingual tablet	Angina pectoris

Itraconazole

Section	Indications
Antifungal medicines Oral > Liquid: 10 mg per mL Oral > Solid: 100 mg	Chronic pulmonary aspergillosis Chromoblastomycosis Histoplasmosis Talaromycosis Sporotrichosis Paracoccidioidomycosis

Ivermectin

Section	Indications
Intestinal anthelminthics Oral > Solid > tablet: 3 mg	Strongyloidiasis Ascariasis Trichuriasis Hookworm diseases Ancylostomiasis
Antifilarials Oral > Solid > tablet: 3 mg	Onchocerciasis Lymphatic filariasis
Medicines for ectoparasitic infections Oral > Solid > tablet: 3 mg	Scabies

Kanamycin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis Therapeutic equivalent to gentamicin pour Infectious blepharitis

Kanamycin (injection)

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

Ketamine

Section	Indications
General anaesthetics and oxygen > Injectable medicines Parenteral > General injections > unspecified: 50 mg per mL in 10 mL vial (as hydrochloride)	Anaesthetics and therapeutic gases

Ketoconazole

Section	Indications
Medicines for endocrine disorders Oral > Solid > tablet: 200 mg	Cushing syndrome

Lamivudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 50 mg per 5 mL Oral > Solid: 150 mg tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + nevirapine + zidovudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals	Therapeutic equivalent to emtricitabine + tenofovir pour Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Therapeutic equivalent to emtricitabine + tenofovir pour Contact with or exposure to human immunodeficiency virus

Lamivudine + zidovudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamotrigine

Section	Indications
Antiseizure medicines Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet; 200 mg tablet; 2 mg tablet (chewable, dispersible); 5 mg tablet (chewable, dispersible); 25 mg tablet (chewable, dispersible); 50 mg tablet (chewable, dispersible); 100 mg tablet (chewable, dispersible); 200 mg tablet (chewable, dispersible)	Epilepsy or seizures

Latanoprost

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 50 µg per mL eye drops	Primary open-angle glaucoma

Ledipasvir + sofosbuvir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 90 mg + 400 mg tablet	Chronic hepatitis C

Lenalidomide

Section	Indications
Immunomodulators Oral > Solid: 25 mg	Plasma cell myeloma

Letrozole

Section	Indications
Ovulation inducers Oral > Solid > dosage form: 2.5 mg	Female infertility without specification whether primary or secondary

Leuprorelin

Section	Indications
Hormones and antihormones Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe	Malignant neoplasms of breast Malignant neoplasms of prostate

Levamisole

Section	Indications
Intestinal anthelminthics Oral > Solid > tablet: 50 mg (as hydrochloride); 150 mg (as hydrochloride)	Ascariasis

Levetiracetam

Section	Indications
Antiseizure medicines Oral > Liquid: 100 mg per mL Oral > Solid > tablet: 250 mg; 500 mg; 750 mg; 1000 mg Parenteral > General injections > IV: 500 mg per 5 mL in 5 mL vial; 5 mg per mL in 100 mL bag; 10 mg per mL in 100 mL bag; 15 mg per mL in 100 mL bag	Epilepsy or seizures Status epilepticus

Levodopa

Section	Indications
Antiseizure medicines Oral > Solid: 250 mg	Parkinson disease

Levodopa + benserazide

Section	Indications
Medicines for parkinsonism	Therapeutic equivalent to levodopa + carbidopa pour Parkinson disease

Levodopa + carbidopa

Section	Indications
Medicines for parkinsonism Oral > Solid: 100 mg + 10 mg tablet; 250 mg + 25 mg tablet; 100 mg + 25 mg tablet	Parkinson disease

Levofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 250 mg; 500 mg; 750 mg	Multi-drug resistant Mycobacterium tuberculosis

Levonorgestrel

Section	Indications
Oral hormonal contraceptives Oral > Solid: 750 µg [2]; 1.5 mg [1] Oral > Solid: 30 µg	Contact with health services for postcoital contraception Contact with health services for contraceptive management
Intrauterine devices Intrauterine system with reservoir containing 52 mg of levonorgestrel.	Contact with health services for insertion of contraceptive device

Levonorgestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 75 mg per rod (two-rods)	Contact with health services for insertion of contraceptive device

Levothyroxine

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 µg (sodium salt); 100 µg (sodium salt); 25 µg (sodium salt)	Hypothyroidism

Lidocaine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)	Ventricular tachyarrhythmia
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 5% in 2 mL ampoule (hydrochloride) + 7.5% glucose solution Parenteral > Locoregional injections > Other: 1% in vial (hydrochloride); 2% in vial (hydrochloride) Local > Topical > unspecified: 2 to 4% (hydrochloride)	Local anaesthetics

Lidocaine + epinephrine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Other: 1% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial; 2% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial Local > Dental > Cartridge: 2% (lidocaine hydrochloride) + 1:80 000 epinephrine	Local anaesthetics

Lindane

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Cream: 1% Local > Topical > Lotion: 1%	Scabies

Linezolid

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 2 mg per mL in 300 mL bag Oral > Liquid: 100 mg per 5 mL powder for oral liquid Oral > Solid: 400 mg tablet; 600 mg tablet Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 600 mg	Other specified bacterial diseases Methicillin resistant Staphylococcus aureus Vancomycin resistant Staphylococcus aureus Vancomycin resistant Enterococcus
Antituberculosis medicines Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 600 mg	Multi-drug resistant Mycobacterium tuberculosis

Liraglutide

Section	Indications
Medicines for endocrine disorders Parenteral > General injections > SC: 6 mg per mL in 3 mL pre-filled pen	Obesity

Lisinopril + amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg	Essential hypertension

Lisinopril + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg	Essential hypertension

Lisocabtagene maraleucel

Section	Indications
Antineoplastics and supportive medicines	Therapeutic equivalent to axicabtagene ciloleucel pour Diffuse large B-cell lymphomas

Lithium carbonate

Section	Indications
Medicines for bipolar disorders Oral > Solid: 300 mg	Bipolar or related disorders

Long-acting insulin analogues

Section	Indications
Insulins Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen	Type 1 diabetes mellitus Type 2 diabetes mellitus

Loperamide

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid > dosage form: 2 mg	Palliative care

Lopinavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid > tablet: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg tablet (heat-stable) Oral > Solid > dosage form: 40 mg + 10 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Loratadine

Section	Indications
Antiallergics and medicines used in anaphylaxis Oral > Liquid: 1 mg per mL Oral > Solid: 10 mg	Allergic or hypersensitivity conditions of unspecified type

Lorazepam

Section	Indications
Antiseizure medicines Parenteral > General injections > IV: 2 mg per mL in 1 mL ampoule; 4 mg per mL in 1 mL ampoule	Status epilepticus
Medicines for anxiety disorders	Therapeutic equivalent to diazepam pour Anxiety

Losartan

Section	Indications
Antihypertensive medicines Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet	Essential hypertension
Medicines used in heart failure Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet	Heart failure

Lovastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Magnesium sulfate

Section	Indications
Antiseizure medicines Parenteral > General injections > IV: 500 mg per mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume) Parenteral > General injections > IM: 500 mg per mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume)	Severe pre-eclampsia Eclampsia

Mannitol

Section	Indications
Diuretics Parenteral > General injections > IV: 10% solution; 20% solution	Ocular hypertension Cerebral oedema

Measles vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against measles alone

Mebendazole

Section	Indications
Intestinal anthelminthics Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)	Ascariasis Enterobiasis Trichuriasis Hookworm diseases
Cysticidal medicines Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)	Other specified echinococcosis

Medroxyprogesterone acetate

Section	Indications
Injectable hormonal contraceptives Parenteral > General injections > IM: 150 mg per mL in 1 mL vial Parenteral > General injections > SC: 104 mg per 0.65 mL in pre-filled syringe; 104 mg per 0.65 mL in single dose injection delivery system	Contact with health services for reasons associated with reproduction
Progestogens Oral > Solid: 5 mg	Excessive menstruation with irregular cycle

Mefloquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 250 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Antimalarial medicines > For chemoprevention Oral > Solid: 250 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum

Meglumine antimoniate

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule	Visceral leishmaniasis Cutaneous leishmaniasis Mucocutaneous leishmaniasis

Meglumine iotroxate

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL	X-ray contrast media, iodinated

Melarsoprol

Section	Indications
Medicines for the treatment of 2nd stage African trypanosomiasis Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)	African trypanosomiasis

Melphalan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection Oral > Solid: 2 mg	Plasma cell myeloma

Meningococcal meningitis vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Mercaptopurine

Section	Indications
Cytotoxic medicines Oral > Solid: 50 mg tablet Oral > Liquid: 20 mg per mL	Unspecified malignant neoplasms of ill-defined or unspecified sites Acute myeloid leukaemia with recurrent genetic abnormalities Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Meropenem

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Multi-drug resistant Mycobacterium tuberculosis
Watch group antibiotics Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Second choix Neutropenia (high-risk) Peritoneal abscess (severe) Peritonitis (severe)

Meropenem + vaborbactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 1 g in vial + 1 g in vial powder for injection	Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales Carbapenem resistant Acinetobacter baumannii

Mesalazine

Section	Indications
Anti-inflammatory medicines	Therapeutic equivalent to sulfasalazine pour Crohn disease site Therapeutic equivalent to sulfasalazine pour Ulcerative colitis

Mesna

Section	Indications
Supportive medicines Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule Oral > Solid: 400 mg; 600 mg	Osteosarcoma of bone and articular cartilage of other specified sites Other specified malignant neoplasms of the ovary Germ cell tumour of testis Ewing sarcoma of bone and articular cartilage of unspecified sites Rhabdomyosarcoma primary site Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis

Metformin

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 500 mg (hydrochloride)	Type 2 diabetes mellitus

Methadone

Section	Indications
Medicines for opioid use disorders Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)	Opioid dependence
Opioid analgesics Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride) Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)	Chronic cancer pain

Methimazole

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 5 mg; 10 mg; 20 mg	Thyrotoxicosis

Methionine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg (DL)	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methotrexate

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (as sodium salt) Oral > Solid: 2.5 mg (as sodium salt) Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified Acute myeloid leukaemia with recurrent genetic abnormalities Malignant trophoblastic neoplasms of placenta Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis
Dermatological medicines > Medicines affecting skin differentiation and proliferation Oral > Solid > tablet: 2.5 mg (as sodium salt); 10 mg (as sodium salt)	Psoriasis of unspecified type
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 2.5 mg (as sodium salt)	Rheumatoid arthritis, serology unspecified
Immunomodulators for non-malignant disease Parenteral > General injections > SC: pre-filled syringe in multiple strengths	Crohn disease site Psoriatic arthritis
Medicines for juvenile joint diseases Oral > Solid > tablet: 2.5 mg (as sodium salt)	Juvenile idiopathic arthritis

Methoxy polyethylene glycol-epoetin beta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Methyldopa

Section	Indications
Antihypertensive medicines Oral > Solid: 250 mg	Essential hypertension Gestational hypertension

Methylergometrine

Section	Indications
Uterotonics	Therapeutic equivalent to ergometrine pour Postpartum haemorrhage

Methylphenidate

Section	Indications
Medicines for mental and behavioural disorders Multiple formulations and strengths.	Attention deficit hyperactivity disorder

Methylprednisolone

Section	Indications
Hormones and antihormones Parenteral > General injections > IV: 40 mg per mL in 1 mL single-dose vial (as sodium succinate); 40 mg per mL in 5 mL multi-dose vial (as sodium succinate); 80 mg per mL in 1 mL single-dose vial (as sodium succinate)	Burkitt lymphoma including Burkitt leukaemia

Methylthioninium chloride

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule	Acquired methaemoglobinaemia

Metoclopramide

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 5 mg per 5 mL Oral > Solid: 10 mg (hydrochloride) Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	Palliative care
Antiemetic medicines Oral > Liquid: 5 mg per 5 mL Oral > Solid: 10 mg (hydrochloride) Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	Nausea or vomiting

Metoprolol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Metronidazole

Section	Indications
Access group antibiotics Oral > Liquid: 200 mg per 5 mL (as benzoate) Oral > Solid: 200 to 500 mg tablet Parenteral > General injections > unspecified: 500 mg per 100 mL in vial Injection Local > Rectal > Suppository: 500 mg; 1 g Parenteral > General injections > unspecified: 500 mg per 100 mL in vial Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg	Premier choix Intestinal infections due to Clostridioides difficile Trichomoniasis co-prescrite avec cefazolin Other specified prophylactic measures co-prescrite avec cefotaxime Peritonitis (severe) co-prescrite avec cefotaxime Peritonitis (mild-moderate) co-prescrite avec cefotaxime Peritoneal abscess (mild-moderate) co-prescrite avec cefotaxime Peritoneal abscess (severe) co-prescrite avec ceftriaxone Necrotising fasciitis co-prescrite avec ceftriaxone Peritonitis (severe) co-prescrite avec ceftriaxone Peritoneal abscess (mild-moderate) co-prescrite avec ceftriaxone Peritoneal abscess (severe) co-prescrite avec ceftriaxone Peritonitis (mild-moderate) Second choix co-prescrite avec ciprofloxacin Peritonitis (mild-moderate) co-prescrite avec ciprofloxacin Peritoneal abscess (mild-moderate) Autres indications Bacterial infection of unspecified site
Antiamoebic and antigiardiasis medicines Oral > Liquid: 200 mg per 5 mL (as benzoate) Parenteral > General injections > unspecified: 500 mg in 100 mL vial Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg	Amoebiasis

Micafungin

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection	Systemic or invasive candidosis

Miconazole

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Cream: 2% (nitrate) Local > Topical > Ointment: 2% (nitrate)	Fungal infection of the skin

Midazolam

Section	Indications
Antiseizure medicines Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration); 5 mg per mL in 1 mL vial (for buccal administration); 5 mg per mL in 3 mL vial (for buccal administration) Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)	Therapeutic equivalent to lorazepam pour Status epilepticus Status epilepticus
Preoperative medication and sedation for short-term procedures Parenteral > General injections > IV: 1 mg per mL Oral > Liquid: 2 mg per mL Oral > Solid > dosage form: 15 mg; 7.5 mg	Sedative-hypnotic and anxiolytic drugs
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per mL Parenteral > General injections > unspecified: 1 mg per mL; 5 mg per mL Oral > Solid > dosage form: 7.5 mg; 15 mg	Palliative care

Mifepristone - misoprostol

Section	Indications
Uterotonics Oral > Solid: 200 mg + 200 µg; 200 mg [1] + 200 µg [4] in co-package	Induced abortion Unspecified time of fetal death, cause not specified

Miltefosine

Section	Indications
Antileishmaniasis medicines Oral > Solid: 10 mg; 50 mg	Visceral leishmaniasis Mucocutaneous leishmaniasis Cutaneous leishmaniasis

Misoprostol

Section	Indications
Uterotonics Oral > Solid: 200 µg Local > Vaginal > tablet: 25 µg	Spontaneous abortion, incomplete, without complication Unspecified obstetric condition Postpartum haemorrhage

Mometasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Mometasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Morphine

Section	Indications
Opioid analgesics Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate) Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate) Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate) Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)	Pain
Preoperative medication and sedation for short-term procedures Parenteral > General injections > IV: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)	Opioids or related analgesics and agents affecting opioid receptors

Moxifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 400 mg	Multi-drug resistant Mycobacterium tuberculosis Tuberculosis

Multiple micronutrient supplement

Section	Indications
Other medicines administered to the mother Tablet containing: Vit A: 800 mcg retinol activity equivalent Vit C: 70 mg Vit D: 5 mcg (200 IU) Vit E: 10 mg alpha tocopherol equivalent Vit B1: 1.4 mg Vit B2: 1.4 mg Vit B3: 18 mg niacin equivalent Vit B6: 1.9 mg Folic acid: 400 mcg Vit B12: 2.6 mg Iron: 30 mg Iodine: 150 mcg Zinc: 15 mg Selenium: 65 mcg Copper: 2 mg	Maternal care for other specified conditions predominantly related to pregnancy

Mumps vaccine

Section	Indications
Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against mumps alone

Mupirocin

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Cream: 2% (as mupirocin calcium) Local > Topical > Ointment: 2%	Superficial bacterial folliculitis Impetigo

Nadroparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin pour Acute ischaemic heart disease Therapeutic equivalent to enoxaparin pour Venous thromboembolism

Nalidixic acid

Section	Indications
Anti-infective medicines Oral > Solid: 250 mg; 500 mg	Shigella resistant to unspecified antibiotic

Naloxone

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)	Unintentional exposure to or harmful effects of opioids or related analgesics

Naltrexone

Section	Indications
Medicines for alcohol use disorders Parenteral > General injections > IM: 380 mg in vial (extended-release) Oral > Solid > tablet: 50 mg	Disorders due to use of alcohol

Natamycin

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Suspension: 5% eye drops	Infectious keratitis

Nelfinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 250 mg (as mesilate) Oral > Other: 50 mg per g oral powder	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Neostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Oral > Solid: 15 mg (neostigmine bromide) Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate); 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate)	Myasthenia gravis Anticholinesterase reversible

Netilmicin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis Therapeutic equivalent to gentamicin pour Infectious blepharitis

Nevirapine

Section	Indications
Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors Oral > Liquid: 50 mg per 5 mL Oral > Solid > dispersible tablet: 50 mg Oral > Solid > tablet: 200 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Niclosamide

Section	Indications
Intestinal anthelminthics Oral > Solid: 500 mg tablet (chewable)	Hymenolepiasis Diphyllobothriasis Taeniasis due to Taenia saginata Taeniasis due to Taenia solium

Nicotinamide

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg	Pellagra

Nicotine replacement therapy

Section	Indications
Medicines for nicotine use disorders Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour Oral > Solid > lozenge: 2 mg; 4 mg lozenge Local > Buccal > Oral spray: 1 mg per actuation Local > Buccal > Chewing gum: 2 mg; 4 mg	Nicotine dependence

Nifedipine

Section	Indications
Antioxytocics (tocolytics) Oral > Solid: 10 mg (immediate-release)	Preterm labour without delivery
Antianginal medicines Oral > Solid: 10 mg	Angina pectoris
Antihypertensive medicines Oral > Solid: 10 mg tablet; sustained-release formulations	Essential hypertension

Nifurtimox

Section	Indications
American trypanosomiasis Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)	Chagas disease
Medicines for the treatment of 2nd stage African trypanosomiasis Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)	African trypanosomiasis

Nilotinib

Section	Indications
Targeted therapies Oral > Solid > capsule: 150 mg; 200 mg	Chronic myeloid leukaemia, not elsewhere classified

Nilutamide

Section	Indications
Hormones and antihormones	Therapeutic equivalent to bicalutamide pour Malignant neoplasms of prostate

Nitrofurantoin

Section	Indications
Access group antibiotics Oral > Liquid: 25 mg per 5 mL oral liquid Oral > Solid > dosage form: 100 mg; 50 mg	Premier choix Infectious cystitis

Nitrous oxide

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation:	Anaesthetics and therapeutic gases

Nivolumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion	Other specified malignant neoplasms of bronchus or lung Melanoma of skin

Norethisterone

Section	Indications
Oral hormonal contraceptives Oral > Solid: 350 µg Progestogens Oral > Solid: 5 mg	Contact with health services for contraceptive management Excessive menstruation with irregular cycle Therapeutic equivalent to medroxyprogesterone acetate pour Excessive menstruation with irregular cycle

Norethisterone enantate

Section	Indications
Injectable hormonal contraceptives Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule oily solution	Contact with health services for reasons associated with reproduction

Normal immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IV: 5% protein solution; 10% protein solution Parenteral > General injections > IM: 16% protein solution Parenteral > General injections > SC: 15% protein solution; 16% protein solution	Primary immunodeficiencies Langerhans cell histiocytosis Mucocutaneous lymph node syndrome

Nystatin

Section	Indications
Antifungal medicines Oral > Liquid: 100000 IU per mL Local > Topical > Other: 100000 IU pessary Oral > Solid > dosage form: 500000 IU Oral > Solid > lozenge: 100000 IU	Candidosis

Ocrelizumab

Section	Indications
Medicines for multiple sclerosis Parenteral > General injections > IV: 30 mg per mL in 10 vial concentrate for solution Parenteral > General injections > IV: 30 mg per mL in 10 mL vial	Multiple sclerosis Relapsing-remitting multiple sclerosis

Octreotide

Section	Indications
Medicines for disorders of the pituitary hormone system Parenteral > General injections > IM: 20 mg in vial (modified-release, as acetate) plus diluent Parenteral > General injections > SC: 0.05 mg per mL in 1 mL vial (immediate-release, as acetate); 0.1 mg per mL in 1 mL vial (immediate-release, as acetate); 0.5 mg per mL in 1 mL vial (immediate-release, as acetate)	Acromegaly or pituitary gigantism

Ofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.3%	Other specified conjunctivitis Infectious keratitis

Olanzapine

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 10 mg in vial powder for injection	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Omadacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 100 mg lyophilized powder for injection Oral > Solid: 300 mg	Methicillin resistant Staphylococcus aureus

Ombitasvir + paritaprevir + ritonavir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 12.5 mg + 75 mg + 50 mg tablet	Chronic hepatitis C

Omeprazole

Section	Indications
Antiulcer medicines Oral > Liquid: 20 mg powder for oral liquid; 40 mg powder for oral liquid Oral > Solid: 10 mg; 20 mg; 40 mg Parenteral > General injections > IV: 40 mg in vial	Peptic ulcer, site unspecified Gastro-oesophageal reflux disease Gastrointestinal bleeding

Ondansetron

Section	Indications
Medicines for other common symptoms in palliative care Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride) Oral > Liquid: 4 mg per 5 mL (ondansetron base) Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base)	Palliative care
Antiemetic medicines Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride) Oral > Liquid: 4 mg per 5 mL (ondansetron base) Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base); 24 mg (ondansetron base)	Nausea or vomiting

Oral rehydration salts

Section	Indications
Oral rehydration Powder for dilution in 200 mL, 500 mL, 1 L: - glucose 75 mEq - sodium 75 mEq or mmol/L - chloride 65 mEq or mmol/L - potassium 20 mEq or mmol/L - citrate 10 mmol/L - osmolarity 245 mOsm/L - glucose 13.5 g/L - sodium chloride 2.6 g/L - potassium chloride 1.5 g/L - trisodium citrate dihydrate 2.9 g/L	Diarrhoea
Solutions correcting water, electrolyte and acid-base disturbances > Oral Powder for dilution in 200 mL, 500 mL, 1 L: - glucose 75 mEq - sodium 75 mEq or mmol/L - chloride 65 mEq or mmol/L - potassium 20 mEq or mmol/L - citrate 10 mmol/L - osmolarity 245 mOsm/L - glucose 13.5 g/L - sodium chloride 2.6 g/L - potassium chloride 1.5 g/L - trisodium citrate dihydrate 2.9 g/L	Other specified disorders of fluid, electrolyte or acid-base balance

Oseltamivir

Section	Indications
Other antivirals Oral > Solid: 30 mg capsule (as phosphate); 45 mg capsule (as phosphate); 75 mg capsule (as phosphate)	Influenza due to identified seasonal influenza virus

Osimertinib

Section	Indications
Targeted therapies Oral > Solid: 40 mg (as mesylate); 80 mg (as mesylate)	Other specified malignant neoplasms of bronchus or lung

Oxaliplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of colon

Oxamniquine

Section	Indications
Antischistosomals and other antitrematode medicines Oral > Liquid: 250 mg per 5 mL Oral > Solid > capsule: 250 mg	Schistosomiasis due to Schistosoma mansoni

Oxazolindinones

Section	Indications
Reserve group antibiotics .	Other specified bacterial diseases

Oxycodone

Section	Indications
Opioid analgesics	Therapeutic equivalent to morphine pour Pain

Oxygen

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation: (medical gas)	Anaesthetics and therapeutic gases
Medical gases Respiratory > Inhalation:	Respiratory failure as acute or chronic

Oxytetracycline

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to tetracycline pour Other specified conjunctivitis Therapeutic equivalent to tetracycline pour Infectious keratitis Therapeutic equivalent to tetracycline pour Infectious blepharitis Therapeutic equivalent to tetracycline pour Trachoma

Oxytocin

Section	Indications
Uterotonics Parenteral > General injections > unspecified: 10 IU per mL	Postpartum haemorrhage

P-aminosalicylate sodium

Section	Indications
Antituberculosis medicines Oral > Liquid: 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid) powder for oral solution	Multi-drug resistant Mycobacterium tuberculosis

P-aminosalicylic acid

Section	Indications
Antituberculosis medicines Oral > Solid: 4 g granules in sachet	Multi-drug resistant Mycobacterium tuberculosis

Paclitaxel

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 6 mg per mL	Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Kaposi sarcoma of unspecified primary site Other specified malignant neoplasms of breast Other specified carcinomas of ovary Malignant neoplasms of breast Malignant neoplasms of cervix uteri

Palbociclib

Section	Indications
Targeted therapies Oral > Solid: 75 mg; 100 mg; 125 mg	Other specified malignant neoplasms of breast

Paliperidone

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 25 mg in pre-filled syringe (as palmitate); 50 mg in pre-filled syringe (as palmitate); 75 mg in pre-filled syringe (as palmitate); 100 mg in pre-filled syringe (as palmitate); 150 mg in pre-filled syringe (as palmitate)	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Palonosetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Paracetamol (acetaminophen)

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 100 mg; 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Pain
Antimigraine medicines > For treatment of acute attack Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Migraine

Paromomycin

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)	Visceral leishmaniasis
Antiamoebic and antigiardiasis medicines Oral > Liquid: 125 mg per 5 mL as sulfate Oral > Solid: 250 mg as sulfate	Amoebiasis

Pegaspargase

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection	Lymphoid leukaemia, not elsewhere classified

Pegfilgrastim

Section	Indications
Immunomodulators Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe	Acquired neutropaenia

Pegylated interferon alfa (2a)

Section	Indications
Medicines for hepatitis C > Other antivirals for hepatitis C Parenteral > General injections > SC: 180 µg in vial; 180 µg in prefilled syringe	Chronic hepatitis C

Pegylated interferon alfa (2b)

Section	Indications
Medicines for hepatitis C > Other antivirals for hepatitis C Parenteral > General injections > SC: 80 µg in prefilled syringe; 100 µg in prefilled syringe	Chronic hepatitis C

Pembrolizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 50 mg powder for injection	Therapeutic equivalent to nivolumab pour Melanoma of skin Other specified malignant neoplasms of bronchus or lung

Penicillamine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 250 mg	Rheumatoid arthritis, serology unspecified

Pentamidine

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)	Visceral leishmaniasis
Medicines for the treatment of 1st stage African trypanosomiasis Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection	African trypanosomiasis
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid > tablet: 200 mg (as isethionate); 300 mg (as isethionate)	Pneumocystosis

Permethrin

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Lotion: 1% Local > Topical > Cream: 5%	Pediculosis corporis Pediculosis capitis Scabies

Pertussis vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against pertussis alone

Pertuzumab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 240 mg per 14 mL in vial concentrated solution	Carcinoma of breast, specialised type

Phenelzine

Section	Indications
Medicines for depressive disorders Oral > Solid > dosage form: 15 mg (as sulfate)	Other specified depressive disorders

Phenobarbital

Section	Indications
Antiseizure medicines Oral > Liquid: 15 mg per 5 mL Oral > Solid: 15 to 100 mg Parenteral > General injections > IV: 200 mg per mL (phenobarbital sodium); 30 mg per mL (phenobarbital sodium); 60 mg per mL (phenobarbital sodium)	Epilepsy or seizures Status epilepticus

Phenoxymethylpenicillin

Section	Indications
Access group antibiotics Oral > Liquid: 250 mg per 5 mL (as potassium salt) powder for oral liquid Oral > Solid: 250 mg (as potassium salt) tablet Oral > Solid: 250 mg (as potassium salt); 500 mg (as potassium salt)	Premier choix Acute pharyngitis Periapical abscess without sinus Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) Autres indications Bacterial infection of unspecified site

Phenytoin

Section	Indications
Antiseizure medicines Oral > Liquid: 30 mg per 5 mL (phenytoin) Parenteral > General injections > unspecified: 50 mg per mL in vial (phenytoin sodium) Oral > Solid > tablet: 50 mg (phenytoin) (chewable) Oral > Solid > dosage form: 25 mg (phenytoin sodium); 50 mg (phenytoin sodium); 100 mg (phenytoin sodium)	Epilepsy or seizures

Phosphorus

Section	Indications
Medicines for endocrine disorders Oral > Other: 500 mg in sachet (elemental phosphorus) granules Oral > Solid > tablet: 250 mg (elemental phosphorus); 500 mg (elemental phosphorus); 500 mg (elemental phosphorus) (effervescent)	Hypophosphataemic rickets

Phytomenadione

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule Oral > Solid: 10 mg tablet	Haemorrhagic disorder due to other specified circulating anticoagulants

Pilocarpine

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 2% (hydrochloride or nitrate); 4% (hydrochloride or nitrate)	Acute angle closure with pupillary block Ocular hypertension Primary open-angle glaucoma

Piperacillin + tazobactam

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 2 g (as sodium salt) + 250 mg (as sodium salt) powder for injection; 4 g (as sodium salt) + 500 mg (as sodium salt) powder for injection	Premier choix Peritoneal abscess (severe) Other specified pneumonia (Hospital-acquired pneumonia) Neutropenia (high-risk) Peritonitis (severe) co-prescrite avec clindamycin Necrotising fasciitis

Platelets

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Plazomicin

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500 mg per 10 mL	Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Pneumococcal vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Podophyllotoxin

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to podophyllum resin pour Anogenital warts Therapeutic equivalent to podophyllum resin pour Plantar warts

Podophyllum resin

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Solution: 10 to 25%	Anogenital warts Plantar warts

Poliomyelitis vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against poliomyelitis

Polygeline

Section	Indications
Plasma substitutes Parenteral > General injections > IV: 3.5% injectable solution Parenteral > General injections > IV: 3.5% in solution	Therapeutic equivalent to dextran 70 pour Hypovolaemia Hypovolaemia

Polymyxin B (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500000 IU in vial powder for injection (equivalent to 50 mg polymyxin B base)	Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales

Polymyxins

Section	Indications
Reserve group antibiotics .	Other specified bacterial diseases

Potassium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Oral Oral > Liquid: powder for solution	Hypokalaemia
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 11.2% in 20 mL ampoule (equivalent to K+ 1.5 mmol/mL and Cl- 1.5 mmol/mL); 7.5% solution for dilution (equivalent to K+ 1 mmol/mL and Cl- 2 mmol/mL); 15% solution for dilution (equivalent to K+ 2 mmol/mL and Cl- 2 mmol/mL)	Hypokalaemia

Potassium ferric hexacyanoferrate (Prussian blue)

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: Powder for oral administration	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium iodide

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 60 mg	Thyrotoxicosis
Antifungal medicines Local > Topical > Solution: Saturated solution	Sporotrichosis

Potassium permanganate

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Solution: aqueous solution: 1:10 000	Tropical phagedaenic ulcer Pemphigus Other specified pyogenic bacterial infection of skin and subcutaneous tissue Impetigo

Povidone iodine

Section	Indications
Antiseptics Local > Topical > Solution: 10% (equivalent to 1% available iodine)	Povidone iodine

Pralidoxime

Section	Indications
Antidotes and other substances used in poisonings > Specific .	Exposure to or harmful effects of undetermined intent of pesticides

Pramipexole

Section	Indications
Antiseizure medicines Oral > Solid: 125 µg immediate-release; 250 mg immediate-release; 500 µg immediate-release; 750 µg immediate-release; 1 mg immediate-release; 1.5 mg immediate-release; 375 µg extended-release; 750 µg extended-release; 1.5 mg extended-release; 2.25 mg extended-release; 3 mg extended-release; 3.75 mg extended-release; 4.5 extended-release	Parkinson disease

Pravastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Praziquantel

Section	Indications
Antischistosomals and other antitrematode medicines Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Paragonimiasis Clonorchiasis Opisthorchiasis Schistosomiasis
Intestinal anthelminthics Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Taeniasis due to Taenia saginata Taeniasis due to Taenia solium Hymenolepiasis Diphyllobothriasis
Cysticidal medicines Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Cysticercosis of central nervous system

Precipitated sulfur

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Ointment:	Therapeutic equivalent to benzyl benzoate pour Pthiriasis Therapeutic equivalent to benzyl benzoate pour Pediculosis corporis Therapeutic equivalent to benzyl benzoate pour Pediculosis capitis Therapeutic equivalent to benzyl benzoate pour Scabies

Prednisolone

Section	Indications
Hormones and antihormones Oral > Liquid: 5 mg per mL Oral > Solid: 5 mg; 25 mg	Unspecified malignant neoplasms of ill-defined or unspecified sites Plasma cell myeloma Follicular lymphoma Lymphoid leukaemia, not elsewhere classified Malignant neoplasms of prostate Chronic lymphocytic leukaemia or small lymphocytic lymphoma Hodgkin lymphoma Diffuse large B-cell lymphomas Burkitt lymphoma including Burkitt leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis
Anti-inflammatory medicines Local > Rectal > Retention enema: Local > Rectal > Retention enema: 20 mg per 100 mL (as sodium phosphate)	Therapeutic equivalent to hydrocortisone pour Crohn disease site Crohn disease site Ulcerative colitis
Antiallergics and medicines used in anaphylaxis Oral > Liquid: 5 mg per mL Oral > Solid: 5 mg; 25 mg	Allergic or hypersensitivity conditions of unspecified type
Ophthalmological preparations > Anti-inflammatory agents Local > Ophthalmological > Solution (eye drops): 0.5% (sodium phosphate)	Scleritis Anterior uveitis

Prednisone

Section	Indications
Hormones and antihormones	Therapeutic equivalent to prednisolone pour Plasma cell myeloma Therapeutic equivalent to prednisolone pour Follicular lymphoma Therapeutic equivalent to prednisolone pour Lymphoid leukaemia, not elsewhere classified Therapeutic equivalent to prednisolone pour Malignant neoplasms of prostate Therapeutic equivalent to prednisolone pour Chronic lymphocytic leukaemia or small lymphocytic lymphoma Therapeutic equivalent to prednisolone pour Hodgkin lymphoma Therapeutic equivalent to prednisolone pour Diffuse large B-cell lymphomas Therapeutic equivalent to prednisolone pour Anaplastic large cell lymphoma, ALK-negative Therapeutic equivalent to prednisolone pour Anaplastic large cell lymphoma, ALK-positive Therapeutic equivalent to prednisolone pour Langerhans cell histiocytosis
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to prednisolone pour Allergic or hypersensitivity conditions of unspecified type

Pretomanid

Section	Indications
Antituberculosis medicines Oral > Solid > tablet: 200 mg	Multi-drug resistant Mycobacterium tuberculosis

Primaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)	Malaria due to Plasmodium ovale Malaria due to Plasmodium vivax

Probenecid

Section	Indications
Medicines used to treat gout Oral > Solid: 500 mg	Gout

Procainamide

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride) Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)	Cardiac arrhythmia

Procaine benzylpenicillin

Section	Indications
Access group antibiotics Parenteral > General injections > IM: 1 g in vial (=1 million IU) powder for injection; 3 g in vial (=3 million IU) powder for injection	Second choix Syphilis

Procarbazine

Section	Indications
Cytotoxic medicines Oral > Solid: 50 mg (as hydrochloride)	Unspecified malignant neoplasms of ill-defined or unspecified sites

Progesterone vaginal ring

Section	Indications
Intravaginal contraceptives Local > Vaginal > vaginal ring: 2.074 g micronized progesterone	Contact with health services for contraceptive management

Proguanil

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 100 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum

Promethazine

Section	Indications
Antiemetic medicines Oral > Liquid: 5 mg per 5 mL (hydrochloride) Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule	Nausea or vomiting

Propanol

Section	Indications
Antiseptics	Therapeutic equivalent to ethanol pour Denatured alcohol

Propofol

Section	Indications
General anaesthetics and oxygen > Injectable medicines Parenteral > General injections > IV: 10 mg per mL; 20 mg per mL	Anaesthetics and therapeutic gases

Propranolol

Section	Indications
Antihypertensive medicines Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Essential hypertension
Antianginal medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Angina pectoris
Antiarrhythmic medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Cardiac arrhythmia
Antimigraine medicines > For prophylaxis Oral > Solid: 40 mg tablet (hydrochloride); 20 mg tablet (hydrochloride)	Migraine

Propylthiouracil

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 mg	Thyrotoxicosis

Protamine sulfate

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 10 mg per mL in 5 mL ampoule	Haemorrhagic disorder due to other specified circulating anticoagulants

Protionamide

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to ethionamide pour Multi-drug resistant Mycobacterium tuberculosis

Pyrantel

Section	Indications
Intestinal anthelminthics Oral > Solid > tablet: 250 mg (chewable) (as embonate or pamoate)	Ascariasis Trichuriasis Enterobiasis Hookworm diseases

Pyrazinamide

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 400 mg; 500 mg	Tuberculosis

Pyridostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 1 mg in 1 mL ampoule (pyridostigmine bromide) Oral > Solid: 60 mg (pyridostigmine bromide)	Myasthenia gravis

Pyridoxine

Section	Indications
Vitamins and minerals Oral > Solid: 25 mg (hydrochloride)	Vitamin B6 deficiency Sideroblastic anaemia Other specified polyneuropathy

Pyrimethamine

Section	Indications
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid: 25 mg	Toxoplasmosis

Quetiapine

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders
Medicines for bipolar disorders Oral > Solid > tablet: 25 mg (immediate-release); 100 mg (immediate-release); 150 mg (immediate-release); 200 mg (immediate-release); 300 mg (immediate-release); 50 mg (modified-release); 150 mg (modified-release); 200 mg (modified-release); 300 mg (modified-release); 400 mg (modified-release)	Bipolar or related disorders

Quinidine

Section	Indications
Antiarrhythmic medicines Oral > Solid: 200 mg (sulfate)	Cardiac arrhythmia

Quinine

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride) Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)	Malaria due to Plasmodium falciparum

Rabies vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against rabies

Raltegravir

Section	Indications
Antiretrovirals > Integrase inhibitors Oral > Liquid: 100 mg granules for oral suspension Oral > Solid: 400 mg tablet; 25 mg tablet (chewable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Ranibizumab

Section	Indications
Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations .	Retinal venous occlusions Diabetic macular oedema Degenerative high myopia Age related macular degeneration

Ranitidine

Section	Indications
Antiulcer medicines Oral > Liquid: 75 mg per 5 mL (as hydrochloride) Oral > Solid: 150 mg (as hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (as hydrochloride)	Gastro-oesophageal reflux disease Peptic ulcer, site unspecified

Rasburicase

Section	Indications
Supportive medicines Parenteral > General injections > IV: 1.5 mg in vial powder and solvent for solution; 7.5 mg in vial powder and solvent for solution	Tumour lysis syndrome

Ravidasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg	Chronic hepatitis C

Realgar-indigo naturalis formulation

Section	Indications
Cytotoxic medicines Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)	Acute myeloid leukaemia with recurrent genetic abnormalities

Red blood cells

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Resin-based composite (high-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use capsule or multi-use syringe	Dental caries

Resin-based composite (low-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use applicator or multi-use bottle	Dental caries

Retinol

Section	Indications
Vitamins and minerals Parenteral > General injections > IM: 100000 IU in 2 mL ampoule (as palmitate) water-miscible injection Oral > Liquid: 100000 IU per mL (as palmitate) oral oily solution in multidose dispenser Oral > Solid: 10000 IU (as palmitate) sugar-coated tablet; 200000 IU (as palmitate) capsule; 50000 IU (as palmitate) capsule; 100000 IU (as palmitate) capsule	Vitamin A deficiency

Ribavirin

Section	Indications
Other antivirals Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution Oral > Solid: 200 mg; 400 mg; 600 mg	Viral haemorrhagic fever, not elsewhere classified
Medicines for hepatitis C > Other antivirals for hepatitis C Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution Oral > Solid: 200 mg; 400 mg; 600 mg	Chronic hepatitis C

Ribociclib

Section	Indications
Targeted therapies	Therapeutic equivalent to palbociclib pour Other specified malignant neoplasms of breast

Riboflavin

Section	Indications
Vitamins and minerals Oral > Solid: 5 mg	Vitamin B2 deficiency

Rifabutin

Section	Indications
Antituberculosis medicines Oral > Solid: 150 mg	Tuberculosis

Rifampicin

Section	Indications
Antituberculosis medicines Oral > Liquid: 20 mg per mL Oral > Solid: 150 mg; 300 mg Parenteral > General injections > IV: 600 mg in vial powder for injection	Tuberculosis Other specified tuberculosis
Antileprosy medicines Oral > Liquid: 20 mg per mL Oral > Solid: 150 mg; 300 mg	Leprosy

Rifapentine

Section	Indications
Antituberculosis medicines Oral > Solid > tablet: 150 mg; 300 mg	Tuberculosis Latent tuberculosis

Risdiplam

Section	Indications
Medicines for diseases of the nervous system Oral > Liquid: 0.75 mg per mL powder for oral liquid	Spinal muscular atrophy

Risperidone

Section	Indications
Medicines for psychotic disorders Oral > Solid: 0.25 mg; 0.5 mg; 1 mg; 2 mg; 3 mg; 4 mg; 6 mg Parenteral > General injections > IM:	Schizophrenia or other primary psychotic disorders Therapeutic equivalent to paliperidone pour Schizophrenia or other primary psychotic disorders

Ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Rituximab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial	Follicular lymphoma Chronic lymphocytic leukaemia or small lymphocytic lymphoma Diffuse large B-cell lymphomas Burkitt lymphoma including Burkitt leukaemia
Medicines for multiple sclerosis Parenteral > General injections > IV: 500 mg per 50 mL in 50 mL vial	Multiple sclerosis

Rivaroxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Ropinirole

Section	Indications
Antiseizure medicines Oral > Solid: 0.25 mg immediate-release; 0.5 mg immediate-release; 1 mg immediate-release; 2 immediate-release; 3 mg immediate-release; 4 mg immediate-release; 5 mg immediate-release; 2 mg extended-release; 4 mg extended-release; 6 mg extended-release; 8 mg extended-release; 12 mg extended-release	Parkinson disease

Rotavirus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Rubella vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against rubella alone

Salbutamol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Parenteral > General injections > IV: 50 µg per mL in 5 mL ampoule (as sulfate) Respiratory > Inhalation > aerosol: 100 µg per dose (as sulfate); 100 µg per dose (as sulfate) metered-dose inhaler Respiratory > Inhalation > solution: 5 mg per mL (as sulfate) for use in nebulizers	Chronic obstructive pulmonary disease Asthma

Salicylic acid

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Solution: 5%	Seborrhoeic dermatitis Psoriasis of unspecified type Ichthyoses

Saquinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 200 mg; 500 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Selenium sulfide

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Suspension: 2% detergent-based	Seborrhoeic dermatitis Pityriasis versicolor

Senna

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 7.5 mg per 5 mL oral liquid	Palliative care
Laxatives Oral > Solid: 7.5 mg (sennosides) (or traditional dosage forms)	Constipation

Sevoflurane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > liquid:	Anaesthetics and therapeutic gases

Silver diamine fluoride

Section	Indications
Dental medicines and preparations Local > Dental > Solution: 38% w/v	Dental caries

Silver nitrate

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 1%	Neonatal conjunctivitis or dacryocystitis

Silver sulfadiazine

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Cream: 1%	Burns

Simeprevir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 150 mg	Chronic hepatitis C

Simvastatin

Section	Indications
Lipid-lowering agents Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg	Mixed hyperlipidaemia Coronary atherosclerosis
Medicines for endocrine disorders Oral > Solid: 20 mg	Polycystic ovary syndrome

Smallpox vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Snake antivenom

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies .	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium calcium edetate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 0.9% isotonic (equivalent to Na+ 154 mmol/L and Cl- 154 mmol/L)	Other specified disorders of fluid, electrolyte or acid-base balance

Sodium hyaluronate

Section	Indications
Ophthalmological preparations	Therapeutic equivalent to hypromellose pour Keratoconjunctivitis sicca

Sodium hydrogen carbonate

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 1.4% isotonic (equivalent to Na+ 167 mmol/L and HCO3- 167 mmol/L); 8.4% in 10 mL ampoule (equivalent to Na+ 1000 mmol/L and HCO3- 1000 mmol/L)	Acidosis

Sodium nitrite

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitroprusside

Section	Indications
Antihypertensive medicines Parenteral > General injections > IV: 50 mg in ampoule powder for infusion	Hypertensive crisis

Sodium stibogluconate

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial	Visceral leishmaniasis Mucocutaneous leishmaniasis Cutaneous leishmaniasis

Sodium thiosulfate

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Solution: 15%	Pityriasis versicolor
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sofosbuvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg; 400 mg	Chronic hepatitis C

Sofosbuvir + velpatasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg + 50 mg; 400 mg + 100 mg	Chronic hepatitis C

Spectinomycin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 2 g in vial (as hydrochloride) powder for injection	Second choix Gonococcal infection Autres indications Bacterial infection of unspecified site

Spironolactone

Section	Indications
Medicines used in heart failure Oral > Solid > tablet: 25 mg	Heart failure
Diuretics Oral > Liquid: 5 mg per 5 mL; 10 mg per 5 mL; 25 mg per 5 mL Oral > Solid > tablet: 25 mg	Oedema Ascites

Stavudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 5 mg per 5 mL powder for oral solution Oral > Solid: 15 mg; 20 mg; 30 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Streptokinase

Section	Indications
Thrombolytic medicines Parenteral > General injections > IV: 1.5 million IU in vial powder for injection	Acute myocardial infarction

Streptomycin (injection)

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1 g powder for injection (as sulfate) as vial	Multi-drug resistant Mycobacterium tuberculosis

Succimer

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 100 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sulfacetamide

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 10% (sodium salt)	Other specified conjunctivitis

Sulfadiazine

Section	Indications
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid: 500 mg	Toxoplasmosis
Anti-infective medicines Oral > Solid: 500 mg Parenteral > General injections > unspecified: 250 mg in 4 mL ampoule (sodium salt)	Bacterial infection of unspecified site

Sulfadimidine

Section	Indications
Anti-infective medicines Oral > Liquid: 500 mg per 5 mL Oral > Solid: 500 mg Parenteral > General injections > unspecified: 1 g in 3 mL ampoule (sodium salt)	Bacterial infection of unspecified site

Sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 500 mg + 25 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Antimalarial medicines > For chemoprevention Oral > Solid: 500 mg + 25 mg tablet	Malaria due to Plasmodium falciparum

Sulfamethoxazole + trimethoprim

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 80 mg + 16 mg per mL in 10 mL ampoule; 80 mg + 16 mg per mL in 5 mL ampoule Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet Oral > Solid > dispersible tablet: 100 mg + 20 mg	Premier choix Infectious cystitis Second choix Infectious gastroenteritis or colitis without specification of infectious agent Autres indications Bacterial infection of unspecified site
Antipneumocystosis and antitoxoplasmosis medicines Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet Oral > Solid > dispersible tablet: 100 mg + 20 mg	Pneumocystosis

Sulfasalazine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 500 mg	Rheumatoid arthritis, serology unspecified
Anti-inflammatory medicines Oral > Solid: 500 mg Local > Rectal > Suppository: 500 mg Local > Rectal > Retention enema: 3 g per 100 mL	Crohn disease site Ulcerative colitis

Sumatriptan

Section	Indications
Antimigraine medicines > For treatment of acute attack Oral > Solid: 50 mg	Migraine

Sunscreen (broad-spectrum)

Section	Indications
Dermatological medicines Topical cream, lotion, gel, ointment, spray with sun protection factor ≥ 50.	Albinism or other specified genetically-determined hypomelanotic disorders Xeroderma pigmentosum

Suramin sodium

Section	Indications
Antifilarials Parenteral > General injections > IV: 1 g in vial	Filariasis
Medicines for the treatment of 1st stage African trypanosomiasis Parenteral > General injections > IV: 1 g in vial	African trypanosomiasis

Suxamethonium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 50 mg per mL in 2 mL ampoule (suxamethonium chloride); powder for injection (suxamethonium chloride) in vial	Muscle relaxants

Tacalcitol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to calcipotriol pour Psoriasis of unspecified type

Tacrolimus

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 5 mg per mL in 1 mL vial Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)	Failure or rejection of transplanted organs or tissues

Tamoxifen

Section	Indications
Hormones and antihormones Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)	Other specified malignant neoplasms of breast Malignant neoplasms of breast

Tedizolid

Section	Indications
Reserve group antibiotics	Therapeutic equivalent to linezolid pour Methicillin resistant Staphylococcus aureus Therapeutic equivalent to linezolid pour Vancomycin resistant Staphylococcus aureus Therapeutic equivalent to linezolid pour Vancomycin resistant Enterococcus

Telmisartan + amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg	Essential hypertension

Telmisartan + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg	Essential hypertension

Tenofovir alafenamide

Section	Indications
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 25 mg	Chronic hepatitis B

Tenofovir disoproxil fumarate

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Contact with or exposure to human immunodeficiency virus
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)	Chronic hepatitis B

Terbinafine

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Cream: 1% terbinafine hydrochloride Local > Topical > Ointment: 1% terbinafine hydrochloride	Fungal infection of the skin

Terbutaline

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to salbutamol pour Chronic obstructive pulmonary disease Therapeutic equivalent to salbutamol pour Asthma

Terizidone

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to cycloserine pour Multi-drug resistant Mycobacterium tuberculosis

Testosterone

Section	Indications
Androgens Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule (enantate)	Testicular dysfunction or testosterone-related disorders

Tetanus antitoxin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IM: 50000 IU in vial	Tetanus

Tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against tetanus alone

Tetracaine

Section	Indications
Ophthalmological preparations > Local anaesthetics Local > Ophthalmological > Solution (eye drops): 0.5% (hydrochloride)	Local anaesthetics

Tetracycline

Section	Indications
Access group antibiotics Oral > Solid: 250 mg (hydrochloride)	Bacterial infection of unspecified site
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)	Other specified conjunctivitis Infectious keratitis Infectious blepharitis Trachoma

Thalidomide

Section	Indications
Immunomodulators Oral > Solid: 50 mg	Plasma cell myeloma

Thiamine

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg (hydrochloride)	Vitamin B1 deficiency

Thiopental

Section	Indications
General anaesthetics and oxygen > Injectable medicines	Therapeutic equivalent to propofol pour Anaesthetics and therapeutic gases

Ticagrelor

Section	Indications
Anti-platelet medicines Oral > Solid > tablet: 60 mg; 90 mg	Acute ischaemic heart disease Presence of coronary angioplasty implant or graft

Tick-borne encephalitis vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against arthropod-borne viral encephalitis

Tigecycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg in vial powder for injection	Other specified bacterial diseases

Timolol

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 0.25% (hydrogen maleate); 0.5% (hydrogen maleate)	Primary open-angle glaucoma Ocular hypertension

Tinidazole

Section	Indications
Antiamoebic and antigiardiasis medicines	Therapeutic equivalent to metronidazole pour Amoebiasis

Tiotropium bromide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > dry powder: 18 µg capsule Respiratory > Inhalation > solution: 1.25 µg per actuation; 2.5 µg per actuation	Chronic obstructive pulmonary disease

Tisagenlecleucel

Section	Indications
Antineoplastics and supportive medicines	Therapeutic equivalent to axicabtagene ciloleucel pour Diffuse large B-cell lymphomas

Tislelizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 100 mg per 10 mL	Malignant neoplasms of urinary tract Hodgkin lymphoma Other specified malignant neoplasms of bronchus or lung

Tobramycin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis Therapeutic equivalent to gentamicin pour Infectious blepharitis

Tocilizumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	Juvenile systemic arthritis

Tolbutamide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 500 mg	Type 2 diabetes mellitus

Torasemide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure
Diuretics	Therapeutic equivalent to furosemide pour Anuria or oliguria Therapeutic equivalent to furosemide pour Oedema

Toripalimab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 240 mg per 6 mL in vial	Malignant neoplasms of nasopharynx Malignant neoplasms of oesophagus

Tramadol

Section	Indications
Opioid analgesics Oral > Liquid: 100 mg per mL (hydrochloride) Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)	Chronic cancer pain

Trametinib

Section	Indications
Targeted therapies Oral > Solid: 0.5 mg; 2 mg	Other specified melanoma of skin

Tranexamic acid

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Haemorrhage, not elsewhere classified
Other medicines administered to the mother Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Postpartum haemorrhage

Trastuzumab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 60 mg in vial powder for injection; 150 mg in vial powder for injection; 440 mg in vial powder for injection	Carcinoma of breast, specialised type

Trastuzumab emtansine

Section	Indications
Targeted therapies Parenteral > General injections > IV: 100 mg in vial powder for injection; 160 mg in vial powder for injection	Carcinoma of breast, specialised type

Triamcinolone hexacetonide

Section	Indications
Medicines for juvenile joint diseases Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial	Juvenile idiopathic arthritis

Triclabendazole

Section	Indications
Antischistosomals and other antitrematode medicines Oral > Solid > tablet: 250 mg (scored)	Paragonimiasis Fascioliasis

Trihexyphenidyl

Section	Indications
Antiseizure medicines Oral > Solid: 2 mg (hydrochloride); 5 mg (hydrochloride)	Parkinson disease
Medicines for parkinsonism	Therapeutic equivalent to biperiden pour Parkinson disease

Trimethoprim

Section	Indications
Access group antibiotics Oral > Liquid: 50 mg per mL Oral > Solid: 100 mg; 200 mg	Premier choix Infectious cystitis

Triptorelin

Section	Indications
Hormones and antihormones	Therapeutic equivalent to leuprorelin pour Malignant neoplasms of breast Therapeutic equivalent to leuprorelin pour Malignant neoplasms of prostate

Tropicamide

Section	Indications
Diagnostic agents > Ophthalmic medicines Local > Ophthalmological > Solution (eye drops): 0.5%	Cycloplegic drug

Tropisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Tuberculin, purified protein derivative

Section	Indications
Immunologicals > Diagnostic agents Parenteral > Locoregional injections > Intradermal:	Mycobacterial diseases

Tubocurarine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg per mL in 1.5 mL ampoule (tubocurarine chloride)	Muscle relaxants

Typhoid vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against typhoid-paratyphoid alone

Ulipristal

Section	Indications
Oral hormonal contraceptives Oral > Solid: 30 mg tablet (ulipristal acetate)	Contact with health services for postcoital contraception

Umeclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Urea

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Cream: 10%; 5% Local > Topical > Ointment: 10%; 5%	Xerosis cutis or asteatosis

Ustekinumab

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Parenteral > General injections > SC: 45 mg per 0.5 mL in vial; 90 mg per mL in pre-filled syringe	Psoriasis of unspecified type

Valaciclovir

Section	Indications
Antiherpes medicines	Therapeutic equivalent to aciclovir pour Zoster Therapeutic equivalent to aciclovir pour Varicella Therapeutic equivalent to aciclovir pour Herpes simplex infections

Valganciclovir

Section	Indications
Other antivirals Oral > Liquid: 50 mg per mL powder for oral solution Oral > Solid > tablet: 450 mg	Cytomegaloviral retinitis

Valproic acid (sodium valproate)

Section	Indications
Antiseizure medicines Oral > Liquid: 200 mg per 5 mL Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated); 100 mg tablet (crushable) Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule; 100 mg per mL in 3 mL ampoule	Epilepsy or seizures Status epilepticus
Medicines for bipolar disorders Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated)	Bipolar or related disorders

Vancomycin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection Oral > Solid > capsule: 125 mg (as hydrochloride); 250 mg (as hydrochloride) Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection Parenteral > Locoregional injections > Intravitreal: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection	Premier choix Necrotising fasciitis co-prescrite avec ceftazidime Endophthalmitis co-prescrite avec ceftriaxone Endophthalmitis Second choix Neutropenia (high-risk) Intestinal infections due to Clostridioides difficile Autres indications Methicillin resistant Staphylococcus aureus

Varenicline

Section	Indications
Medicines for nicotine use disorders Oral > Solid: 0.5 mg; 1 mg	Nicotine dependence

Varicella vaccine

Section	Indications
Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Vecuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg in vial powder for injection (vecuronium bromide)	Muscle relaxants

Vemurafenib

Section	Indications
Targeted therapies	Therapeutic equivalent to dabrafenib pour Other specified melanoma of skin

Verapamil

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride) Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)	Supraventricular tachyarrhythmia
Antianginal medicines Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)	Angina pectoris

Vinblastine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate) Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Germ cell tumour of testis Anaplastic large cell lymphoma, ALK-positive Anaplastic large cell lymphoma, ALK-negative Langerhans cell histiocytosis

Vincristine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate) Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Malignant trophoblastic neoplasms of placenta Follicular lymphoma Kaposi sarcoma of unspecified primary site Rhabdomyosarcoma primary site Hodgkin lymphoma Diffuse large B-cell lymphomas Retinoblastoma Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Vinorelbine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Oral > Solid > capsule: 20 mg; 30 mg; 80 mg	Other specified malignant neoplasms of bronchus or lung Other specified malignant neoplasms of breast Rhabdomyosarcoma primary site

Voriconazole

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 200 mg in vial powder for injection Oral > Liquid: 40 mg per mL powder for oral liquid Oral > Solid: 50 mg tablet; 200 mg tablet	Chronic pulmonary aspergillosis Invasive aspergillosis

Warfarin

Section	Indications
Medicines affecting coagulation Oral > Solid: 5 mg (sodium salt); 1 mg (sodium salt); 2 mg (sodium salt); 0.5 mg (sodium salt)	Other specified diseases of arteries or arterioles Cerebral ischaemic stroke due to embolic occlusion Venous thromboembolism Atrial fibrillation

Water for injection

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Miscellaneous Parenteral > General injections > unspecified: 2 mL ampoule; 5 mL ampoule; 10 mL ampoule	Unspecified appropriateness of dosing or administration without injury or harm

Whole blood

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Yellow fever vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Zanubrutinib

Section	Indications
Targeted therapies Oral > Solid: 80 mg	Mantle cell lymphoma Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Zidovudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Parenteral > General injections > IV: 10 mg per mL in 20 mL vial solution for IV infusion Oral > Liquid: 50 mg per 5 mL Oral > Solid > tablet: 300 mg Oral > Solid > capsule: 250 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Zinc sulfate

Section	Indications
Medicines for diarrhoea Oral > Solid: 20 mg	Diarrhoea

Zoledronic acid

Section	Indications
Supportive medicines Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle	Malignant neoplasm metastasis in bone or bone marrow
Medicines for endocrine disorders Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle	Osteogenesis imperfecta

Zuclopenthixol decanoate

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to fluphenazine pour Schizophrenia or other primary psychotic disorders