



Liste modèle des médicaments essentiels

Caffeine citrate

Section	Indications
<p>Medicines administered to the neonate [c]</p> <p>Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL)</p> <p>Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)</p>	Apnoea of newborn

Cefotaxime

Section	Indications
<p>Watch group antibiotics</p> <p>Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)</p>	Bacterial infection of unspecified site

Chlorhexidine

Section	Indications
<p>Medicines administered to the neonate [c]</p> <p>Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)</p> <p>Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)</p>	Omphalitis of newborn

Erythromycin

Section	Indications
<p>Ophthalmological preparations > Anti-infective agents</p> <p>Local > Ophthalmological > Ointment: 0.5%</p>	<p>Chlamydial conjunctivitis</p> <p>Gonococcal infection of eye</p>

Gentamicin

Section	Indications
<p>Access group antibiotics</p> <p>Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)</p>	<p>Premier choix</p> <p>Neonatal meningitis</p>

Ibuprofen

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 5 mg per mL	Patent arterial duct

Indometacin

Section	Indications
Medicines administered to the neonate [c]	Therapeutic equivalent to ibuprofen pour Patent arterial duct

Phytomenadione

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule	Haemorrhagic or haematological disorders of fetus or newborn

Prostaglandin E1

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 0.5 mg per mL in alcohol	Other specified congenital anomaly of great arteries including arterial duct

Prostaglandin E2

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 1 mg per mL	Therapeutic equivalent to prostaglandin E1 pour Other specified congenital anomaly of great arteries including arterial duct
	Other specified congenital anomaly of great arteries including arterial duct

Surfactant

Section	Indications
Medicines administered to the neonate [c] Respiratory > Suspension: 80 mg per mL for intratracheal instillation; 25 mg per mL for intratracheal instillation	Respiratory distress syndrome of newborn

Tetracycline

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)	Neonatal conjunctivitis or dacryocystitis