

Alprostadil (prostaglandin E1)

Section

Medicines administered to the neonate [c]
Parenteral > General injections > IV: 0.5 mg per mL in alcohol

Indications

Other specified congenital anomaly of great arteries including arterial duct

Beractant

Section

Medicines administered to the neonate [c]
Respiratory > Suspension: 25 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Caffeine citrate

Section

Medicines administered to the neonate [c]
Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL)
Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)

Indications

Apnoea of newborn

Cefotaxime

Section

Watch group antibiotics
Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)

Indications

Bacterial infection of unspecified site

Chlorhexidine

Section

Medicines administered to the neonate [c]
Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)
Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)

Indications

Omphalitis of newborn

Dinoprostone (prostaglandin E2)

Section

Medicines administered to the neonate [c]
Parenteral > General injections > IV: 1 mg per mL

Indications

Other specified congenital anomaly of great arteries including arterial duct

Therapeutic equivalent to alprostadil (prostaglandin E1) pour

Other specified congenital anomaly of great arteries including arterial duct

Erythromycin

Section

Ophthalmological preparations > Anti-infective agents
Local > Ophthalmological > Ointment: 0.5%

Indications

Chlamydial conjunctivitis

Gonococcal infection of eye

Gentamicin

Section

Access group antibiotics
Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Medicines for bacterial central nervous system infections
Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Indications

Premier choix

Neonatal meningitis

Premier choix

Neonatal meningitis

Ibuprofen

Section

Medicines administered to the neonate [c]
Parenteral > General injections > IV: 5 mg per mL

Indications

Patent arterial duct

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to ibuprofen pour

Patent arterial duct

Phytomenadione

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 1 mg per 0.5 mL in ampoule; 10 mg per mL in ampoule; 1 mg per mL in ampoule; 2 mg per 0.2 mL in ampoule mixed micelle solution; 10 mg per mL in ampoule mixed micelle solution

Indications

Haemorrhagic or haematological disorders of fetus or newborn

Porcatant alfa

Section

Medicines administered to the neonate [c]

Respiratory > Suspension: 80 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Tetracycline

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)

Indications

Neonatal conjunctivitis or dacryocystitis