

Alprostadil (prostaglandin E1)

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 0.5 mg per mL in alcohol	Other specified congenital anomaly of great arteries including arterial duct

Beractant

Section	Indications
Medicines administered to the neonate [c] Respiratory > Suspension: 25 mg per mL for intratracheal instillation	Respiratory distress syndrome of newborn

Caffeine citrate

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL) Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)	Apnoea of newborn

Cefotaxime

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)	Bacterial infection of unspecified site

Chlorhexidine

Section	Indications
Medicines administered to the neonate [c] Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine) Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)	Omphalitis of newborn

Dinoprostone (prostaglandin E2)

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 1 mg per mL

Indications

Other specified congenital anomaly of great arteries including arterial duct

Therapeutic equivalent to alprostadil (prostaglandin E1) for

Other specified congenital anomaly of great arteries including arterial duct

Erythromycin

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Ointment: 0.5%

Indications

Chlamydial conjunctivitis

Gonococcal infection of eye

Gentamicin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Medicines for bacterial central nervous system infections

Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Indications

First choice

Neonatal meningitis

First choice

Neonatal meningitis

Ibuprofen

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 5 mg per mL

Indications

Patent arterial duct

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to ibuprofen for

Patent arterial duct

Phytomenadione

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 1 mg per 0.5 mL in ampoule; 10 mg per mL in ampoule; 1 mg per mL in ampoule; 2 mg per 0.2 mL in ampoule mixed micelle solution; 10 mg per mL in ampoule mixed micelle solution

Indications

Haemorrhagic or haematological disorders of fetus or newborn

Porcatant alfa

Section

Medicines administered to the neonate [c]

Respiratory > Suspension: 80 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Tetracycline

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)

Indications

Neonatal conjunctivitis or dacryocystitis