

Caffeine citrate

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL) Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)	Apnoea of newborn

Cefotaxime

Section	Indications
Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)	Bacterial infection of unspecified site

Chlorhexidine

Section	Indications
Medicines administered to the neonate [c] Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine) Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)	Omphalitis of newborn

Erythromycin

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 0.5%	Chlamydial conjunctivitis Gonococcal infection of eye

Gentamicin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)	First choice Neonatal meningitis

Ibuprofen

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 5 mg per mL

Indications

Patent arterial duct

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to **ibuprofen** for

Patent arterial duct

Phytomenadione

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule

Indications

Haemorrhagic or haematological disorders of fetus or newborn

Prostaglandin E1

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 0.5 mg per mL in alcohol

Indications

Other specified congenital anomaly of great arteries including arterial duct

Prostaglandin E2

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 1 mg per mL

Indications

Therapeutic equivalent to **prostaglandin E1** for

Other specified congenital anomaly of great arteries including arterial duct

Other specified congenital anomaly of great arteries including arterial duct

Surfactant

Section

Medicines administered to the neonate [c]

Respiratory > Suspension: 80 mg per mL for intratracheal instillation; 25 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Tetracycline

Section

[Ophthalmological preparations > Anti-infective agents](#)

Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)

Indications

Neonatal conjunctivitis or dacryocystitis