

Codeine

Section	Indications
<p>Opioid analgesics</p> <p>Oral > Solid: 30 mg tablet (codeine phosphate)</p>	<p>Pain</p>

Fentanyl

Section	Indications
<p>Opioid analgesics</p> <p>Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour</p> <p>Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate)</p> <p>Oral > Other: 200 to 1200 µg buccal film (as citrate)</p>	<p>Chronic cancer pain</p> <p>Other specified chronic cancer related pain</p>

Hydromorphone

Section	Indications
<p>Opioid analgesics</p>	<p>Therapeutic equivalent to <i>morphine</i> for</p> <p>Pain</p>

Methadone

Section	Indications
<p>Opioid analgesics</p> <p>Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)</p> <p>Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)</p>	<p>Chronic cancer pain</p>

Morphine

Section

Opioid analgesics

Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate)

Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate)

Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate)

Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)

Indications

Pain

Oxycodone

Section

Opioid analgesics

Indications

Therapeutic equivalent to [morphine](#) for

Pain

Tramadol

Section

Opioid analgesics

Oral > Liquid: 100 mg per mL (hydrochloride)

Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release

Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)

Indications

Chronic cancer pain