

Amphotericin B

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	<input type="text" value="Blastomycosis"/> <input type="text" value="Cryptococcosis"/> <input type="text" value="Candidosis"/> <input type="text" value="Coccidioidomycosis"/> <input type="text" value="Histoplasmosis"/> <input type="text" value="Sporotrichosis"/> <input type="text" value="Aspergillosis"/>

Anidulafungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin for <input type="text" value="Systemic or invasive candidosis"/>

Caspofungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin for <input type="text" value="Systemic or invasive candidosis"/>

Clotrimazole

Section	Indications
Antifungal medicines Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet	<input type="text" value="Vulvovaginal candidosis"/>

Fluconazole

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 2 mg per mL in vial Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid Oral > Solid > capsule: 50 mg	<input type="text" value="Cryptococcosis"/> <input type="text" value="Candidosis"/>

Flucytosine

Section

Antifungal medicines

Parenteral > General injections > IV: 2.5 g per 250 mL infusion

Oral > Solid: 250 mg capsule

Indications

Cryptococcosis

Griseofulvin

Section

Antifungal medicines

Oral > Liquid: 125 mg per 5 mL

Oral > Solid: 125 mg; 250 mg

Indications

Dermatophytosis

Itraconazole

Section

Antifungal medicines

Oral > Liquid: 10 mg per mL

Oral > Solid: 100 mg

Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Histoplasmosis

Talaromycosis

Sporotrichosis

Paracoccidioidomycosis

Micafungin

Section

Antifungal medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection

Indications

Systemic or invasive candidosis

Nystatin

Section

Antifungal medicines

Oral > Liquid: 100000 IU per mL

Local > Topical > Other: 100000 IU pessary

Oral > Solid > dosage form: 500000 IU

Oral > Solid > lozenge: 100000 IU

Indications

Candidosis

Potassium iodide

Section

Antifungal medicines

Local > Topical > Solution: Saturated solution

Indications

Sporotrichosis

Voriconazole

Section

Antifungal medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection

Oral > Liquid: 40 mg per mL powder for oral liquid

Oral > Solid: 50 mg tablet; 200 mg tablet

Indications

Chronic pulmonary aspergillosis

Invasive aspergillosis