

Amodiaquine

Section

Antimalarial medicines > Medicines for curative treatment
 Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artemether

Section

Antimalarial medicines > Medicines for curative treatment
 Parenteral > General injections > IM: 20 mg per mL in 1 mL ampoule (oily injection); 40 mg per mL in 1 mL ampoule (oily injection)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artemether + lumefantrine

Section

Antimalarial medicines > Medicines for curative treatment
 Oral > Solid > dispersible tablet: 20 mg + 120 mg
 Oral > Solid > tablet: 20 mg + 120 mg

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate

Section

Antimalarial medicines > Medicines for curative treatment
 Parenteral > General injections > IV: 60 mg in vial powder for injection; 30 mg in vial powder for injection; 120 mg in vial powder for injection
 Local > Rectal > Other: 100 mg capsules

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + amodiaquine

Section

Antimalarial medicines > Medicines for curative treatment
 Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg
 Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + mefloquine

Section

Antimalarial medicines > Medicines for curative treatment
 Oral > Solid > tablet: 25 mg + 50 mg (as hydrochloride); 100 mg + 200 mg (as hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + pyronaridine

Section

[Antimalarial medicines](#) > [Medicines for curative treatment](#)
Oral > Solid > tablet: 60 mg + 180 mg (tetraphosphate)
Oral > Solid > granules: 20 mg + 60 mg (tetraphosphate)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate - sulfadoxine + pyrimethamine

Section

[Antimalarial medicines](#) > [Medicines for curative treatment](#)
Oral > Solid > tablet: 50 mg [3] + 500 mg + 25 mg [1] in copackage (scored)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Chloroquine

Section

[Antimalarial medicines](#) > [For curative treatment](#)
Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)
Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)
Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

[Antimalarial medicines](#) > [Medicines for curative treatment](#)
Oral > Liquid: 50 mg per 5 mL (base)
Oral > Solid > tablet: 150 mg (base)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Dihydroartemisinin + piperaquine

Section

[Antimalarial medicines](#) > [Medicines for curative treatment](#)
Oral > Solid > dispersible tablet: 20 mg + 160 mg (phosphate); 40 mg + 320 mg (phosphate)
Oral > Solid > tablet: 40 mg + 320 mg (phosphate); 20 mg + 160 mg (phosphate); 60 mg + 480 (phosphate); 80 mg + 640 mg (phosphate)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Doxycycline

Section

[Antimalarial medicines](#) > [Medicines for curative treatment](#)
Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Indications

Malaria due to *Plasmodium falciparum*

Mefloquine

Section

Antimalarial medicines > Medicines for curative treatment
Oral > Solid: 250 mg tablet (hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Primaquine

Section

Antimalarial medicines > Medicines for curative treatment
Oral > Solid > tablet: 7.5 mg (as phosphate); 15 mg (as phosphate)

Indications

Malaria due to *Plasmodium ovale*

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Quinine

Section

Antimalarial medicines > Medicines for curative treatment
Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion; 60 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion

Indications

Malaria due to *Plasmodium falciparum*

Sulfadoxine + pyrimethamine

Section

Antimalarial medicines > Medicines for curative treatment
Oral > Solid: 500 mg + 25 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*