

Adalimumab

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 2px;">Axial spondyloarthritis</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 2px;">Crohn disease site</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rheumatoid arthritis</div>

Azathioprine

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg in vial powder for injection (as sodium salt); 100 mg in vial powder for injection (as sodium salt) Oral > Liquid: 10 mg per mL Oral > Solid > tablet: 25 mg; 50 mg (scored) Oral > Solid > tablet: 50 mg (scored)	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 2px;">Failure or rejection of transplanted organs or tissues</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Relapsing-remitting multiple sclerosis</div>

Certolizumab pegol

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 2px;">Axial spondyloarthritis</div> Therapeutic equivalent to adalimumab for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 2px;">Crohn disease site</div> Therapeutic equivalent to adalimumab for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rheumatoid arthritis</div>

Ciclosporin

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection Oral > Liquid: 100 mg per mL Oral > Solid > capsule: 25 mg	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Failure or rejection of transplanted organs or tissues</div>

Etanercept

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Fingolimod

Section

Immunomodulators for non-malignant disease

Oral > Solid: 0.25 mg; 0.5 mg

Indications

Relapsing-remitting multiple sclerosis

Golimumab

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Infliximab

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Methotrexate

Section

Immunomodulators for non-malignant disease

Parenteral > General injections > SC: pre-filled syringe in multiple strengths

Indications

Crohn disease site

Psoriatic arthritis

Tacrolimus

Section

Immunomodulators for non-malignant disease

Parenteral > General injections > IV: 5 mg per mL in 1 mL vial

Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension

Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)

Indications

Failure or rejection of transplanted organs or tissues