

**Acetylsalicylic acid**

## Section

Anti-platelet medicines  
Oral > Solid: 100 mg

## Indications

Acute myocardial infarction

Cerebral ischaemic stroke

**Acetylsalicylic acid + atorvastatin + ramipril**

## Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease  
Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg

## Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

**Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide**

## Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease  
Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg

## Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

**Alteplase**

## Section

Thrombolytic medicines  
Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection

## Indications

Cerebral ischaemic stroke

**Amiodarone**

## Section

Antiarrhythmic medicines  
Parenteral > General injections > IV: 50 mg per mL in 3 ampoules (hydrochloride)  
Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)

## Indications

Cardiac arrhythmia

## Amlodipine

### Section

Antihypertensive medicines

Oral > Solid: 5 mg (as maleate, mesylate or besylate)

### Indications

Essential hypertension

## Atenolol

### Section

Antianginal medicines

Oral > Solid: 50 mg; 100 mg

Antihypertensive medicines

Oral > Solid: 50 mg; 100 mg

### Indications

Angina pectoris

Essential hypertension

Therapeutic equivalent to *bisoprolol* for

Essential hypertension

## Atorvastatin

### Section

Lipid-lowering agents

### Indications

Therapeutic equivalent to *simvastatin* for

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* for

Coronary atherosclerosis

## Atorvastatin + perindopril + amlodipine

### Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg

### Indications

Chronic ischaemic heart disease

Atherosclerotic chronic arterial occlusive disease

Cerebral ischaemic stroke

## Bisoprolol

### Section

Antianginal medicines

Oral > Solid: 1.25 mg; 5 mg

Medicines used in heart failure

Oral > Solid: 1.25 mg; 5 mg

Antihypertensive medicines

Oral > Solid: 1.25 mg; 5 mg

Antiarrhythmic medicines

Oral > Solid: 1.25 mg; 5 mg

### Indications

Angina pectoris

Heart failure

Essential hypertension

Cardiac arrhythmia

## Bumetanide

### Section

Medicines used in heart failure

### Indications

Therapeutic equivalent to [furosemide](#) for

Heart failure

## Captopril

### Section

Medicines used in heart failure

Oral > Solid: 25 mg

### Indications

Heart failure

## Carvedilol

### Section

Antianginal medicines

Medicines used in heart failure

Antihypertensive medicines

Antiarrhythmic medicines

### Indications

Therapeutic equivalent to [bisoprolol](#) for

Angina pectoris

Therapeutic equivalent to [bisoprolol](#) for

Heart failure

Therapeutic equivalent to [bisoprolol](#) for

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) for

Cardiac arrhythmia

## Chlorothiazide

### Section

Antihypertensive medicines

Medicines used in heart failure

### Indications

Therapeutic equivalent to hydrochlorothiazide for

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide for

Heart failure

## Chlortalidone

### Section

Antihypertensive medicines

Medicines used in heart failure

### Indications

Therapeutic equivalent to hydrochlorothiazide for

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide for

Heart failure

## Clopidogrel

### Section

Anti-platelet medicines

Oral > Solid: 75 mg; 300 mg

### Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

## Digitoxin

### Section

Medicines used in heart failure

Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule

Oral > Solid: 50 µg; 100 µg

### Indications

Heart failure

## Digoxin

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid: 62.5 µg; 250 µg

#### Medicines used in heart failure

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg

### Indications

Cardiac arrhythmia

Heart failure

## Dopamine

### Section

#### Medicines used in heart failure

Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)

### Indications

Heart failure

Cardiogenic shock

## Enalapril

### Section

#### Antihypertensive medicines

Oral > Liquid: 1 mg per mL (as hydrogen maleate)

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

#### Medicines used in heart failure

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

### Indications

Essential hypertension

Heart failure

## Epinephrine

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)

### Indications

Cardiac arrest

## Fluvastatin

### Section

Lipid-lowering agents

### Indications

Therapeutic equivalent to *simvastatin* for

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* for

Coronary atherosclerosis

## Furosemide

### Section

Medicines used in heart failure

Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL

Oral > Solid > tablet: 40 mg; 20 mg

### Indications

Heart failure

## Glyceryl trinitrate

### Section

Antianginal medicines

Oral > Solid: 500 µg sublingual tablet

### Indications

Angina pectoris

## Hydralazine

### Section

Antihypertensive medicines

Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)

Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)

Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)

### Indications

Essential hypertension

Gestational hypertension

## Hydrochlorothiazide

### Section

Antihypertensive medicines

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg; 12.5 mg

Medicines used in heart failure

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg

### Indications

Essential hypertension

Heart failure

## Indapamide

### Section

Antihypertensive medicines

Medicines used in heart failure

### Indications

Therapeutic equivalent to hydrochlorothiazide for

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide for

Heart failure

## Isoprenaline

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)

Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)

### Indications

Bradycardia

Cardiogenic shock

## Isosorbide dinitrate

### Section

Antianginal medicines

Oral > Solid: 5 mg sublingual tablet

### Indications

Angina pectoris

## Lidocaine

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)

### Indications

Ventricular tachyarrhythmia

## Lisinopril + amlodipine

### Section

Antihypertensive medicines

Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg

### Indications

Essential hypertension

## Lisinopril + hydrochlorothiazide

### Section

Antihypertensive medicines

Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg

### Indications

Essential hypertension

## Losartan

### Section

Antihypertensive medicines

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Medicines used in heart failure

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

### Indications

Essential hypertension

Heart failure

## Lovastatin

### Section

Lipid-lowering agents

### Indications

Therapeutic equivalent to *simvastatin* for

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* for

Coronary atherosclerosis

## Methyldopa

### Section

Antihypertensive medicines

Oral > Solid: 250 mg

### Indications

Essential hypertension

Gestational hypertension

## Metoprolol

### Section

Antianginal medicines

Therapeutic equivalent to *bisoprolol* for

Angina pectoris

Medicines used in heart failure

Therapeutic equivalent to *bisoprolol* for

Heart failure

Antihypertensive medicines

Therapeutic equivalent to *bisoprolol* for

Essential hypertension

Antiarrhythmic medicines

Therapeutic equivalent to *bisoprolol* for

Cardiac arrhythmia



## Nifedipine

### Section

#### Antianginal medicines

Oral > Solid: 10 mg

#### Antihypertensive medicines

Oral > Solid: 10 mg tablet; sustained-release formulations

### Indications

Angina pectoris

Essential hypertension

## Pravastatin

### Section

#### Lipid-lowering agents

### Indications

Therapeutic equivalent to *simvastatin* for

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* for

Coronary atherosclerosis

## Procainamide

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)

Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)

### Indications

Cardiac arrhythmia

## Propranolol

### Section

#### Antihypertensive medicines

Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

#### Antianginal medicines

Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)

Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)

Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

### Indications

Essential hypertension

Angina pectoris

Cardiac arrhythmia

## Quinidine

### Section

Antiarrhythmic medicines  
Oral > Solid: 200 mg (sulfate)

### Indications

Cardiac arrhythmia

## Simvastatin

### Section

Lipid-lowering agents  
Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg

### Indications

Mixed hyperlipidaemia

Coronary atherosclerosis

## Sodium nitroprusside

### Section

Antihypertensive medicines  
Parenteral > General injections > IV: 50 mg in ampoule powder for infusion

### Indications

Hypertensive crisis

## Spironolactone

### Section

Medicines used in heart failure  
Oral > Solid > tablet: 25 mg

### Indications

Heart failure

## Streptokinase

### Section

Thrombolytic medicines  
Parenteral > General injections > IV: 1.5 million IU in vial powder for injection

### Indications

Acute myocardial infarction

## Telmisartan + amlodipine

### Section

Antihypertensive medicines  
Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg

### Indications

Essential hypertension

## Telmisartan + hydrochlorothiazide

### Section

Antihypertensive medicines  
Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg

### Indications

Essential hypertension

## Ticagrelor

### Section

Anti-platelet medicines

Oral > Solid > tablet: 60 mg; 90 mg

### Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

## Torsemide

### Section

Medicines used in heart failure

### Indications

Therapeutic equivalent to furosemide for

Heart failure

## Verapamil

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Antianginal medicines

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

### Indications

Supraventricular tachyarrhythmia

Angina pectoris