



Model List of Essential Medicines

Amiodarone

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride) Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)	Cardiac arrhythmia

Bisoprolol

Section	Indications
Antiarrhythmic medicines Oral > Solid: 1.25 mg; 5 mg	Cardiac arrhythmia

Carvedilol

Section	Indications
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol for Cardiac arrhythmia

Digoxin

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid: 62.5 µg; 250 µg	Cardiac arrhythmia

Epinephrine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)	Cardiac arrest

Isoprenaline

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)	Bradycardia Cardiogenic shock
Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)	

Lidocaine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)	Ventricular tachyarrhythmia

Metoprolol

Section	Indications
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol for Cardiac arrhythmia

Procainamide

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)	Cardiac arrhythmia
Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)	

Propranolol

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)	Cardiac arrhythmia
Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	

Quinidine

Section	Indications
Antiarrhythmic medicines Oral > Solid: 200 mg (sulfate)	Cardiac arrhythmia

Verapamil

Section	Indications
<p>Antiarrhythmic medicines</p> <p>Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)</p> <p>Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)</p>	<p>Supraventricular tachyarrhythmia</p>