

Amidotrizoate

Section

Diagnostic agents > Radiocontrast media
Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)

Indications

Amidotrizoate

Atropine

Section

Diagnostic agents > Ophthalmic medicines

Indications

Therapeutic equivalent to tropicamide for

Cycloplegic drug

Barium sulfate

Section

Diagnostic agents > Radiocontrast media
Oral > Liquid: Aqueous suspension

Indications

Barium sulfate with suspending agents

Cyclopentolate

Section

Diagnostic agents > Ophthalmic medicines

Indications

Therapeutic equivalent to tropicamide for

Cycloplegic drug

Fluorescein

Section

Diagnostic agents > Ophthalmic medicines
Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)

Indications

Fluorescein

Iohexol

Section

Diagnostic agents > Radiocontrast media
Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule

Indications

X-ray contrast media, iodinated

Meglumine iotroxate

Section

Diagnostic agents > Radiocontrast media

Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL

Indications

X-ray contrast media, iodinated

Tropicamide

Section

Diagnostic agents > Ophthalmic medicines

Local > Ophthalmological > Solution (eye drops): 0.5%

Indications

Cycloplegic drug