

**Aprepitant**

Section	Indications
Antiemetic medicines Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid > capsule: 80 mg; 125 mg; 165 mg	<input type="text" value="Nausea or vomiting"/>

**Bisacodyl**

Section	Indications
Laxatives	Therapeutic equivalent to <i>senna</i> for <input type="text" value="Constipation"/>

**Cimetidine**

Section	Indications
Antiulcer medicines Oral > Solid: 200 mg Parenteral > General injections > unspecified: 200 mg in 2 mL ampoule	<input type="text" value="Peptic ulcer, site unspecified"/> <input type="text" value="Gastro-oesophageal reflux disease"/>

**Codeine**

Section	Indications
Medicines for diarrhoea Oral > Solid: 30 mg tablet (codeine phosphate)	<input type="text" value="Diarrhoea"/>

**Dexamethasone**

Section	Indications
Antiemetic medicines Oral > Liquid: 0.5 mg per 5 mL (as sodium phosphate); 2 mg per 5 mL (as sodium phosphate) Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule dexamethasone phosphate (as sodium phosphate, equivalent to 3.3 mg/mL dexamethasone base) Oral > Solid > tablet: 0.5 mg (as dexamethasone base); 0.75 mg (as dexamethasone base); 1.5 mg (as dexamethasone base); 4 mg (as dexamethasone base); 2 mg (as dexamethasone base)	<input type="text" value="Nausea or vomiting"/>

## Dolasetron

### Section

Antiemetic medicines

### Indications

Therapeutic equivalent to [ondansetron](#) for

Nausea or vomiting

## Granisetron

### Section

Antiemetic medicines

### Indications

Therapeutic equivalent to [ondansetron](#) for

Nausea or vomiting

## Hydrocortisone

### Section

Anti-inflammatory medicines

Local > Rectal > Suppository: 25 mg (acetate)

Local > Rectal > Retention enema: 100 mg per 60 mL

### Indications

Crohn disease site

Ulcerative colitis

## Mesalazine

### Section

Anti-inflammatory medicines

### Indications

Therapeutic equivalent to [sulfasalazine](#) for

Crohn disease site

Therapeutic equivalent to [sulfasalazine](#) for

Ulcerative colitis

## Metoclopramide

### Section

Antiemetic medicines

Oral > Liquid: 5 mg per 5 mL

Oral > Solid: 10 mg (scored) (hydrochloride)

Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)

### Indications

Nausea or vomiting

## Omeprazole

### Section

#### Antiulcer medicines

Parenteral > General injections > IV: 40 mg in vial

Oral > Liquid: 1 mg per mL powder for oral liquid; 4 mg per mL powder for oral liquid

Oral > Solid > dosage form: 10 mg; 20 mg; 40 mg

### Indications

Gastrointestinal bleeding

Peptic ulcer, site unspecified

Gastro-oesophageal reflux disease

## Ondansetron

### Section

#### Antiemetic medicines

Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (as hydrochloride dihydrate); 2 mg per mL in 4 mL ampoule

Oral > Liquid: 4 mg per 5 mL (as hydrochloride dihydrate)

Oral > Solid > dosage form: 4 mg (as hydrochloride dihydrate); 8 mg (as hydrochloride dihydrate); 24 mg (as hydrochloride dihydrate)

### Indications

Nausea or vomiting

## Oral rehydration salts

### Section

#### Oral rehydration

Powder for dilution in 200 mL, 500 mL, 1 L:

- glucose 75 mEq or mmol/L
- sodium 75 mEq or mmol/L
- chloride 65 mEq or mmol/L
- potassium 20 mEq or mmol/L
- citrate 10 mEq or mmol/L
- osmolarity 245 mOsm/L
- glucose 13.5 g/L
- sodium chloride 2.6 g/L
- potassium chloride 1.5 g/L
- trisodium citrate dihydrate 2.9 g/L

### Indications

Diarrhoea

## Oral rehydration salts - zinc sulfate

### Section

#### Medicines used in diarrhoea

Oral > Other: ORS powder for dilution and zinc sulfate dispersible tablet 20 mg (co-packaged)

### Indications

Diarrhoea

## Palonosetron

### Section

Antiemetic medicines

### Indications

Therapeutic equivalent to ondansetron for

Nausea or vomiting

## Pancreatic enzymes

### Section

Gastrointestinal medicines

Capsule (modified release): 10 000 lipase units + 8000 amylase units + 600 protease units; 25 000 lipase units + 18 000 amylase units + 1000 protease units.

\*Units expressed in Ph. Eur

### Indications

Enzyme intestinal

## Prednisolone

### Section

Anti-inflammatory medicines

Local > Rectal > Retention enema:

Local > Rectal > Retention enema: 20 mg per 100 mL (as sodium phosphate)

### Indications

Therapeutic equivalent to hydrocortisone for

Crohn disease site

Crohn disease site

Ulcerative colitis

## Promethazine

### Section

Antiemetic medicines

Oral > Liquid: 5 mg per 5 mL (hydrochloride)

Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)

Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule

### Indications

Nausea or vomiting

## Ranitidine

### Section

Antiulcer medicines

Oral > Liquid: 75 mg per 5 mL (as hydrochloride)

Oral > Solid: 150 mg (as hydrochloride)

Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (as hydrochloride)

### Indications

Gastro-oesophageal reflux disease

Peptic ulcer, site unspecified

## Senna

### Section

Laxatives

Oral > Solid: 7.5 mg (sennosides) (or traditional dosage forms)

### Indications

Constipation

## Sulfasalazine

### Section

Anti-inflammatory medicines

Oral > Solid: 500 mg

Local > Rectal > Suppository: 500 mg

Local > Rectal > Retention enema: 3 g per 100 mL

### Indications

Crohn disease site

Ulcerative colitis

## Tropisetron

### Section

Antiemetic medicines

### Indications

Therapeutic equivalent to ondansetron for

Nausea or vomiting

## Zinc sulfate

### Section

Medicines for diarrhoea

Oral > Solid > dispersible tablet: 20 mg (scored)

### Indications

Diarrhoea