

Alfacalcidol

Section	Indications
Medicines for endocrine disorders Oral > Liquid: 2 µg per mL Oral > Solid > capsule: 0.25 µg; 1 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

Bromocriptine

Section	Indications
Medicines for disorders of the pituitary hormone system	Therapeutic equivalent to cabergoline for Other specified benign neoplasm of endocrine glands

Cabergoline

Section	Indications
Medicines for disorders of the pituitary hormone system Oral > Solid > tablet: 0.5 mg; 1 mg	Other specified benign neoplasm of endocrine glands

Calcitriol

Section	Indications
Medicines for endocrine disorders Oral > Solid > capsule: 0.25 µg; 0.5 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

Canagliflozin

Section	Indications
Oral hypoglycaemic agents	Therapeutic equivalent to empagliflozin for Type 2 diabetes mellitus

Carbimazole

Section	Indications
Thyroid hormones and antithyroid medicines	Therapeutic equivalent to methimazole for Thyrotoxicosis

Dapagliflozin

Section

Oral hypoglycaemic agents

Indications

Therapeutic equivalent to empagliflozin for

Type 2 diabetes mellitus

Diazoxide

Section

Medicines for hypoglycaemia

Oral > Liquid: 50 mg per mL

Oral > Solid: 50 mg

Indications

Persistent hyperinsulinaemic hypoglycaemia of infancy

Empagliflozin

Section

Oral hypoglycaemic agents

Oral > Solid: 10 mg; 25 mg

Indications

Type 2 diabetes mellitus

Estradiol

Section

Estrogens

Oral > Solid > tablet: 0.5 mg; 1 mg; 2 mg

Indications

Gonadotropin deficiency

Hypopituitarism

Turner syndrome

Fludrocortisone

Section

Adrenal hormones and synthetic substitutes

Oral > Solid: 100 µg (acetate)

Indications

Adrenocortical insufficiency

Glibenclamide

Section

Oral hypoglycaemic agents

Oral > Solid: 5 mg; 2.5 mg

Indications

Type 2 diabetes mellitus

Gliclazide

Section

Oral hypoglycaemic agents

Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release

Indications

Type 2 diabetes mellitus

Glucagon

Section

Medicines for hypoglycaemia

Parenteral > General injections > unspecified: 1 mg per mL

Indications

Hypoglycaemia in the context of diabetes

Hydrocortisone

Section

Adrenal hormones and synthetic substitutes

Oral > Solid: 5 mg; 10 mg; 20 mg

Indications

Adrenocortical insufficiency

Insulin

Section

Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen (soluble)

Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

Insulin analogues

Section

Insulins

Not specified

Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

Insulin degludec

Section

Insulins

Parenteral > General injections > SC: 100 units per mL

Indications

Therapeutic equivalent to [insulin glargine](#) for

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues for

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues for

Type 2 diabetes mellitus

Insulin detemir

Section	Indications
Insulins	Therapeutic equivalent to insulin glargine for Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues for Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues for Type 2 diabetes mellitus

Insulin glargine

Section	Indications
Insulins Parenteral > General injections > SC: 100 units per mL	Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues for Type 1 diabetes mellitus Therapeutic equivalent to long-acting insulin analogues for Type 2 diabetes mellitus

Intermediate-acting insulin

Section	Indications
Insulins Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin	Type 1 diabetes mellitus Type 2 diabetes mellitus

Ketoconazole

Section	Indications
Medicines for endocrine disorders Oral > Solid > tablet: 200 mg	Cushing syndrome

Levothyroxine

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 µg (sodium salt); 100 µg (sodium salt); 25 µg (sodium salt)	Hypothyroidism

Liraglutide

Section

Medicines for endocrine disorders

Parenteral > General injections > SC: 6 mg per mL in 3 mL pre-filled pen

Indications

Obesity

Long-acting insulin analogues

Section

Insulins

Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

Lugol's solution

Section

Thyroid hormones and antithyroid medicines

Oral > Liquid: 13% w/v (about 130 mg/mL total iodine)

Indications

Thyrotoxicosis

Medroxyprogesterone acetate

Section

Progestogens

Oral > Solid: 5 mg

Indications

Excessive menstruation with irregular cycle

Metformin

Section

Oral hypoglycaemic agents

Oral > Solid: 500 mg (hydrochloride)

Indications

Type 2 diabetes mellitus

Methimazole

Section

Thyroid hormones and antithyroid medicines

Oral > Solid: 5 mg; 10 mg; 20 mg

Indications

Thyrotoxicosis

Norethisterone

Section

Progestogens

Oral > Solid: 5 mg

Indications

Excessive menstruation with irregular cycle

Therapeutic equivalent to medroxyprogesterone acetate for

Excessive menstruation with irregular cycle

Octreotide

Section

Medicines for disorders of the pituitary hormone system

Parenteral > General injections > IM: 20 mg in vial (modified-release, as acetate) plus diluent

Parenteral > General injections > SC: 0.05 mg per mL in 1 mL vial (immediate-release, as acetate); 0.1 mg per mL in 1 mL vial (immediate-release, as acetate); 0.5 mg per mL in 1 mL vial (immediate-release, as acetate)

Indications

Acromegaly or pituitary gigantism

Phosphorus

Section

Medicines for endocrine disorders

Oral > Other: 500 mg in sachet (elemental phosphorus) granules

Oral > Solid > tablet: 250 mg (elemental phosphorus); 500 mg (elemental phosphorus); 500 mg (elemental phosphorus) (effervescent)

Indications

Hypophosphataemic rickets

Potassium iodide

Section

Thyroid hormones and antithyroid medicines

Oral > Solid: 60 mg

Indications

Thyrotoxicosis

Propylthiouracil

Section

Thyroid hormones and antithyroid medicines

Oral > Solid: 50 mg

Indications

Thyrotoxicosis

Simvastatin

Section

Medicines for endocrine disorders

Oral > Solid: 20 mg

Indications

Polycystic ovary syndrome

Somatropin

Section

Medicines for endocrine disorders

Parenteral > General injections > SC: 5 mg in cartridge powder for injection; 5 mg per mL in 2 mL cartridge

Indications

Hypoglycaemia without associated diabetes

Testosterone

Section

Androgens

Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule (enantate)

Indications

Testicular dysfunction or testosterone-related disorders

Tolbutamide

Section

Oral hypoglycaemic agents

Oral > Solid: 500 mg

Indications

Type 2 diabetes mellitus

Zoledronic acid

Section

Medicines for endocrine disorders

Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle

Indications

Osteogenesis imperfecta