

Arsenic trioxide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion	Acute myeloid leukaemia with recurrent genetic abnormalities

Asparaginase

Section	Indications
Cytotoxic medicines Powder for injection: 10 000 IU in vial. Parenteral > General injections > unspecified: 10000 IU in vial powder for injection	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified

Azacitidine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection	Myeloid leukaemia

Bendamustine

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL	Chronic lymphocytic leukaemia or small lymphocytic lymphoma Follicular lymphoma

Bleomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)	Germ cell tumour of testis Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Unspecified malignant neoplasms of ill-defined or unspecified sites

Calcium folinate (leucovorin calcium)

Section

Cytotoxic medicines

Oral > Solid: 15 mg tablet

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Solid > tablet: 15 mg; 5 mg; 25 mg

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasm metastasis in large intestine

Malignant trophoblastic neoplasms of placenta

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of rectum

Malignant neoplasms of colon

Capecitabine

Section

Cytotoxic medicines

Oral > Solid > tablet: 150 mg; 500 mg

Indications

Other specified malignant neoplasms of breast

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of rectum

Malignant neoplasms of colon

Carboplatin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL

Indications

Other specified gliomas of brain

Other specified carcinomas of ovary

Retinoblastoma

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Malignant neoplasms of kidney, except renal pelvis

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Chlorambucil

Section

Cytotoxic medicines

Oral > Solid: 2 mg tablet

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Cisplatin

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection

Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Germ cell tumour of testis

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Cladribine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial

Indications

Langerhans cell histiocytosis

Cyclophosphamide

Section

Cytotoxic medicines

Oral > Solid: 25 mg tablet

Parenteral > General injections > unspecified: 500 mg in vial powder for injection

Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection

Oral > Solid > dosage form: 25 mg; 50 mg

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified malignant neoplasms of breast

Malignant trophoblastic neoplasms of placenta

Follicular lymphoma

Malignant neoplasms of breast

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Rhabdomyosarcoma primary site

Hodgkin lymphoma

Diffuse large B-cell lymphomas

Plasma cell myeloma

Other specified gliomas of brain

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of kidney, except renal pelvis

Ewing sarcoma of bone and articular cartilage of unspecified sites

Lymphoid leukaemia, not elsewhere classified

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Cytarabine

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 100 mg in vial powder for injection

Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Dacarbazine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection

Indications

Hodgkin lymphoma

Dactinomycin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 500 µg in vial powder for injection

Indications

Malignant trophoblastic neoplasms of placenta

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Ewing sarcoma of bone and articular cartilage of unspecified sites

Daunorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)

Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Docetaxel

Section

Cytotoxic medicines

Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL

Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Malignant neoplasms of prostate

Unspecified malignant neoplasms of ill-defined or unspecified sites

Doxorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)

Parenteral > General injections > IV: 10 mg in vial (hydrochloride) powder for injection; 50 mg in vial (hydrochloride) powder for injection; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Rhabdomyosarcoma primary site

Malignant neoplasms of breast

Hodgkin lymphoma

Diffuse large B-cell lymphomas

Plasma cell myeloma

Follicular lymphoma

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of kidney, except renal pelvis

Osteosarcoma of bone and articular cartilage of unspecified sites

Lymphoid leukaemia, not elsewhere classified

Anaplastic large cell lymphoma, ALK-negative

Other specified malignant neoplasms of breast

Anaplastic large cell lymphoma, ALK-positive

Ewing sarcoma of bone and articular cartilage of unspecified sites

Kaposi sarcoma of unspecified primary site

Doxorubicin (as pegylated liposomal)

Section

Cytotoxic medicines

Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Indications

Kaposi sarcoma of unspecified primary site

Etoposide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule Oral > Solid: 100 mg capsule Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate) Oral > Solid: 100 mg capsule; 50 mg capsule	Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Germ cell tumour of testis Malignant trophoblastic neoplasms of placenta Malignant neoplasms of kidney, except renal pelvis Osteosarcoma of bone and articular cartilage of unspecified sites Hodgkin lymphoma Retinoblastoma Anaplastic large cell lymphoma, ALK-positive Anaplastic large cell lymphoma, ALK-negative

Fludarabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection Oral > Solid: 10 mg tablet (fludarabine phosphate)	Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Fluorouracil

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule Parenteral > General injections > IV: 50 mg per mL in vial	Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of nasopharynx Malignant neoplasm metastasis in large intestine Malignant neoplasms of breast Malignant neoplasms of rectum Malignant neoplasms of colon

Gemcitabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection	Other specified carcinomas of ovary Other specified malignant neoplasms of bronchus or lung

Hydroxycarbamide (hydroxyurea)

Section	Indications
Cytotoxic medicines Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg	Squamous cell carcinoma of oropharynx Chronic myeloid leukaemia, not elsewhere classified

Ifosfamide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified malignant neoplasms of the ovary Germ cell tumour of testis Ewing sarcoma of bone and articular cartilage of unspecified sites Osteosarcoma of bone and articular cartilage of unspecified sites Rhabdomyosarcoma primary site Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Irinotecan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site

Melphalan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection Oral > Solid: 2 mg	Plasma cell myeloma

Mercaptopurine

Section	Indications
Cytotoxic medicines Oral > Solid: 50 mg tablet Oral > Liquid: 20 mg per mL	Unspecified malignant neoplasms of ill-defined or unspecified sites Acute myeloid leukaemia with recurrent genetic abnormalities Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Methotrexate

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (as sodium salt) Oral > Solid: 2.5 mg (as sodium salt) Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified Acute myeloid leukaemia with recurrent genetic abnormalities Malignant trophoblastic neoplasms of placenta Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis

Oxaliplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of colon

Paclitaxel

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 6 mg per mL	Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Kaposi sarcoma of unspecified primary site Other specified malignant neoplasms of breast Other specified carcinomas of ovary Malignant neoplasms of breast Malignant neoplasms of cervix uteri

Pegaspargase

Section

Cytotoxic medicines

Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection

Indications

Lymphoid leukaemia, not elsewhere classified

Procarbazine

Section

Cytotoxic medicines

Oral > Solid: 50 mg (as hydrochloride)

Oral > Solid > capsule: 50 mg (as hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Hodgkin lymphoma

Realgar-indigo naturalis formulation

Section

Cytotoxic medicines

Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)

Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

Tioguanine

Section

Cytotoxic medicines

Oral > Solid: 40 mg

Indications

Lymphoid leukaemia, not elsewhere classified

Vinblastine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate)

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Anaplastic large cell lymphoma, ALK-positive

Anaplastic large cell lymphoma, ALK-negative

Langerhans cell histiocytosis

Vincristine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate) Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Malignant trophoblastic neoplasms of placenta Follicular lymphoma Kaposi sarcoma of unspecified primary site Rhabdomyosarcoma primary site Hodgkin lymphoma Diffuse large B-cell lymphomas Retinoblastoma Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Vinorelbine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Oral > Solid > capsule: 20 mg; 30 mg; 80 mg	Other specified malignant neoplasms of breast Other specified malignant neoplasms of bronchus or lung Rhabdomyosarcoma primary site