

Acetylsalicylic acid

Section

Medicines for juvenile joint diseases
Oral > Solid: 100 to 500 mg
Local > Rectal > Suppository: 50 to 150 mg

Indications

Acute rheumatic fever without mention of heart involvement

Mucocutaneous lymph node syndrome

Juvenile idiopathic arthritis

Adalimumab

Section

Medicines for juvenile joint diseases
Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL;
10 mg per 0.2 mL; 20 mg per 0.4 mL

Indications

Juvenile idiopathic arthritis

Anakinra

Section

Medicines for juvenile joint diseases
Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe

Indications

Juvenile systemic arthritis

Certolizumab pegol

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Etanercept

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Golimumab

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Infliximab

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to adalimumab for

Juvenile idiopathic arthritis

Methotrexate

Section

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

Juvenile idiopathic arthritis

Tocilizumab

Section

Medicines for juvenile joint diseases

Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial

Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

Indications

Juvenile systemic arthritis

Triamcinolone hexacetonide

Section

Medicines for juvenile joint diseases

Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

Indications

Juvenile idiopathic arthritis