

Amikacin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate); 50 mg per mL in 2 mL vial (as sulfate)

Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)

Indications

Premier choix

Neutropenia (high-risk)

co-prescrire avec cefotaxime

Inflammatory and other diseases of prostate (severe)

co-prescrire avec cefotaxime

Acute pyelonephritis (severe)

co-prescrire avec ceftriaxone

Inflammatory and other diseases of prostate (severe)

co-prescrire avec ceftriaxone

Acute pyelonephritis (severe)

Second choix

co-prescrire avec cloxacillin

Sepsis without septic shock

Amoxicillin

Section

Access group antibiotics

Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid

Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection

Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)

Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate); 1 g (as trihydrate)

Indications

Premier choix

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Acute malnutrition in infants, children or adolescents (uncomplicated) [children]

Acute malnutrition in infants, children or adolescents (complicated) [children]

Acute pharyngitis

Chronic obstructive pulmonary disease with acute exacerbation

Periapical abscess without sinus

Acute otitis media

Acute sinusitis

co-prescrire avec gentamicin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec gentamicin

Sepsis without septic shock

Second choix

Bacterial meningitis

Autres indications

Infectious cystitis

Bacterial infection of unspecified site

Amoxicillin + clavulanic acid

Section

Access group antibiotics

Parenteral > General injections > IV: 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection

Oral > Liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt) / 5 mL powder for oral liquid; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid

Oral > Solid > dispersible tablet: 200 mg (as trihydrate) + 28.5 mg (as potassium salt); 250 mg (as trihydrate) + 62.5 mg (as potassium salt)

Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt)

Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)

Indications

Premier choix

Peritoneal abscess (mild-moderate)

Neutropenia (low-risk)

Other specified pneumonia (Hospital-acquired pneumonia)

Infectious cystitis

Chronic obstructive pulmonary disease with acute exacerbation

Peritonitis (mild-moderate)

Acute sinusitis

Bacterial cellulitis, erysipelas or lymphangitis

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Second choix

Osteomyelitis or osteitis

Acute otitis media

Other specified prophylactic measures

Bacterial infection of joint

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

co-prescrire avec clarithromycin

Bacterial pneumonia (Community-acquired pneumonia - severe)

Autres indications

Bacterial infection of unspecified site

Ampicillin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 1 g in vial (as sodium salt) powder for injection

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral suspension

Oral > Solid: 250 mg (anhydrous); 500 mg (anhydrous)

Indications

Premier choix

co-prescrire avec [gentamicin](#)

Peritoneal abscess (severe)

co-prescrire avec [gentamicin](#)

Sepsis without septic shock

co-prescrire avec [gentamicin](#)

Peritoneal abscess (mild-moderate)

co-prescrire avec [gentamicin](#)

Peritonitis (severe)

co-prescrire avec [gentamicin](#)

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec [gentamicin](#)

Peritonitis (mild-moderate)

co-prescrire avec [gentamicin](#)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Benzathine benzylpenicillin

Section

Access group antibiotics

Parenteral > General injections > IM: 1.2 million IU in vial (\approx 900 mg) powder for injection; 2.4 million IU in vial (\approx 1.8 g) powder for injection

Indications

Premier choix

Syphilis

Congenital syphilis [children]

Autres indications

Bacterial infection of unspecified site

Benzympenicillin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 600 mg in vial powder for injection (=1 million IU as sodium or potassium salt); 3 g in vial powder for injection (=5 million IU as sodium or potassium salt)

Parenteral > General injections > unspecified: 600 mg in vial powder for injection (= 1 million IU as sodium or potassium salt); 3 g in vial powder for injection (= 5 million IU as sodium or potassium salt)

Indications

Premier choix

Congenital syphilis [children]

Syphilis

co-prescrire avec gentamicin

Sepsis without septic shock

co-prescrire avec gentamicin

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec gentamicin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Cefalexin

Section

Access group antibiotics

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid: 250 mg (as monohydrate)

Oral > Solid: 250 mg (as monohydrate); 500 mg (as monohydrate)

Oral > Solid > dispersible tablet: 125 mg; 250 mg

Indications

Premier choix

Bacterial cellulitis, erysipelas or lymphangitis

Second choix

Acute pharyngitis

Chronic obstructive pulmonary disease with acute exacerbation

Autres indications

Bacterial infection of unspecified site

Cefazolin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 1 g in vial (as sodium salt) powder for injection

Indications

Premier choix

Other specified prophylactic measures

co-prescrire avec metronidazole

Other specified prophylactic measures

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Autres indications

Surgical site infection

Chloramphenicol

Section

Access group antibiotics

Parenteral > General injections > IV: 1 g in vial powder for injection (sodium succinate)

Parenteral > General injections > IM: 500 mg in 2 mL ampoule oily suspension for injection

Oral > Liquid: 150 mg per 5 mL (as palmitate)

Oral > Solid: 250 mg capsule

Indications

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Clindamycin

Section

Access group antibiotics

Oral > Liquid: 75 mg per 5 mL (as palmitate hydrochloride)

Oral > Solid: 150 mg (as hydrochloride) capsule

Parenteral > General injections > unspecified: 150 mg per mL (as phosphate) injection

Parenteral > General injections > IV: 150 mg per mL (as phosphate); 600 mg per 4 mL (as phosphate); 900 mg per 6 mL (as phosphate)

Oral > Liquid: 75 mg per 5 mL powder for oral liquid (as palmitate hydrochloride)

Oral > Solid: 150 mg (as hydrochloride)

Indications

Premier choix

co-prescrire avec piperacillin + tazobactam

Necrotising fasciitis

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Autres indications

Bacterial infection of unspecified site

Cloxacillin

Section

Access group antibiotics

Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid

Oral > Solid: 500 mg (as sodium salt) capsule; 1 g (as sodium salt) capsule

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection

Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid; 250 mg per 5 mL (as sodium salt) powder for oral liquid

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 250 mg in vial (as sodium salt) powder for injection

Oral > Solid > capsule: 500 mg (as sodium salt); 1 g (as sodium salt); 250 mg (as sodium salt)

Indications

Premier choix

Bacterial infection of joint

Osteomyelitis or osteitis

Bacterial cellulitis, erysipelas or lymphangitis

Second choix

co-prescrire avec amikacin

Sepsis without septic shock

Autres indications

Bacterial infection of unspecified site

Doxycycline

Section

Access group antibiotics

Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous)

Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate)

Parenteral > General injections > unspecified: 100 mg in vial powder for injection

Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium)

Oral > Solid > dispersible tablet: 100 mg (as monohydrate)

Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)

Oral > Solid > dosage form: 100 mg (as hyclate)

Indications

Premier choix

Chlamydia trachomatis

Cholera

Second choix

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Chronic obstructive pulmonary disease with acute exacerbation

Cholera [children]

Autres indications

Bacterial infection of unspecified site

Gentamicin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Indications

Premier choix

Neonatal meningitis

co-prescrire avec amoxicillin

Sepsis without septic shock

co-prescrire avec amoxicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec ampicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec ampicillin

Sepsis without septic shock

co-prescrire avec ampicillin

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec ampicillin

Peritoneal abscess (severe)

co-prescrire avec ampicillin

Peritonitis (mild-moderate)

co-prescrire avec ampicillin

Peritonitis (severe)

co-prescrire avec ampicillin

Peritoneal abscess (mild-moderate)

co-prescrire avec benzylpenicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec benzylpenicillin

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec benzylpenicillin

Sepsis without septic shock

Second choix

Other specified prophylactic measures

Gonococcal infection

Autres indications

Bacterial infection of unspecified site

Metronidazole

Section

Access group antibiotics

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Oral > Solid: 200 to 500 mg tablet

Parenteral > General injections > unspecified: 500 mg per 100 mL in vial

Injection

Local > Rectal > Suppository: 500 mg; 1 g

Parenteral > General injections > unspecified: 500 mg per 100 mL in vial

Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg

Indications

Premier choix

Intestinal infections due to *Clostridioides difficile*

Trichomoniasis

co-prescrire avec cefazolin

Other specified prophylactic measures

co-prescrire avec cefotaxime

Peritonitis (severe)

co-prescrire avec cefotaxime

Peritonitis (mild-moderate)

co-prescrire avec cefotaxime

Peritoneal abscess (mild-moderate)

co-prescrire avec cefotaxime

Peritoneal abscess (severe)

co-prescrire avec ceftriaxone

Necrotising fasciitis

co-prescrire avec ceftriaxone

Peritonitis (severe)

co-prescrire avec ceftriaxone

Peritoneal abscess (mild-moderate)

co-prescrire avec ceftriaxone

Peritoneal abscess (severe)

co-prescrire avec ceftriaxone

Peritonitis (mild-moderate)

Second choix

co-prescrire avec ciprofloxacin

Peritonitis (mild-moderate)

co-prescrire avec ciprofloxacin

Peritoneal abscess (mild-moderate)

Autres indications

Bacterial infection of unspecified site

Nitrofurantoin

Section

Access group antibiotics

Oral > Liquid: 25 mg per 5 mL oral liquid

Oral > Solid > dosage form: 100 mg; 50 mg

Indications

Premier choix

Infectious cystitis

Phenoxymethylpenicillin

Section

Access group antibiotics

Oral > Liquid: 250 mg per 5 mL (as potassium salt) powder for oral liquid

Oral > Solid: 250 mg (as potassium salt) tablet

Oral > Solid: 250 mg (as potassium salt); 500 mg (as potassium salt)

Indications

Premier choix

Acute pharyngitis

Periapical abscess without sinus

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Autres indications

Bacterial infection of unspecified site

Procaine benzylpenicillin

Section

Access group antibiotics

Parenteral > General injections > IM: 1 g in vial (=1 million IU) powder for injection; 3 g in vial (=3 million IU) powder for injection

Indications

Premier choix

Congenital syphilis [children]

Second choix

Syphilis

Spectinomycin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 2 g in vial (as hydrochloride) powder for injection

Indications

Second choix

Gonococcal infection

Autres indications

Bacterial infection of unspecified site

Sulfamethoxazole + trimethoprim

Section

Access group antibiotics

Parenteral > General injections > IV: 80 mg + 16 mg per mL in 10 mL ampoule; 80 mg + 16 mg per mL in 5 mL ampoule

Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid

Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet

Oral > Solid > dispersible tablet: 100 mg + 20 mg

Indications

Premier choix

Infectious cystitis

Second choix

Infectious gastroenteritis or colitis without specification of infectious agent

Autres indications

Bacterial infection of unspecified site

Tetracycline

Section

Access group antibiotics

Oral > Solid: 250 mg (hydrochloride)

Indications

Bacterial infection of unspecified site

Trimethoprim

Section

Access group antibiotics

Oral > Liquid: 50 mg per mL

Oral > Solid: 100 mg; 200 mg

Indications

Premier choix

Infectious cystitis