

Azithromycin

Section

Watch group antibiotics

Oral > Liquid: 200 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid > dosage form: 250 mg (anhydrous); 500 mg (anhydrous)

Indications

Premier choix

Chlamydia trachomatis

Typhoid fever

Yaws

Paratyphoid fever

Cholera

Cholera [children]

Trachoma

co-prescrire avec ceftriaxone

Gonococcal infection

Second choix

Infectious gastroenteritis or colitis without specification of infectious agent

co-prescrire avec cefixime

Gonococcal infection

Cefepime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection; 2 g in vial (as hydrochloride) powder for injection

Indications

Other specified bacterial diseases

Cefixime

Section

Watch group antibiotics

Oral > Liquid: 100 mg per 5 mL powder for oral liquid

Oral > Solid: 200 mg (as trihydrate); 400 mg (as trihydrate)

Indications

Second choix

Infectious gastroenteritis or colitis without specification of infectious agent

co-prescrire avec azithromycin

Gonococcal infection

Cefotaxime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)

Indications

Premier choix

Other specified pneumonia (Hospital-acquired pneumonia)

Bacterial meningitis

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec amikacin

Inflammatory and other diseases of prostate (severe)

co-prescrire avec amikacin

Acute pyelonephritis (severe)

co-prescrire avec clarithromycin

Bacterial pneumonia (Community-acquired pneumonia - severe)

co-prescrire avec metronidazole

Peritonitis (severe)

co-prescrire avec metronidazole

Peritonitis (mild-moderate)

co-prescrire avec metronidazole

Peritoneal abscess (severe)

co-prescrire avec metronidazole

Peritoneal abscess (mild-moderate)

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Inflammatory and other diseases of prostate (mild to moderate)

Acute pyelonephritis (mild to moderate)

Sepsis without septic shock

Autres indications

Bacterial infection of unspecified site

Ceftazidime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial (as pentahydrate) powder for injection; 1 g in vial (as pentahydrate) powder for injection

Indications

Premier choix

co-prescrire avec [vancomycin](#)

[Endophthalmitis](#)

Autres indications

[Pseudomonas aeruginosa resistant to other antibiotic](#)

Ceftriaxone

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)

Parenteral > Locoregional injections > Intravitreal: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 2 g in vial; 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)

Indications

Premier choix

[Typhoid fever](#)

[Bacterial pneumonia \(Community-acquired pneumonia - severe\) \[children\]](#)

[Paratyphoid fever](#)

[Other specified pneumonia \(Hospital-acquired pneumonia\)](#)

[Bacterial meningitis](#)

co-prescrire avec [amikacin](#)

[Acute pyelonephritis \(severe\)](#)

co-prescrire avec [amikacin](#)

[Inflammatory and other diseases of prostate \(severe\)](#)

co-prescrire avec [azithromycin](#)

[Gonococcal infection](#)

co-prescrire avec [clarithromycin](#)

[Bacterial pneumonia \(Community-acquired pneumonia - severe\)](#)

co-prescrire avec [metronidazole](#)

[Necrotising fasciitis](#)

co-prescrire avec [metronidazole](#)

[Peritonitis \(mild-moderate\)](#)

co-prescrire avec [metronidazole](#)

[Peritonitis \(severe\)](#)

co-prescrire avec [metronidazole](#)

[Peritoneal abscess \(mild-moderate\)](#)

co-prescrire avec [metronidazole](#)

[Peritoneal abscess \(severe\)](#)

co-prescrire avec [vancomycin](#)

Endophthalmitis

Second choix

Bacterial infection of joint

Osteomyelitis or osteitis

Infectious gastroenteritis or colitis without specification of infectious agent

Sepsis without septic shock

Inflammatory and other diseases of prostate (mild to moderate)

Acute pyelonephritis (mild to moderate)

Autres indications

Bacterial infection of unspecified site

Cefuroxime

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg in vial (as sodium salt) powder for injection; 750 mg in vial (as sodium salt) powder for injection; 1.5 g in vial (as sodium salt) powder for injection

Indications

Second choix

Other specified prophylactic measures

Ciprofloxacin

Section

Watch group antibiotics

Parenteral > General injections > IV: 2 mg per mL (as hyclate)

Oral > Liquid: 250 mg per 5 mL (anhydrous)

Oral > Solid: 250 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride)

Indications

Premier choix

Neutropenia (low-risk)

Inflammatory and other diseases of prostate (mild to moderate)

Typhoid fever

Paratyphoid fever

Infectious gastroenteritis or colitis without specification of infectious agent

Acute pyelonephritis (mild to moderate)

Second choix

Cholera

Cholera [children]

co-prescrire avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrire avec metronidazole

Peritonitis (mild-moderate)

Autres indications

Bacterial infection of unspecified site

Clarithromycin

Section

Watch group antibiotics

Oral > Liquid: 125 mg per 5 mL powder for oral liquid; 250 mg per 5 mL powder for oral liquid

Parenteral > General injections > unspecified: 500 mg in vial powder for injection

Oral > Solid > dosage form: 500 mg; 250 mg

Oral > Solid > dosage form: 500 mg

Indications

Premier choix

co-prescrire avec cefotaxime

Bacterial pneumonia (Community-acquired pneumonia - severe)

co-prescrire avec ceftriaxone

Bacterial pneumonia (Community-acquired pneumonia - severe)

Second choix

Acute pharyngitis

co-prescrire avec amoxicillin + clavulanic acid

Bacterial pneumonia (Community-acquired pneumonia - severe)

Autres indications

Helicobacter pylori associated gastric ulcer

Delafloxacin

Section

Watch group antibiotics

Parenteral > General injections > IV: 300 mg lyophilized powder for injection

Oral > Solid: 450 mg

Indications

Methicillin resistant *Staphylococcus aureus*

Erythromycin

Section

Watch group antibiotics

Parenteral > General injections > IV: 500 mg in vial powder for injection (as lactobionate)

Oral > Liquid: 125 mg per 5 mL (as stearate or ethylsuccinate)

Oral > Solid: 250 mg (as stearate or ethylsuccinate)

Indications

Second choix

Therapeutic equivalent to *clarithromycin* pour

Acute pharyngitis

Autres indications

Bacterial infection of unspecified site

Flomoxef

Section

Watch group antibiotics

Parenteral > General injections > IV: 0.5 g in vial powder for injection; 1 g in vial powder for injection

Indications

Other specified other antibiotic resistant *Enterobacterales*

Fourth generation cephalosporins

Section

Watch group antibiotics

.

Indications

Other specified bacterial diseases

Imipenem + cilastatin

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg (as monohydrate) + 250 mg (as sodium salt) powder for injection; 500 mg (as monohydrate) + 500 mg (as sodium salt) powder for injection

Indications

Second choix

Therapeutic equivalent to meropenem pour

Neutropenia (high-risk)

Therapeutic equivalent to meropenem pour

Peritoneal abscess (severe)

Therapeutic equivalent to meropenem pour

Peritonitis (severe)

Autres indications

Pseudomonas aeruginosa resistant to other antibiotic

Antibiotic resistant Acinetobacter baumannii

Meropenem

Section

Watch group antibiotics

Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection

Indications

Second choix

Neutropenia (high-risk)

Peritoneal abscess (severe)

Peritonitis (severe)

Neonatal meningitis

Piperacillin + tazobactam

Section

Watch group antibiotics

Parenteral > General injections > IV: 2 g (as sodium salt) + 250 mg (as sodium salt) powder for injection; 4 g (as sodium salt) + 500 mg (as sodium salt) powder for injection

Indications

Premier choix

Peritoneal abscess (severe)

Other specified pneumonia (Hospital-acquired pneumonia)

Neutropenia (high-risk)

Peritonitis (severe)

co-prescrire avec clindamycin

Necrotising fasciitis

Vancomycin

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection

Oral > Solid > capsule: 125 mg (as hydrochloride); 250 mg (as hydrochloride)

Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection

Parenteral > Locoregional injections > Intravitreal: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection

Indications

Premier choix

Necrotising fasciitis

co-prescrire avec ceftazidime

Endophthalmitis

co-prescrire avec ceftriaxone

Endophthalmitis

Second choix

Neutropenia (high-risk)

Intestinal infections due to Clostridioides difficile

Autres indications

Methicillin resistant Staphylococcus aureus