



# Liste modèle des médicaments essentiels

## Amikacin

Section	Indications
Antituberculosis medicines Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)	Multi-drug resistant Mycobacterium tuberculosis

## Amoxicillin + clavulanic acid

Section	Indications
Antituberculosis medicines Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	Multi-drug resistant Mycobacterium tuberculosis

## Bedaquiline

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg tablet; 20 mg tablet	Multi-drug resistant Mycobacterium tuberculosis

## Capreomycin

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

## Clofazimine

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg; 50 mg	Multi-drug resistant Mycobacterium tuberculosis

## Cycloserine

Section	Indications
Antituberculosis medicines Oral > Solid: 250 mg; 125 mg	Multi-drug resistant Mycobacterium tuberculosis

## Delamanid

Section	Indications
Antituberculosis medicines	Multi-drug resistant Mycobacterium tuberculosis
Oral > Solid > dispersible tablet: 25 mg	
Oral > Solid > tablet: 50 mg	

## Ethambutol

Section	Indications
Antituberculosis medicines	Other specified tuberculosis
PARENTERAL > General injections > IV: 1000 mg per 10 mL; 2000 mg per 20 mL	Tuberculosis
Oral > Solid > dispersible tablet: 100 mg	
Oral > Solid > tablet: 100 mg (hydrochloride); 400 mg (hydrochloride)	

## Ethambutol + isoniazid

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 400 mg + 150 mg tablet	

## Ethambutol + isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet	

## Ethambutol + isoniazid + rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 275 mg + 75 mg + 150 mg tablet	

## Ethionamide

Section	Indications
Antituberculosis medicines	Multi-drug resistant Mycobacterium tuberculosis
Oral > Solid > dispersible tablet: 125 mg	
Oral > Solid > tablet: 250 mg	Tuberculous meningitis

## Gatifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis

## Imipenem + cilastatin

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to <a href="#">meropenem</a> pour Multi-drug resistant Mycobacterium tuberculosis

## Isoniazid

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 100 mg per mL Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 100 mg; 300 mg	Other specified tuberculosis Tuberculosis

## Isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 50 mg + 150 mg + 75 mg	Tuberculosis

## Isoniazid + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 50 mg + 75 mg Oral > Solid > tablet: 75 mg + 150 mg; 150 mg + 300 mg	Tuberculosis

## Isoniazid + rifapentine

Section	Indications
Antituberculosis medicines Oral > Solid: 300 mg + 300 mg tablet (scored)	Latent tuberculosis

## Kanamycin (injection)

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

## Levofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 250 mg; 500 mg; 750 mg	Multi-drug resistant Mycobacterium tuberculosis

## Linezolid

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 600 mg	Multi-drug resistant Mycobacterium tuberculosis

## Meropenem

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Multi-drug resistant Mycobacterium tuberculosis

## Moxifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 400 mg	Multi-drug resistant Mycobacterium tuberculosis Tuberculosis

## Ofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis

## P-aminosalicylate sodium

Section	Indications
Antituberculosis medicines Oral > Liquid: 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid) powder for oral solution	Multi-drug resistant Mycobacterium tuberculosis

## P-aminosalicylic acid

Section	Indications
Antituberculosis medicines Oral > Solid: 4 g granules in sachet	Multi-drug resistant Mycobacterium tuberculosis

## Pretomanid

Section	Indications
Antituberculosis medicines Oral > Solid > tablet: 200 mg	Multi-drug resistant Mycobacterium tuberculosis

## Protonamide

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to ethionamide pour Multi-drug resistant Mycobacterium tuberculosis

## Pyrazinamide

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 400 mg; 500 mg	Tuberculosis

## Rifabutin

Section	Indications
Antituberculosis medicines Oral > Solid: 150 mg	Tuberculosis

## Rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Liquid: 20 mg per mL	Other specified tuberculosis
Oral > Solid: 150 mg; 300 mg	
PARENTERAL > General injections > IV: 600 mg in vial powder for injection	

## Rifapentine

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid > tablet: 150 mg; 300 mg	Latent tuberculosis

## Streptomycin (injection)

Section	Indications
Antituberculosis medicines PARENTERAL > General injections > IV: 1 g powder for injection (as sulfate) as vial	Multi-drug resistant Mycobacterium tuberculosis

## Terizidone

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to cycloserine pour Multi-drug resistant Mycobacterium tuberculosis