

**Atazanavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

**Atazanavir + ritonavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

**Darunavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

**Indinavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid: 400 mg (as sulfate)

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

**Lopinavir + ritonavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid > tablet: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg  
 tablet (heat-stable)  
 Oral > Solid > dosage form: 40 mg + 10 mg

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

**Nelfinavir**

## Section

Antiretrovirals > Protease inhibitors  
 Oral > Solid: 250 mg (as mesilate)  
 Oral > Other: 50 mg per g oral powder

## Indications

Human immunodeficiency virus disease  
 without mention of associated disease or  
 condition, clinical stage unspecified

## Ritonavir

### Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)

### Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

## Saquinavir

### Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 200 mg; 500 mg

### Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified