

Amphotericin B

Section	Indications
<p>Antileishmaniasis medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)</p>	Visceral leishmaniasis

Meglumine antimoniate

Section	Indications
<p>Antileishmaniasis medicines Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule</p>	Visceral leishmaniasis
	Cutaneous leishmaniasis
	Mucocutaneous leishmaniasis

Miltefosine

Section	Indications
<p>Antileishmaniasis medicines Oral > Solid: 10 mg; 50 mg</p>	Visceral leishmaniasis
	Mucocutaneous leishmaniasis
	Cutaneous leishmaniasis

Paromomycin

Section	Indications
<p>Antileishmaniasis medicines Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)</p>	Visceral leishmaniasis

Pentamidine

Section	Indications
<p>Antileishmaniasis medicines Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)</p>	Visceral leishmaniasis

Sodium stibogluconate

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial	Visceral leishmaniasis Mucocutaneous leishmaniasis Cutaneous leishmaniasis