

## Adalimumab

## Section

Immunomodulators for non-malignant disease  
 Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL;  
 10 mg per 0.2 mL; 20 mg per 0.4 mL

## Indications

Axial spondyloarthritis

Crohn disease site

Rheumatoid arthritis

## Azathioprine

## Section

Immunomodulators for non-malignant disease  
 Parenteral > General injections > IV: 50 mg in vial powder for injection (as sodium salt); 100 mg in vial powder for injection (as sodium salt)  
 Oral > Liquid: 10 mg per mL  
 Oral > Solid > tablet: 25 mg; 50 mg (scored)  
 Oral > Solid > tablet: 50 mg (scored)

## Indications

Failure or rejection of transplanted organs or tissues

Relapsing-remitting multiple sclerosis

## Certolizumab pegol

## Section

Immunomodulators for non-malignant disease

## Indications

Therapeutic equivalent to adalimumab pour

Axial spondyloarthritis

Therapeutic equivalent to adalimumab pour

Crohn disease site

Therapeutic equivalent to adalimumab pour

Rheumatoid arthritis

## Ciclosporin

## Section

Immunomodulators for non-malignant disease  
 Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection  
 Oral > Liquid: 100 mg per mL  
 Oral > Solid > capsule: 25 mg

## Indications

Failure or rejection of transplanted organs or tissues

## Etanercept

### Section

Immunomodulators for non-malignant disease

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) pour

Crohn disease site

Therapeutic equivalent to [adalimumab](#) pour

Rheumatoid arthritis

## Fingolimod

### Section

Immunomodulators for non-malignant disease

Oral > Solid: 0.25 mg; 0.5 mg

### Indications

Relapsing-remitting multiple sclerosis

## Golimumab

### Section

Immunomodulators for non-malignant disease

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) pour

Crohn disease site

Therapeutic equivalent to [adalimumab](#) pour

Rheumatoid arthritis

## Infliximab

### Section

Immunomodulators for non-malignant disease

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) pour

Crohn disease site

Therapeutic equivalent to [adalimumab](#) pour

Rheumatoid arthritis

## Methotrexate

### Section

Immunomodulators for non-malignant disease

Parenteral > General injections > SC: pre-filled syringe in multiple strengths

### Indications

Crohn disease site

Psoriatic arthritis

## Tacrolimus

### Section

Immunomodulators for non-malignant disease

Parenteral > General injections > IV: 5 mg per mL in 1 mL vial

Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension

Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)

### Indications

Failure or rejection of transplanted organs or tissues