

**Canagliflozin**

## Section

Oral hypoglycaemic agents

## Indications

Therapeutic equivalent to empagliflozin pour

Type 2 diabetes mellitus

**Dapagliflozin**

## Section

Oral hypoglycaemic agents

## Indications

Therapeutic equivalent to empagliflozin pour

Type 2 diabetes mellitus

**Empagliflozin**

## Section

Oral hypoglycaemic agents

Oral &gt; Solid: 10 mg; 25 mg

## Indications

Type 2 diabetes mellitus

**Glibenclamide**

## Section

Oral hypoglycaemic agents

Oral &gt; Solid: 5 mg; 2.5 mg

## Indications

Type 2 diabetes mellitus

**Gliclazide**

## Section

Oral hypoglycaemic agents

Oral &gt; Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release

## Indications

Type 2 diabetes mellitus

**Insulin**

## Section

Insulins

Parenteral &gt; General injections &gt; SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen (soluble)

## Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

## Insulin analogues

### Section

Insulins  
Not specified

### Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

## Insulin degludec

### Section

Insulins  
Parenteral > General injections > SC: 100 units per mL

### Indications

Therapeutic equivalent to insulin glargine pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 2 diabetes mellitus

## Insulin detemir

### Section

Insulins

### Indications

Therapeutic equivalent to insulin glargine pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 2 diabetes mellitus

## Insulin glargine

### Section

Insulins  
Parenteral > General injections > SC: 100 units per mL

### Indications

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 2 diabetes mellitus

## Intermediate-acting insulin

### Section

#### Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin

### Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

## Long-acting insulin analogues

### Section

#### Insulins

Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

### Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

## Metformin

### Section

#### Oral hypoglycaemic agents

Oral > Solid: 500 mg (hydrochloride)

### Indications

Type 2 diabetes mellitus

## Tolbutamide

### Section

#### Oral hypoglycaemic agents

Oral > Solid: 500 mg

### Indications

Type 2 diabetes mellitus