

**Beinaglutide**

## Section

Hypoglycaemic agents

## Indications

Therapeutic equivalent to [semaglutide](#) pour

Obesity

**Canagliflozin**

## Section

Oral hypoglycaemic agents

## Indications

Therapeutic equivalent to [empagliflozin](#) pour

Type 2 diabetes mellitus

**Dapagliflozin**

## Section

Oral hypoglycaemic agents

## Indications

Therapeutic equivalent to [empagliflozin](#) pour

Type 2 diabetes mellitus

**Dulaglutide**

## Section

Hypoglycaemic agents

## Indications

Therapeutic equivalent to [semaglutide](#) pour

Obesity

Therapeutic equivalent to [semaglutide](#) pour

Type 2 diabetes mellitus

**Empagliflozin**

## Section

Oral hypoglycaemic agents

Oral &gt; Solid: 10 mg; 25 mg

## Indications

Type 2 diabetes mellitus

**Exenatide**

## Section

Hypoglycaemic agents

## Indications

Therapeutic equivalent to [semaglutide](#) pour

Obesity

## Glibenclamide

### Section

Oral hypoglycaemic agents

Oral > Solid: 5 mg; 2.5 mg

### Indications

Type 2 diabetes mellitus

## Gliclazide

### Section

Oral hypoglycaemic agents

Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release

### Indications

Type 2 diabetes mellitus

## Insulin (analogue, long-acting)

### Section

Insulins

Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge solution; 100 IU per mL in 3 mL pre-filled pen solution; 100 IU per mL in 10 mL vial solution

### Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

## Insulin (analogue, rapid-acting)

### Section

Insulins

Parenteral > General injections > SC: 100 IU per 1 mL in 10 mL vial; 100 IU per 1 mL in 3 mL cartridge; 100 IU per 1 mL in 3 mL pre-filled pen

### Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

## Insulin (human, intermediate-acting)

### Section

Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin

### Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

## Insulin (human, short-acting)

### Section

Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial solution; 100 IU per mL in 10 mL vial solution; 100 IU per mL in 3 mL cartridge solution; 100 IU per mL in 3 mL pre-filled pen solution

### Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

## Insulin analogues

### Section

Insulins  
Not specified

### Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

## Insulin aspart

### Section

Insulins  
Parenteral > General injections > SC: 100 IU per mL in 110 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen  
Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

### Indications

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 2 diabetes mellitus

## Insulin degludec

### Section

Insulins  
Parenteral > General injections > SC: 100 units per mL

### Indications

Therapeutic equivalent to insulin glargine pour

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 2 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 1 diabetes mellitus

## Insulin detemir

### Section

Insulins

### Indications

Therapeutic equivalent to insulin glargine pour

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 2 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 1 diabetes mellitus

## Insulin glargine

### Section

#### Insulins

Parenteral > General injections > SC: 100 units per mL

### Indications

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 2 diabetes mellitus

Therapeutic equivalent to insulin (analogue, long-acting) pour

Type 1 diabetes mellitus

## Insulin glulisine

### Section

#### Insulins

Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

### Indications

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 2 diabetes mellitus

## Insulin lispro

### Section

#### Insulins

Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

### Indications

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 1 diabetes mellitus

Therapeutic equivalent to insulin (analogue, rapid-acting) pour

Type 2 diabetes mellitus

## Liraglutide

### Section

#### Hypoglycaemic agents

### Indications

Therapeutic equivalent to semaglutide pour

Obesity

Therapeutic equivalent to semaglutide pour

Type 2 diabetes mellitus

## Metformin

### Section

Oral hypoglycaemic agents

Oral > Solid: 500 mg (hydrochloride)

### Indications

Type 2 diabetes mellitus

## Semaglutide

### Section

Hypoglycaemic agents

Parenteral > General injections > SC: 0.25 mg in pre-filled pen injection solution; 0.5 mg in pre-filled pen injection solution; 1 mg in pre-filled pen injection solution; 2 mg in pre-filled pen injection solution; 4 mg in pre-filled pen injection solution

Oral > Solid > tablet: 3 mg

Parenteral > General injections > SC: 0.68 mg per mL; 1.34 mg per mL; 2.68 mg per mL

### Indications

Obesity

Type 2 diabetes mellitus

## Tirzepatide

### Section

Hypoglycaemic agents

### Indications

Therapeutic equivalent to [semaglutide](#) pour

Obesity

Therapeutic equivalent to [semaglutide](#) pour

Type 2 diabetes mellitus

## Tolbutamide

### Section

Oral hypoglycaemic agents

Oral > Solid: 500 mg

### Indications

Type 2 diabetes mellitus