

Acamprosate calcium

Section

Medicines for alcohol use disorders
Oral > Solid > tablet: 333 mg

Indications

Disorders due to use of alcohol

Buprenorphine

Section

Medicines for opioid use disorders

Indications

Therapeutic equivalent to methadone pour

Opioid dependence

Bupropion

Section

Medicines for nicotine use disorders
Oral > Solid: 150 mg sustained-release (hydrochloride)

Indications

Nicotine dependence

Methadone

Section

Medicines for opioid use disorders
Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

Indications

Opioid dependence

Naltrexone

Section

Medicines for alcohol use disorders
Parenteral > General injections > IM: 380 mg in vial (extended-release)
Oral > Solid > tablet: 50 mg

Indications

Disorders due to use of alcohol

Nicotine replacement therapy

Section

Medicines for nicotine use disorders

Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour

Oral > Solid > lozenge: 2 mg; 4 mg lozenge

Local > Buccal > Oral spray: 1 mg per actuation

Local > Buccal > Chewing gum: 2 mg; 4 mg

Indications

Nicotine dependence

Varenicline

Section

Medicines for nicotine use disorders

Oral > Solid: 0.5 mg; 1 mg

Indications

Nicotine dependence