

**Acamprosate calcium**

## Section

Medicines for alcohol use disorders  
Oral > Solid > tablet: 333 mg

## Indications

Disorders due to use of alcohol

**Buprenorphine**

## Section

Medicines for opioid use disorders

## Indications

Therapeutic equivalent to methadone pour

Opioid dependence

**Bupropion**

## Section

Medicines for nicotine use disorders  
Oral > Solid: 150 mg sustained-release (hydrochloride)

## Indications

Nicotine dependence

**Methadone**

## Section

Medicines for opioid use disorders  
Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

## Indications

Opioid dependence

**Naltrexone**

## Section

Medicines for alcohol use disorders  
Parenteral > General injections > IM: 380 mg in vial (extended-release)  
Oral > Solid > tablet: 50 mg

## Indications

Disorders due to use of alcohol

## Nicotine replacement therapy

### Section

#### Medicines for nicotine use disorders

Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour

Oral > Solid > lozenge: 2 mg; 4 mg lozenge

Local > Buccal > Oral spray: 1 mg per actuation

Local > Buccal > Chewing gum: 2 mg; 4 mg

### Indications

Nicotine dependence

## Varenicline

### Section

#### Medicines for nicotine use disorders

Oral > Solid: 0.5 mg; 1 mg

### Indications

Nicotine dependence