

**Acetylsalicylic acid**

## Section

Medicines for juvenile joint diseases  
 Oral > Solid: 100 to 500 mg  
 Local > Rectal > Suppository: 50 to 150 mg

## Indications

Acute rheumatic fever without mention of heart involvement

Mucocutaneous lymph node syndrome

Juvenile idiopathic arthritis

**Adalimumab**

## Section

Medicines for juvenile joint diseases  
 Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL;  
 10 mg per 0.2 mL; 20 mg per 0.4 mL

## Indications

Juvenile idiopathic arthritis

**Allopurinol**

## Section

Medicines used to treat gout  
 Oral > Solid: 100 mg

## Indications

Gout

**Anakinra**

## Section

Medicines for juvenile joint diseases  
 Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe

## Indications

Juvenile systemic arthritis

**Azathioprine**

## Section

Disease-modifying anti-rheumatic drugs (DMARDs)  
 Oral > Solid > tablet: 50 mg (scored)

## Indications

Rheumatoid arthritis, serology unspecified

Psoriatic arthritis

**Certolizumab pegol**

## Section

Medicines for juvenile joint diseases

## Indications

Therapeutic equivalent to adalimumab pour

Juvenile idiopathic arthritis

## Chloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

### Indications

Rheumatoid arthritis

## Colchicine

### Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

### Indications

Gout

## Etanercept

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Golimumab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Hydroxychloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

### Indications

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

## Infliximab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Methotrexate

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

### Indications

Rheumatoid arthritis, serology unspecified

Juvenile idiopathic arthritis

## Penicillamine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 250 mg

### Indications

Rheumatoid arthritis, serology unspecified

## Probenecid

### Section

Medicines used to treat gout

Oral > Solid: 500 mg

### Indications

Gout

## Sulfasalazine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 500 mg

### Indications

Rheumatoid arthritis, serology unspecified

## Tocilizumab

### Section

Medicines for juvenile joint diseases

Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial

Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

### Indications

Juvenile systemic arthritis

## Triamcinolone hexacetonide

### Section

Medicines for juvenile joint diseases

Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

### Indications

Juvenile idiopathic arthritis