



Liste modèle des médicaments essentiels

BCG vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against tuberculosis

Japanese encephalitis vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Medicines for COVID-19

Section	Indications
Medicines for COVID-19 Refer to WHO living guidelines	COVID-19

Abacavir

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 300 mg tablet (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abacavir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abacavir + lamivudine + lopinavir + ritonavir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 15 mg + 40 mg + 10 mg capsule containing oral granules	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abemaciclib

Section	Indications
Targeted therapies	Therapeutic equivalent to palbociclib pour Other specified malignant neoplasms of breast

Abiraterone

Section	Indications
Hormones and antihormones Oral > Solid: 250 mg; 500 mg	Malignant neoplasms of prostate

Acamprostate calcium

Section	Indications
Medicines for alcohol use disorders Oral > Solid > tablet: 333 mg	Disorders due to use of alcohol

Acetazolamide

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Oral > Solid: 250 mg	Primary open-angle glaucoma

Acetic acid

Section	Indications
Ear, nose and throat medicines [c] Local > Otological > drops: 2% in alcohol	Infectious diseases of external ear

Acetylcysteine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule Oral > Liquid: 10%; 20%	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs Acute or subacute hepatic failure

Acetylsalicylic acid

Section	Indications
Antimigraine medicines > For treatment of acute attack Oral > Solid: 300 to 500 mg	Migraine
Anti-platelet medicines Oral > Solid: 100 mg	Acute myocardial infarction Cerebral ischaemic stroke
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Pain

Acetylsalicylic acid + atorvastatin + ramipril

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Aciclovir

Section	Indications
Antiherpes medicines	Zoster
Parenteral > General injections > IV: 250 mg in vial powder for injection (as sodium salt)	Varicella
Oral > Liquid: 200 mg per 5 mL	Herpes simplex infections
Oral > Solid: 200 mg tablet	
Ophthalmological preparations > Anti-infective agents	Herpes simplex keratitis
Local > Ophthalmological > Ointment: 3% w/w	

Aclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Activated charcoal

Section	Indications
Antidotes and other substances used in poisonings > Non-specific Oral > Other: powder	Unspecified injury, poisoning or certain other consequences of external causes

Adalimumab

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Axial spondyloarthritis Crohn disease site Rheumatoid arthritis
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Afatinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib pour Other specified malignant neoplasms of bronchus or lung

Albendazole

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 400 mg (chewable, scored)	Echinococcosis Enterobiasis Trichuriasis Hookworm diseases Cysticercosis Ascariasis
Cysticidal medicines Oral > Solid > tablet: 400 mg (chewable, scored); 200 mg (chewable)	Other specified echinococcosis Cysticercosis of central nervous system
Antifilarials Oral > Solid > tablet: 400 mg (chewable, scored)	Lymphatic filariasis

Alcohol based hand rub

Section	Indications
Disinfectants Local > Topical > Solution: 80% v/v ethanol; 75% v/v isopropyl alcohol	Denatured alcohol

Alcuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 5 mg per mL in 2 mL ampoule (alcuronium chloride)	Muscle relaxants

Alfacalcidol

Section	Indications
Medicines for endocrine disorders Oral > Liquid: 2 µg per mL Oral > Solid > capsule: 0.25 µg; 1 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

All-trans retinoic acid

Section	Indications
Targeted therapies Oral > Solid: 10 mg capsule	Acute myeloid leukaemia with recurrent genetic abnormalities

Allopurinol

Section	Indications
Medicines used to treat gout Oral > Solid: 100 mg	Gout
Supportive medicines Oral > Solid: 100 mg; 300 mg	Tumour lysis syndrome

Alteplase

Section	Indications
Thrombolytic medicines Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection	Cerebral ischaemic stroke

Amidotrizoate

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)	Amidotrizoate

Amikacin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis
Antituberculosis medicines	Therapeutic equivalent to gentamicin pour Infectious blepharitis
Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)	Multi-drug resistant Mycobacterium tuberculosis
Access group antibiotics	Premier choix
Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate); 50 mg per mL in 2 mL vial (as sulfate)	Neutropenia (high-risk) co-prescrite avec cefotaxime
Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)	Inflammatory and other diseases of prostate (severe) co-prescrite avec cefotaxime
	Acute pyelonephritis (severe) co-prescrite avec ceftriaxone
	Inflammatory and other diseases of prostate (severe) co-prescrite avec ceftriaxone
	Acute pyelonephritis (severe) co-prescrite avec cloxacillin
	Sepsis without septic shock

Amiloride

Section	Indications
Diuretics	Oedema Ascites

Amiodarone

Section	Indications
Antiarrhythmic medicines	Cardiac arrhythmia
Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride)	
Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)	

Amitriptyline

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 25 mg; 10 mg; 75 mg	Palliative care
Medicines for depressive disorders Oral > Solid: 25 mg; 75 mg	Depressive disorders

Amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 5 mg (as maleate, mesylate or besylate)	Essential hypertension

Amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	Malaria due to Plasmodium falciparum
	Malaria due to Plasmodium vivax

Amodiaquine + sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	Malaria

Amoxicillin

Section	Indications
Access group antibiotics	
Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid	Premier choix
Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection	Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)
Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)	Acute malnutrition in infants, children or adolescents (uncomplicated) [children]
Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)	Acute malnutrition in infants, children or adolescents (complicated) [children]
Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid	Acute pharyngitis
Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate); 1 g (as trihydrate)	Chronic obstructive pulmonary disease with acute exacerbation
	Periapical abscess without sinus
	Acute otitis media
	Acute sinusitis
	co-prescrite avec gentamicin
	Bacterial pneumonia (Community-acquired pneumonia - severe) [children]
	co-prescrite avec gentamicin
	Sepsis without septic shock
	Second choix
	Bacterial meningitis
	Autres indications
	Infectious cystitis
	Bacterial infection of unspecified site

Amoxicillin + clavulanic acid

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection Oral > Liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt) / 5 mL powder for oral liquid; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid Oral > Solid > dispersible tablet: 200 mg (as trihydrate) + 28.5 mg (as potassium salt); 250 mg (as trihydrate) + 62.5 mg (as potassium salt) Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt) Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	Premier choix Peritoneal abscess (mild-moderate) Neutropenia (low-risk) Other specified pneumonia (Hospital-acquired pneumonia) Infectious cystitis Chronic obstructive pulmonary disease with acute exacerbation Peritonitis (mild-moderate) Acute sinusitis Bacterial cellulitis, erysipelas or lymphangitis Bacterial pneumonia (Community-acquired pneumonia - severe) [children] Second choix Osteomyelitis or osteitis Acute otitis media Other specified prophylactic measures Bacterial infection of joint Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) co-prescrite avec clarithromycin Bacterial pneumonia (Community-acquired pneumonia - severe)
Antituberculosis medicines Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid Oral > Solid > tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	Autres indications Bacterial infection of unspecified site Multi-drug resistant Mycobacterium tuberculosis

Amphotericin B

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	Blastomycosis Cryptococcosis Candidosis Coccidioidomycosis Histoplasmosis Sporotrichosis Aspergillosis
Antileishmaniasis medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	Visceral leishmaniasis

Ampicillin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 1 g in vial (as sodium salt) powder for injection Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral suspension Oral > Solid: 250 mg (anhydrous); 500 mg (anhydrous)	Premier choix co-prescrite avec gentamicin Peritoneal abscess (severe) co-prescrite avec gentamicin Sepsis without septic shock co-prescrite avec gentamicin Peritoneal abscess (mild-moderate) co-prescrite avec gentamicin Peritonitis (severe) co-prescrite avec gentamicin Acute malnutrition in infants, children or adolescents (complicated) [children] co-prescrite avec gentamicin Peritonitis (mild-moderate) co-prescrite avec gentamicin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] Second choix Bacterial meningitis Autres indications Bacterial infection of unspecified site

Anakinra

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

Anastrozole

Section	Indications
Hormones and antihormones Oral > Solid: 1 mg tablet	Other specified malignant neoplasms of breast Malignant neoplasms of breast

Anidulafungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin pour Systemic or invasive candidosis

Ansuvimab

Section	Indications
Medicines for Ebola virus disease Parenteral > General injections > IV: 400 mg powder for injection	Ebola virus disease

Anti-D immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IV: 250 µg in single-dose vial Parenteral > General injections > IM: 250 µg in single-dose vial	Rh incompatibility reaction Maternal care for red cell antibodies

Anti-rabies immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 150 IU per mL in vial Parenteral > Locoregional injections > Other: 150 IU per mL in vial	Rabies

Anti-rabies virus monoclonal antibodies

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > Locoregional injections > Intradermal: 40 IU per mL in 1.25 mL vial (human); 40 IU per mL in 2.5 mL vial (human); 100 IU per mL in 2.5 mL vial (human); 300 IU per mL in 10 mL vial (murine); 600 IU per mL in 1 mL vial (murine); 600 IU per mL in 2.5 mL vial (murine); 600 IU per mL in 5 mL vial (murine)	Rabies

Anti-tetanus immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 500 IU in vial	Tetanus

Antirabies hyperimmune serum

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > unspecified: 1000 IU in 5 mL ampoule	Rabies

Antivenom immunoglobulin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IV:	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Antivenom sera

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IV:	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Apixaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Aprepitant

Section	Indications
Antiemetic medicines Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid: 80 mg; 125 mg; 165 mg	Nausea or vomiting

Aripiprazole

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders

Arsenic trioxide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion	Acute myeloid leukaemia with recurrent genetic abnormalities

Artemether

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artemether + lumefantrine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution Oral > Solid: 50 mg tablet Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + mefloquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + pyronaridine tetraphosphate

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Ascorbic acid

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg	Scurvy

Asparaginase

Section	Indications
Cytotoxic medicines Powder for injection: 10 000 IU in vial.	Unspecified malignant neoplasms of ill-defined or unspecified sites
Parenteral > General injections > unspecified: 10000 IU in vial powder for injection	Lymphoid leukaemia, not elsewhere classified

Atazanavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atazanavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atenolol

Section	Indications
Antianginal medicines Oral > Solid: 50 mg; 100 mg	Angina pectoris
Antihypertensive medicines Oral > Solid: 50 mg; 100 mg	Essential hypertension Therapeutic equivalent to bisoprolol pour Essential hypertension

Atezolizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion	Other specified malignant neoplasms of bronchus or lung Therapeutic equivalent to pembrolizumab pour Other specified malignant neoplasms of bronchus or lung

Atoltivimab + maftivimab + odesivimab

Section	Indications
Medicines for Ebola virus disease Parenteral > General injections > IV: 241.7 mg + 241.7 mg + 241.7 mg in 14.5 mL vial	Ebola virus disease

Atorvastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg	Chronic ischaemic heart disease Atherosclerotic chronic arterial occlusive disease Cerebral ischaemic stroke

Atracurium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg per mL (besylate)	Muscle relaxants Therapeutic equivalent to vecuronium pour Muscle relaxants

Atropine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	Exposure to or harmful effects of undetermined intent of pesticides
Preoperative medication and sedation for short-term procedures Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	Parasympatholytics [anticholinergics and antimuscarinics] and spasmolytics
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour Cycloplegic drug
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 0.1% (sulfate); 0.5% (sulfate); 1% (sulfate)	Anterior uveitis

Axicabtagene ciloleucel

Section	Indications
Antineoplastics and supportive medicines Cell suspension for infusion: 2×10^6 anti-CD19 CAR-positive viable T-cells per kg of body weight.	Diffuse large B-cell lymphomas

Azacitidine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection	Myeloid leukaemia

Azathioprine

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg in vial powder for injection (as sodium salt); 100 mg in vial powder for injection (as sodium salt) Oral > Liquid: 10 mg per mL Oral > Solid > tablet: 25 mg; 50 mg (scored) Oral > Solid > tablet: 50 mg (scored) Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 50 mg (scored)	Failure or rejection of transplanted organs or tissues Relapsing-remitting multiple sclerosis Rheumatoid arthritis, serology unspecified Psoriatic arthritis

Azithromycin

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 1.5% eye drops	Trachoma
Watch group antibiotics Oral > Liquid: 200 mg per 5 mL (anhydrous) powder for oral liquid Oral > Solid > dosage form: 250 mg (anhydrous); 500 mg (anhydrous)	Premier choix Chlamydia trachomatis Typhoid fever Yaws Paratyphoid fever Cholera Cholera [children] Trachoma co-prescrite avec ceftriaxone Gonococcal infection Second choix Infectious gastroenteritis or colitis without specification of infectious agent co-prescrite avec cefixime Gonococcal infection

Aztreonam

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified bacterial diseases

Barium sulfate

Section	Indications
Diagnostic agents > Radiocontrast media Oral > Liquid: Aqueous suspension	Barium sulfate with suspending agents

Beclometasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 50 µg per dose (dipropionate); 250 µg per dose (dipropionatae)	Therapeutic equivalent to budesonide pour Asthma Asthma

Beclometasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Bedaquiline

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg tablet; 20 mg tablet	Multi-drug resistant Mycobacterium tuberculosis

Bendamustine

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL	Chronic lymphocytic leukaemia or small lymphocytic lymphoma Follicular lymphoma

Benzathine benzylpenicillin

Section	Indications
Access group antibiotics Parenteral > General injections > IM: 1.2 million IU in vial (\approx 900 mg) powder for injection; 2.4 million IU in vial (\approx 1.8 g) powder for injection	Premier choix Syphilis Congenital syphilis [children]
	Autres indications Bacterial infection of unspecified site

Benznidazole

Section	Indications
American trypanosomiasis Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg	Chagas disease

Benzoic acid + salicylic acid

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Cream: 6% + 3% Local > Topical > Ointment: 6% + 3%	Fungal infection of the skin

Benzoyl peroxide

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Cream: 5% Local > Topical > Lotion: 5%	Acne

Benzyl benzoate

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Lotion: 25%	Pthiriasis Pediculosis corporis Pediculosis capitis Scabies

Benzylpenicillin

Section	Indications
Access group antibiotics	Premier choix
Parenteral > General injections > unspecified: 600 mg in vial powder for injection (=1 million IU as sodium or potassium salt); 3 g in vial powder for injection (=5 million IU as sodium or potassium salt)	Congenital syphilis [children] Syphilis co-prescrite avec gentamicin
Parenteral > General injections > unspecified: 600 mg in vial powder for injection (= 1 million IU as sodium or potassium salt); 3 g in vial powder for injection (= 5 million IU as sodium or potassium salt)	Sepsis without septic shock co-prescrite avec gentamicin
	Acute malnutrition in infants, children or adolescents (complicated) [children]
	co-prescrite avec gentamicin
	Bacterial pneumonia (Community-acquired pneumonia - severe) [children]
	Second choix
	Bacterial meningitis
	Autres indications
	Bacterial infection of unspecified site

Betamethasone

Section	Indications
Dermatological medicines > Anti-inflammatory and antipruritic medicines	
Local > Topical > Cream: 0.1% (as valerate)	Pruritus due to skin disorder
Local > Topical > Ointment: 0.1% (as valerate)	Psoriasis of unspecified type
	Dermatitis or eczema
	Lichen planus

Bevacizumab

Section	Indications
Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations	Age related macular degeneration
Parenteral > Locoregional injections > Intravitreal: 25 mg per mL	

Bicalutamide

Section	Indications
Hormones and antihormones	
Oral > Solid: 50 mg	Malignant neoplasms of prostate

Binimetinib

Section	Indications
Targeted therapies	Therapeutic equivalent to trametinib pour Other specified melanoma of skin

Biperiden

Section	Indications
Medicines for parkinsonism Parenteral > General injections > IV: 5 mg per mL in 1 mL ampoule (lactate) Oral > Solid: 2 mg tablet (hydrochloride)	Parkinson disease

Bisacodyl

Section	Indications
Laxatives	Therapeutic equivalent to senna pour Constipation

Bisoprolol

Section	Indications
Antianginal medicines Oral > Solid: 1.25 mg; 5 mg	Angina pectoris
Medicines used in heart failure Oral > Solid: 1.25 mg; 5 mg	Heart failure
Antihypertensive medicines Oral > Solid: 1.25 mg; 5 mg	Essential hypertension
Antiarrhythmic medicines Oral > Solid: 1.25 mg; 5 mg	Cardiac arrhythmia

Bleomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)	Germ cell tumour of testis Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Unspecified malignant neoplasms of ill-defined or unspecified sites

Bortezomib

Section	Indications
Targeted therapies Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection	Plasma cell myeloma

Bromocriptine

Section	Indications
Antiseizure medicines Oral > Solid: 2.5 mg; 5 mg	Parkinson disease
Medicines for disorders of the pituitary hormone system	Therapeutic equivalent to cabergoline pour Other specified benign neoplasm of endocrine glands

Budesonide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 100 µg per dose; 200 µg per dose	Asthma
Ear, nose and throat medicines [c] Local > Nasal > Spray: 100 µg per dose	Allergic rhinitis

Budesonide + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > dry powder: 100 µg + 6 µg per dose; 200 µg + 6 µg per dose	Asthma

Budesonide + salmeterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Bumetanide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure
Diuretics	Therapeutic equivalent to furosemide pour Anuria or oliguria
	Therapeutic equivalent to furosemide pour Oedema

Bupivacaine

Section	Indications
Local anaesthetics	Local anaesthetics
Parenteral > Locoregional injections > Spinal anaesthesia: 0.5% in 4 mL ampoule (hydrochloride) + 7.5% glucose solution	
Parenteral > General injections > unspecified: 0.25% in vial (hydrochloride); 0.5% in vial (hydrochloride)	

Buprenorphine

Section	Indications
Medicines for opioid use disorders	Therapeutic equivalent to methadone pour Opioid dependence

Bupropion

Section	Indications
Medicines for nicotine use disorders	Nicotine dependence

Oral > Solid: 150 mg sustained-release (hydrochloride)

C1 esterase inhibitor

Section	Indications
Plasma-derived medicines Parenteral > General injections > IV: 500 IU in vial powder for injection	Hereditary angioedema

Cabergoline

Section	Indications
Medicines for disorders of the pituitary hormone system Oral > Solid > tablet: 0.5 mg; 1 mg	Other specified benign neoplasm of endocrine glands
Antiseizure medicines Oral > Solid > tablet: 0.5 mg	Parkinson disease

Caffeine citrate

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL) Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)	Apnoea of newborn

Calamine

Section	Indications
Dermatological medicines > Anti-inflammatory and antipruritic medicines Local > Topical > Lotion:	Pruritus

Calcipotriol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Cream: 50 µg per mL (0.005%) Local > Topical > Lotion: 50 µg per mL (0.005%) Local > Topical > Ointment: 50 µg per mL (0.005%)	Psoriasis of unspecified type

Calcitriol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to calcipotriol pour Psoriasis of unspecified type
Medicines for endocrine disorders	Chronic kidney disease, stage unspecified
Oral > Solid > capsule: 0.25 µg; 0.5 µg	Disorders of vitamin D metabolism or transport

Calcium

Section	Indications
Vitamins and minerals Oral > Solid: 500 mg (elemental calcium)	Calcium deficiency

Calcium folinate (leucovorin calcium)

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Oral > Solid: 15 mg tablet	Burkitt lymphoma including Burkitt leukaemia
Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule	Malignant neoplasm metastasis in large intestine
Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule	Malignant trophoblastic neoplasms of placenta
Oral > Solid > tablet: 15 mg; 5 mg; 25 mg	Osteosarcoma of bone and articular cartilage of unspecified sites
	Malignant neoplasms of rectum
	Malignant neoplasms of colon

Calcium gluconate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Hypermagnesaemia
Vitamins and minerals Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Tetany due to acute calcium deficiency

Canagliflozin

Section	Indications
Oral hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Capecitabine

Section	Indications
Cytotoxic medicines Oral > Solid > tablet: 150 mg; 500 mg	Other specified malignant neoplasms of breast Malignant neoplasm metastasis in large intestine Malignant neoplasms of rectum Malignant neoplasms of colon

Capreomycin

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

Captopril

Section	Indications
Medicines used in heart failure Oral > Solid: 25 mg	Heart failure

Carbachol

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines	Therapeutic equivalent to pilocarpine pour Acute angle closure with pupillary block Therapeutic equivalent to pilocarpine pour Ocular hypertension Therapeutic equivalent to pilocarpine pour Primary open-angle glaucoma

Carbamazepine

Section	Indications
Medicines for bipolar disorders Oral > Solid > tablet: 100 mg (scored); 200 mg (scored); 400 mg	Bipolar or related disorders
Antiseizure medicines Oral > Liquid: 100 mg per 5 mL Oral > Solid > tablet: 100 mg (scored); 200 mg (scored); 100 mg (chewable); 200 mg (chewable); 400 mg (scored)	Epilepsy or seizures

Carbetocin

Section	Indications
Uterotonics Parenteral > General injections > IV: 100 µg per mL (heat stable)	Postpartum haemorrhage

Carbimazole

Section	Indications
Thyroid hormones and antithyroid medicines	Therapeutic equivalent to methimazole pour Thyotoxicosis

Carboplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL	Other specified gliomas of brain Other specified carcinomas of ovary Retinoblastoma Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Malignant neoplasms of kidney, except renal pelvis Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of lip, oral cavity or pharynx Malignant neoplasms of cervix uteri Other specified malignant neoplasms of the ovary Germ cell tumour of testis

Carmellose

Section	Indications
Ophthalmological preparations	Therapeutic equivalent to hypromellose pour Keratoconjunctivitis sicca

Carvedilol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Caspofungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin pour Systemic or invasive candidosis

Cefalexin

Section	Indications
Access group antibiotics	Premier choix Bacterial cellulitis, erysipelas or lymphangitis
Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid	Second choix Acute pharyngitis
Oral > Solid: 250 mg (as monohydrate)	Chronic obstructive pulmonary disease with acute exacerbation
Oral > Solid: 250 mg (as monohydrate); 500 mg (as monohydrate)	
Oral > Solid > dispersible tablet: 125 mg; 250 mg	Autres indications Bacterial infection of unspecified site

Cefazolin

Section	Indications
<p>Access group antibiotics Parenteral > General injections > unspecified: 1 g in vial (as sodium salt) powder for injection</p>	<p>Premier choix</p> <ul style="list-style-type: none">Other specified prophylactic measures<ul style="list-style-type: none">co-prescrite avec metronidazole <p>Second choix</p> <ul style="list-style-type: none">Osteomyelitis or osteitisBacterial infection of joint <p>Autres indications</p> <ul style="list-style-type: none">Surgical site infection

Cefepime

Section	Indications
<p>Watch group antibiotics Parenteral > General injections > unspecified: 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection; 2 g in vial (as hydrochloride) powder for injection</p>	<p>Other specified bacterial diseases</p>

Cefiderocol

Section	Indications
<p>Reserve group antibiotics Parenteral > General injections > IV: 1 g in vial (as sulfate tosylate) powder for injection</p>	<p>Carbapenem-resistant <i>Pseudomonas aeruginosa</i></p> <p>Carbapenem resistant Enterobacteriales</p>

Cefixime

Section	Indications
<p>Watch group antibiotics Oral > Liquid: 100 mg per 5 mL powder for oral liquid Oral > Solid: 200 mg (as trihydrate); 400 mg (as trihydrate)</p>	<p>Second choix</p> <ul style="list-style-type: none">Infectious gastroenteritis or colitis without specification of infectious agent<ul style="list-style-type: none">co-prescrite avec azithromycin <p>Gonococcal infection</p>

Cefotaxime

Section	Indications
Watch group antibiotics	Premier choix
Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)	Other specified pneumonia (Hospital-acquired pneumonia)
Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)	Bacterial meningitis
	Bacterial pneumonia (Community-acquired pneumonia - severe) [children]
	co-prescrite avec amikacin
	Inflammatory and other diseases of prostate (severe)
	co-prescrite avec amikacin
	Acute pyelonephritis (severe)
	co-prescrite avec clarithromycin
	Bacterial pneumonia (Community-acquired pneumonia - severe)
	co-prescrite avec metronidazole
	Peritonitis (severe)
	co-prescrite avec metronidazole
	Peritonitis (mild-moderate)
	co-prescrite avec metronidazole
	Peritoneal abscess (severe)
	co-prescrite avec metronidazole
	Peritoneal abscess (mild-moderate)
	Second choix
	Osteomyelitis or osteitis
	Bacterial infection of joint
	Inflammatory and other diseases of prostate (mild to moderate)
	Acute pyelonephritis (mild to moderate)
	Sepsis without septic shock
	Autres indications
	Bacterial infection of unspecified site

Ceftaroline

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases
Parenteral > General injections > IV: 400 mg in vial (as fosamil) powder for injection; 600 mg in vial (as fosamil) powder for injection	

Ceftazidime

Section	Indications
<p>Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial (as pentahydrate) powder for injection; 1 g in vial (as pentahydrate) powder for injection</p>	<p>Premier choix co-prescrite avec vancomycin</p> <p>Endophthalmitis</p> <p>Autres indications</p> <p>Pseudomonas aeruginosa resistant to other antibiotic</p>

Ceftazidime + avibactam

Section	Indications
<p>Reserve group antibiotics Parenteral > General injections > IV: 2 g in vial + 0.5 g in vial powder for injection</p>	<p>Carbapenem-resistant Pseudomonas aeruginosa</p> <p>Carbapenem resistant Enterobacteriales</p>

Ceftolozane + tazobactam

Section	Indications
<p>Reserve group antibiotics Parenteral > General injections > IV: 1 g + 0.5 g powder for injection</p>	<p>Carbapenem-resistant Pseudomonas aeruginosa</p>

Ceftriaxone

Section	Indications
<p>Watch group antibiotics Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt) Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt) Parenteral > Locoregional injections > Intravitreal: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt) Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt) Parenteral > General injections > unspecified: 2 g in vial; 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)</p>	<p>Premier choix</p> <p>Typhoid fever</p> <p>Bacterial pneumonia (Community-acquired pneumonia - severe) [children]</p> <p>Paratyphoid fever</p> <p>Other specified pneumonia (Hospital-acquired pneumonia)</p> <p>Bacterial meningitis</p> <p>co-prescrite avec amikacin</p> <p>Acute pyelonephritis (severe)</p> <p>co-prescrite avec amikacin</p> <p>Inflammatory and other diseases of prostate (severe)</p> <p>co-prescrite avec azithromycin</p> <p>Gonococcal infection</p> <p>co-prescrite avec clarithromycin</p> <p>Bacterial pneumonia (Community-acquired pneumonia - severe)</p>

co-prescrite avec metronidazole

Necrotising fasciitis

co-prescrite avec metronidazole

Peritonitis (mild-moderate)

co-prescrite avec metronidazole

Peritonitis (severe)

co-prescrite avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrite avec metronidazole

Peritoneal abscess (severe)

co-prescrite avec vancomycin

Endophthalmitis

Second choix

Bacterial infection of joint

Osteomyelitis or osteitis

Infectious gastroenteritis or colitis without specification of infectious agent

Sepsis without septic shock

Inflammatory and other diseases of prostate (mild to moderate)

Acute pyelonephritis (mild to moderate)

Autres indications

Bacterial infection of unspecified site

Cefuroxime

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg in vial (as sodium salt) powder for injection; 750 mg in vial (as sodium salt) powder for injection; 1.5 g in vial (as sodium salt) powder for injection

Indications

Second choix

Other specified prophylactic measures

Cemiplimab

Section

Immunomodulators

Indications

Therapeutic equivalent to pembrolizumab pour

Other specified malignant neoplasms of bronchus or lung

Certolizumab pegol

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis
	Therapeutic equivalent to adalimumab pour Crohn disease site
	Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Cetirizine

Section	Indications
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to loratadine pour Allergic or hypersensitivity conditions of unspecified type

Chlorambucil

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Oral > Solid: 2 mg tablet	Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Chloramphenicol

Section	Indications
Access group antibiotics	
Parenteral > General injections > IV: 1 g in vial powder for injection (sodium succinate)	Second choix Bacterial meningitis
Parenteral > General injections > IM: 500 mg in 2 mL ampoule oily suspension for injection	Autres indications
Oral > Liquid: 150 mg per 5 mL (as palmitate)	Bacterial infection of unspecified site
Oral > Solid: 250 mg capsule	

Chlorhexidine

Section	Indications
Medicines administered to the neonate [c]	Omphalitis of newborn
Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)	
Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)	
Antiseptics	Chlorhexidine
Local > Topical > Solution: 5% (digluconate) for dilution	

Chlorine base compound

Section	Indications
Disinfectants	Disinfectant, not elsewhere classified
Local > Topical > Powder: 0.1% available chlorine for solution	
Local > Topical > Liquid: 0.1% available chlorine for solution	
Local > Topical > Solid: 0.1% available chlorine for solution	

Chloroquine

Section	Indications
Antimalarial medicines > For chemoprevention	Malaria due to Plasmodium falciparum
Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)	Malaria due to Plasmodium vivax
Oral > Solid: 150 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium ovale
	Malaria due to Plasmodium malariae
Antimalarial medicines > For curative treatment	Malaria due to Plasmodium falciparum
Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)	Malaria due to Plasmodium vivax
Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)	
Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)	
Disease-modifying anti-rheumatic drugs (DMARDs)	Rheumatoid arthritis
Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)	

Chlorothiazide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chloroxylenol

Section	Indications
Disinfectants Local > Topical > Solution: 4.8%	Phenol disinfectant, not elsewhere classified

Chlorpromazine

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to haloperidol pour
Oral > Liquid: 25 mg per 5 mL (hydrochloride)	Schizophrenia or other primary psychotic disorders
Oral > Solid: 100 mg (hydrochloride)	Schizophrenia or other primary psychotic disorders
Oral > Solid: 100 mg (hydrochloride); 10 mg (hydrochloride); 25 mg (hydrochloride); 50 mg (hydrochloride)	Schizophrenia or other primary psychotic disorders
Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (hydrochloride)	

Chlortalidone

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chlortetracycline

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to tetracycline pour Other specified conjunctivitis
	Therapeutic equivalent to tetracycline pour Infectious keratitis
	Therapeutic equivalent to tetracycline pour Infectious blepharitis
	Therapeutic equivalent to tetracycline pour Trachoma

Cholera vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against cholera alone

Ciclesonide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Ciclosporin

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection Oral > Liquid: 100 mg per mL Oral > Solid > capsule: 25 mg	Failure or rejection of transplanted organs or tissues

Cimetidine

Section	Indications
Antiulcer medicines Oral > Solid: 200 mg Parenteral > General injections > unspecified: 200 mg in 2 mL ampoule	Peptic ulcer, site unspecified
	Gastro-oesophageal reflux disease

Ciprofloxacin

Section

Watch group antibiotics

Parenteral > General injections > IV: 2 mg per mL (as hydrate)

Oral > Liquid: 250 mg per 5 mL (anhydrous)

Oral > Solid: 250 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride)

Ear, nose and throat medicines [c]

Local > Otological > drops: 0.3% (as hydrochloride)

Indications

Premier choix

Neutropenia (low-risk)

Inflammatory and other diseases of prostate (mild to moderate)

Typhoid fever

Paratyphoid fever

Infectious gastroenteritis or colitis without specification of infectious agent

Acute pyelonephritis (mild to moderate)

Second choix

Cholera

Cholera [children]

co-prescrite avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrite avec metronidazole

Peritonitis (mild-moderate)

Autres indications

Bacterial infection of unspecified site

Infectious diseases of external ear

Cisplatin

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection

Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Germ cell tumour of testis

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Cladribine

Section	Indications
Medicines for multiple sclerosis Oral > Solid > tablet: 10 mg	Multiple sclerosis
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial	Langerhans cell histiocytosis

Clarithromycin

Section	Indications
Watch group antibiotics Oral > Liquid: 125 mg per 5 mL powder for oral liquid; 250 mg per 5 mL powder for oral liquid Parenteral > General injections > unspecified: 500 mg in vial powder for injection Oral > Solid > dosage form: 500 mg; 250 mg Oral > Solid > dosage form: 500 mg	Premier choix co-prescrite avec cefotaxime Bacterial pneumonia (Community-acquired pneumonia - severe) co-prescrite avec ceftriaxone Bacterial pneumonia (Community-acquired pneumonia - severe) Second choix Acute pharyngitis co-prescrite avec amoxicillin + clavulanic acid Bacterial pneumonia (Community-acquired pneumonia - severe) Autres indications Helicobacter pylori associated gastric ulcer

Clindamycin

Section	Indications
Access group antibiotics Oral > Liquid: 75 mg per 5 mL (as palmitate hydrochloride) Oral > Solid: 150 mg (as hydrochloride) capsule Parenteral > General injections > unspecified: 150 mg per mL (as phosphate) injection Parenteral > General injections > IV: 150 mg per mL (as phosphate); 600 mg per 4 mL (as phosphate); 900 mg per 6 mL (as phosphate) Oral > Liquid: 75 mg per 5 mL powder for oral liquid (as palmitate hydrochloride) Oral > Solid: 150 mg (as hydrochloride)	Premier choix co-prescrite avec piperacillin + tazobactam Necrotising fasciitis Second choix Osteomyelitis or osteitis Bacterial infection of joint Autres indications Bacterial infection of unspecified site

Clofazimine

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg; 50 mg	Multi-drug resistant <i>Mycobacterium tuberculosis</i>
Antileprosy medicines Oral > Solid: 100 mg; 50 mg	Leprosy

Clomifene

Section	Indications
Ovulation inducers Oral > Solid: 50 mg (citrate)	Female infertility without specification whether primary or secondary

Clomipramine

Section	Indications
Medicines for obsessive compulsive disorders Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)	Obsessive-compulsive disorder

Clonazepam

Section	Indications
Antiseizure medicines Oral > Solid: 500 µg	Generalised myoclonic seizure

Clopidogrel

Section	Indications
Anti-platelet medicines Oral > Solid: 75 mg; 300 mg	Acute ischaemic heart disease
	Presence of coronary angioplasty implant or graft

Clotrimazole

Section	Indications
Antifungal medicines Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet	Vulvovaginal candidosis

Cloxacillin

Section	Indications
Access group antibiotics	Premier choix
Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid	Bacterial infection of joint
Oral > Solid: 500 mg (as sodium salt) capsule; 1 g (as sodium salt) capsule	Osteomyelitis or osteitis
Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection	Bacterial cellulitis, erysipelas or lymphangitis
Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid; 250 mg per 5 mL (as sodium salt) powder for oral liquid	Second choix
Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 250 mg in vial (as sodium salt) powder for injection	co-prescrite avec amikacin
Oral > Solid > capsule: 500 mg (as sodium salt); 1 g (as sodium salt); 250 mg (as sodium salt)	Sepsis without septic shock
	Autres indications
	Bacterial infection of unspecified site

Clozapine

Section	Indications
Medicines for psychotic disorders	Schizophrenia or other primary psychotic disorders

Coagulation factor IX

Section	Indications
Coagulation factors	Coagulation defects, purpura or other haemorrhagic or related conditions Haemophilia B

Coagulation factor IX complex

Section	Indications
Coagulation factors	Therapeutic equivalent to coagulation factor IX pour Coagulation defects, purpura or other haemorrhagic or related conditions Therapeutic equivalent to coagulation factor IX pour Haemophilia B

Coagulation factor VIII

Section	Indications
Coagulation factors Parenteral > General injections > IV: 250 IU in vial powder for injection; 500 IU in vial powder for injection; 1000 IU in vial powder for injection	Haemophilia A

Coal tar

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Solution: 5%	Psoriasis of unspecified type

Cobicistat + elvitegravir + emtricitabine + tenofovir disoproxil fumarate

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobicistat + elvitegravir + emtricitabine+ tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 10 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobimetinib

Section	Indications
Targeted therapies	Therapeutic equivalent to trametinib pour Other specified melanoma of skin

Codeine

Section	Indications
Opioid analgesics Oral > Solid: 30 mg tablet (codeine phosphate)	Pain
Antitussives Oral > Solid: 10 mg tablet (codeine phosphate)	Cough
Medicines for diarrhoea Oral > Solid: 30 mg tablet (codeine phosphate)	Diarrhoea

Colchicine

Section	Indications
Medicines used to treat gout Oral > Solid: 0.5 mg	Gout

Colecalciferol

Section	Indications
Vitamins and minerals Oral > Liquid: 400 IU per mL Oral > Solid: 400 IU; 1000 IU	Calcium deficiency Vitamin D deficiency Therapeutic equivalent to ergocalciferol pour Vitamin D deficiency Therapeutic equivalent to ergocalciferol pour Calcium deficiency

Colistin (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 million IU in vial (as colistimethate sodium) Parenteral > General injections > unspecified: 1 million IU in vial (as colistimethate sodium) (equivalent to 34 mg colistin base activity)	Other specified bacterial diseases Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacteriales

Compound sodium lactate solution

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: injectable solution	Other specified disorders of fluid, electrolyte or acid-base balance Hypovolaemic shock

Condoms

Section	Indications
Barrier methods	Contact with health services for contraceptive management

Copper-containing intrauterine device

Section	Indications
Intrauterine devices	Contact with health services for insertion of contraceptive device

Crizotinib

Section	Indications
Targeted therapies	Other specified malignant neoplasms of bronchus or lung
Oral > Solid: 200 mg; 250 mg	Anaplastic large cell lymphoma, ALK-positive
Oral > Solid > capsule: 200 mg; 250 mg	

Cryoprecipitate (not pathogen-reduced)

Section	Indications
Blood and blood components	Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour
	Haemophilia A
	Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour
	Coagulation defects, purpura or other haemorrhagic or related conditions
	Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour
	Von Willebrand disease
	Therapeutic equivalent to cryoprecipitate (pathogen-reduced) pour
	Obstetric haemorrhage

Cryoprecipitate (pathogen-reduced)

Section	Indications
Blood and blood components	Haemophilia A
Injection: frozen liquid in bag or lyophilized powder in vial containing:	Coagulation defects, purpura or other haemorrhagic or related conditions
> 50 IU Factor VIII	Von Willebrand disease
> 100 IU vWF	
> 140 mg clottable fibrinogen per unit	Obstetric haemorrhage

Cyanocobalamin

Section	Indications
Antianaemia medicines	Anaemias or other erythrocyte disorders

Cyclizine

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 50 mg tablet Parenteral > General injections > unspecified: 50 mg per mL	Palliative care

Cyclopentolate

Section	Indications
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour Cycloplegic drug
Ophthalmological preparations > Mydriatics	Therapeutic equivalent to atropine pour Anterior uveitis

Cyclophosphamide

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Oral > Solid: 25 mg tablet	Other specified malignant neoplasms of breast
Parenteral > General injections > unspecified: 500 mg in vial powder for injection	Malignant trophoblastic neoplasms of placenta
Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection	Follicular lymphoma
Oral > Solid > dosage form: 25 mg; 50 mg	Malignant neoplasms of breast
	Chronic lymphocytic leukaemia or small lymphocytic lymphoma
	Rhabdomyosarcoma primary site
	Hodgkin lymphoma
	Diffuse large B-cell lymphomas
	Plasma cell myeloma
	Other specified gliomas of brain
	Burkitt lymphoma including Burkitt leukaemia
	Malignant neoplasms of kidney, except renal pelvis
	Ewing sarcoma of bone and articular cartilage of unspecified sites
	Lymphoid leukaemia, not elsewhere classified
	Anaplastic large cell lymphoma, ALK-negative
	Anaplastic large cell lymphoma, ALK-positive

Cycloserine

Section	Indications
Antituberculosis medicines Oral > Solid: 250 mg; 125 mg	Multi-drug resistant Mycobacterium tuberculosis

Cytarabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 100 mg in vial powder for injection Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Acute myeloid leukaemia with recurrent genetic abnormalities Burkitt lymphoma including Burkitt leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis

Dabigatran

Section	Indications
Medicines affecting coagulation Oral > Solid: 110 mg; 150 mg	Venous thromboembolism Other specified atrial fibrillation

Dabrafenib

Section	Indications
Targeted therapies Oral > Solid: 50 mg; 75 mg	Other specified melanoma of skin

Dacarbazine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection	Hodgkin lymphoma

Daclatasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 30 mg (as hydrochloride); 60 mg (as hydrochloride)	Chronic hepatitis C

Daclatasvir + sofosbuvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 60 mg + 400 mg	Chronic hepatitis C

Dactinomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 µg in vial powder for injection	Malignant trophoblastic neoplasms of placenta Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site Ewing sarcoma of bone and articular cartilage of unspecified sites

Dalteparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin pour Acute ischaemic heart disease Therapeutic equivalent to enoxaparin pour Venous thromboembolism

Dapagliflozin

Section	Indications
Oral hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Dapsone

Section	Indications
Antileprosy medicines Oral > Solid: 100 mg tablet; 50 mg tablet; 25 mg tablet	Leprosy

Daptomycin

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 350 mg in vial powder for injection; 500 mg in vial powder for injection	Other specified bacterial diseases

Daratumumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL	Plasma cell myeloma

Darbepoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Darunavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dasabuvir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 250 mg tablet	Chronic hepatitis C

Dasatinib

Section	Indications
<p>Targeted therapies</p> <p>Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet</p>	Chronic myeloid leukaemia, not elsewhere classified

Daunorubicin

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)</p>	Acute myeloid leukaemia with recurrent genetic abnormalities Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia

Deferasirox

Section	Indications
<p>Other medicines for haemoglobinopathies</p> <p>Oral > Solid > dispersible tablet: 100 mg; 125 mg; 250 mg; 400 mg; 500 mg</p> <p>Oral > Solid > tablet: 90 mg; 180 mg; 360 mg</p>	Other specified sickle cell disorders or other haemoglobinopathies Thalassaemias Therapeutic equivalent to deferoxamine pour Thalassaemias

Deferiprone

Section	Indications
<p>Other medicines for haemoglobinopathies</p>	Therapeutic equivalent to deferasirox pour Other specified sickle cell disorders or other haemoglobinopathies Therapeutic equivalent to deferasirox pour Thalassaemias

Deferoxamine

Section	Indications
<p>Antidotes and other substances used in poisonings > Specific</p> <p>Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)</p>	Harmful effects of drugs, medicaments or biological substances, not elsewhere classified
<p>Other medicines for haemoglobinopathies</p> <p>Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)</p>	Other specified sickle cell disorders or other haemoglobinopathies Thalassaemias

Delafloxacin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 300 mg lyophilized powder for injection Oral > Solid: 450 mg	Methicillin resistant <i>Staphylococcus aureus</i>

Delamanid

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 25 mg Oral > Solid > tablet: 50 mg	Multi-drug resistant <i>Mycobacterium tuberculosis</i>

Dengue vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Desmopressin

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 4 µg per mL in 1 mL ampoule (acetate) Local > Nasal > Spray: 10 µg per dose (acetate)	Haemophilia A Von Willebrand disease

Dexamethasone

Section	Indications
<p>Other medicines administered to the mother</p> <p>Parenteral > General injections > unspecified: 4 mg per mL (dexamethasone phosphate (as sodium))</p>	<p>Respiratory distress syndrome of the newborn, altered by maternal corticosteroid therapy</p>
<p>Hormones and antihormones</p> <p>Oral > Solid: 4 mg</p> <p>Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)</p> <p>Oral > Liquid: 2 mg per 5 mL</p> <p>Oral > Solid: 2 mg; 4 mg</p>	<p>Plasma cell myeloma</p> <p>Lymphoid leukaemia, not elsewhere classified</p> <p>Burkitt lymphoma including Burkitt leukaemia</p> <p>Anaplastic large cell lymphoma, ALK-negative</p> <p>Anaplastic large cell lymphoma, ALK-positive</p>
<p>Antiallergics and medicines used in anaphylaxis</p> <p>Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)</p>	<p>Allergic or hypersensitivity conditions of unspecified type</p>
<p>Medicines for other common symptoms in palliative care</p> <p>Oral > Liquid: 2 mg per 5 mL</p> <p>Oral > Solid: 2 mg; 4 mg</p> <p>Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)</p>	<p>Palliative care</p>
<p>Antiemetic medicines</p> <p>Oral > Liquid: 0.5 mg per 5 mL; 2 mg per 5 mL</p> <p>Oral > Solid: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg</p> <p>Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)</p>	<p>Nausea or vomiting</p>

Dextran 40

Section	Indications
Plasma substitutes	Hypovolaemia

Dextran 70

Section	Indications
Plasma substitutes	Hypovolaemia

Parenteral > General injections > IV: 6% injectable solution

Diaphragms

Section	Indications
Barrier methods -	Contact with health services for contraceptive management

Diazepam

Section	Indications
Antiseizure medicines	Therapeutic equivalent to lorazepam pour
Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system	Status epilepticus
Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube	Status epilepticus
Medicines for other common symptoms in palliative care	Palliative care
Oral > Liquid: 2 mg per 5 mL	
Oral > Solid: 5 mg; 10 mg	
Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system	
Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube	
Parenteral > General injections > unspecified: 5 mg per mL	
Medicines for anxiety disorders	Anxiety
Oral > Solid: 5 mg (scored); 2 mg (scored)	

Diazoxide

Section	Indications
Medicines for hypoglycaemia	Persistent hyperinsulinaemic hypoglycaemia of infancy
Oral > Liquid: 50 mg per mL	
Oral > Solid: 50 mg	

Didanosine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 100 mg buffered powder for oral solution; 167 mg buffered powder for oral solution; 250 mg buffered powder for oral solution Oral > Solid: 25 mg buffered chewable dispersible tablet; 50 mg buffered chewable dispersible tablet; 100 mg buffered chewable dispersible tablet; 150 mg buffered chewable dispersible tablet; 200 mg buffered chewable dispersible tablet; 125 mg unbuffered enteric coated capsule; 200 mg unbuffered enteric coated capsule; 250 mg unbuffered enteric coated capsule; 400 mg unbuffered enteric coated capsule	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Diethylcarbamazine

Section	Indications
Antifilarials Oral > Solid: 50 mg tablet (dihydrogen citrate); 100 mg tablet (dihydrogen citrate)	Lymphatic filariasis

Digitoxin

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule Oral > Solid: 50 µg; 100 µg	Heart failure

Digoxin

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid: 62.5 µg; 250 µg	Cardiac arrhythmia
Medicines used in heart failure Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg	Heart failure

Dihydroartemisinin + piperaquine phosphate

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Dihydroergocryptine mesylate

Section	Indications
Antiseizure medicines Oral > Solid:	Parkinson disease

Diloxanide

Section	Indications
Antiamoebic and antigiardiasis medicines Oral > Solid: 500 mg (furoate)	Amoebiasis

Dimercaprol

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Diphtheria antitoxin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IM: 10000 IU in vial; 20000 IU in vial	Need for immunization against diphtheria alone

Diphtheria vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against diphtheria alone

Diphtheria-pertussis-tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against diphtheria-tetanus-pertussis, combined

Diphtheria-tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against combinations of infectious diseases

Docetaxel

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL	Other specified malignant neoplasms of breast Malignant neoplasms of breast Malignant neoplasms of prostate Unspecified malignant neoplasms of ill-defined or unspecified sites

Docusate sodium

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 50 mg per 5 mL oral liquid Oral > Solid: 100 mg capsule	Palliative care

Dolasetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Dolutegravir

Section	Indications
Antiretrovirals > Integrase inhibitors Oral > Solid: 50 mg tablet; 10 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dolutegravir + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Donepezil

Section	Indications
Medicines for diseases of the nervous system Oral > Liquid: 1 mg per mL (hydrochloride) Oral > Solid > tablet: 5 mg (hydrochloride); 10 mg (hydrochloride); 5 mg (hydrochloride) (orodispersible); 10 mg (hydrochloride) (orodispersible)	Dementia due to Alzheimer disease

Dopamine

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)	Heart failure Cardiogenic shock

Doxorubicin

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)</p> <p>Parenteral > General injections > IV: 10 mg in vial (hydrochloride) powder for injection; 50 mg in vial (hydrochloride) powder for injection; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)</p>	<p>Unspecified malignant neoplasms of ill-defined or unspecified sites</p> <p>Rhabdomyosarcoma primary site</p> <p>Malignant neoplasms of breast</p> <p>Hodgkin lymphoma</p> <p>Diffuse large B-cell lymphomas</p> <p>Plasma cell myeloma</p> <p>Follicular lymphoma</p> <p>Burkitt lymphoma including Burkitt leukaemia</p> <p>Malignant neoplasms of kidney, except renal pelvis</p> <p>Osteosarcoma of bone and articular cartilage of unspecified sites</p> <p>Lymphoid leukaemia, not elsewhere classified</p> <p>Anaplastic large cell lymphoma, ALK-negative</p> <p>Other specified malignant neoplasms of breast</p> <p>Anaplastic large cell lymphoma, ALK-positive</p> <p>Ewing sarcoma of bone and articular cartilage of unspecified sites</p> <p>Kaposi sarcoma of unspecified primary site</p>

Doxorubicin (as pegylated liposomal)

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)</p>	<p>Kaposi sarcoma of unspecified primary site</p>

Doxycycline

Section	Indications
Access group antibiotics	Premier choix
Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous)	Chlamydia trachomatis
Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate)	Cholera
Parenteral > General injections > unspecified: 100 mg in vial powder for injection	Second choix
Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium)	Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)
Oral > Solid > dispersible tablet: 100 mg (as monohydrate)	Chronic obstructive pulmonary disease with acute exacerbation
Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)	Cholera [children]
Oral > Solid > dosage form: 100 mg (as hyclate)	Autres indications
Antimalarial medicines > For curative treatment	Bacterial infection of unspecified site
Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)	Malaria due to Plasmodium falciparum
Antimalarial medicines > For chemoprevention	Malaria due to Plasmodium falciparum
Oral > Solid: 100 mg (hydrochloride or hyclate)	

Durvalumab

Section	Indications
Immunomodulators	Other specified malignant neoplasms of bronchus or lung
Parenteral > General injections > IV: 120 mg per 2.4 mL in vial; 500 mg per 10 mL in vial	

Edoxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism
	Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Efavirenz

Section	Indications
Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified
Oral > Solid: 600 mg tablet	

Efavirenz + emtricitabine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Efavirenz + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Therapeutic equivalent to efavirenz + emtricitabine + tenofovir pour Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Eflornithine

Section	Indications
Medicines for the treatment of 2nd stage African trypanosomiasis Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)	African trypanosomiasis

Elbasvir + grazoprevir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 50 mg + 100 mg	Chronic hepatitis C

Empagliflozin

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 10 mg; 25 mg	Type 2 diabetes mellitus

Emtricitabine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 10 mg per mL Oral > Solid: 200 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 25 mg + 25 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir disoproxil fumarate

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 25 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified Contact with or exposure to human immunodeficiency virus

Emtricitabine + tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 200 mg + 10 mg; 200 mg + 25 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Enalapril

Section	Indications
Antihypertensive medicines Oral > Liquid: 1 mg per mL (as hydrogen maleate) Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate) Medicines used in heart failure Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Essential hypertension Heart failure

Encorafenib

Section	Indications
Targeted therapies	Therapeutic equivalent to dabrafenib pour Other specified melanoma of skin

Enoxaparin

Section	Indications
Medicines affecting coagulation Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule; 120 mg per 0.8 mL in ampoule	Acute ischaemic heart disease Venous thromboembolism

Entecavir

Section	Indications
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 0.05 mg per mL Oral > Solid: 0.5 mg; 1 mg	Chronic hepatitis B

Enzalutamide

Section	Indications
Hormones and antihormones Oral > Solid: 40 mg	Therapeutic equivalent to abiraterone pour Malignant neoplasms of prostate Malignant neoplasms of prostate

Ephedrine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 30 mg per mL in 1 mL ampoule (ephedrine hydrochloride)	Cardiac complications of anaesthesia during labour or delivery

Epinephrine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)	Cardiac arrest
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 2% (as hydrochloride)	Primary open-angle glaucoma
Antiasthmatic and medicines for chronic obstructive pulmonary disease Parenteral > General injections > unspecified: 1 mg per mL in ampoule (as hydrochloride or hydrogen tartrate)	Asthma Chronic obstructive pulmonary disease
Antiallergics and medicines used in anaphylaxis Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (as hydrochloride or hydrogen tartrate)	Anaphylaxis

Epoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Epoetin beta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Epoetin theta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Equine rabies immunoglobulin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > Locoregional injections > Intradermal: 150 IU per mL in vial; 200 IU per mL in vial; 300 IU per mL in vial; 400 IU per mL in vial	Rabies

Eravacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg lyophilized powder for injection	Carbapenem resistant Enterobacteriales

Ergocalciferol

Section	Indications
Vitamins and minerals Oral > Liquid: 250 µg per mL (10 000 IU per mL)	Therapeutic equivalent to colecalciferol pour Calcium deficiency
Oral > Solid: 1.25 mg (50 000 IU)	Therapeutic equivalent to colecalciferol pour Vitamin D deficiency
	Vitamin D deficiency
	Calcium deficiency

Ergometrine

Section	Indications
Uterotonics Parenteral > General injections > unspecified: 200 µg in 1 mL ampoule (hydrogen maleate)	Postpartum haemorrhage

Erlotinib

Section	Indications
Targeted therapies Oral > Solid: 100 mg; 150 mg	Other specified malignant neoplasms of bronchus or lung

Erythromycin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 500 mg in vial powder for injection (as lactobionate)	Second choix Therapeutic equivalent to clarithromycin pour Acute pharyngitis
Oral > Liquid: 125 mg per 5 mL (as stearate or ethylsuccinate)	Autres indications
Oral > Solid: 250 mg (as stearate or ethylsuccinate)	Bacterial infection of unspecified site
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 0.5%	Chlamydial conjunctivitis Gonococcal infection of eye

Erythropoiesis-stimulating agents

Section	Indications
<p>Antianaemia medicines</p> <p>Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe</p>	<p>Anaemia due to chronic disease</p>

Estradiol

Section	Indications
<p>Estrogens</p> <p>Oral > Solid > tablet: 0.5 mg; 1 mg; 2 mg</p>	<p>Gonadotropin deficiency</p> <p>Hypopituitarism</p> <p>Turner syndrome</p>

Estradiol cypionate + medroxyprogesterone acetate

Section	Indications
<p>Injectable hormonal contraceptives</p> <p>Parenteral > General injections > IM: 5 mg + 25 mg</p>	<p>Contact with health services for reasons associated with reproduction</p>

Etanercept

Section	Indications
<p>Immunomodulators for non-malignant disease</p>	<p>Therapeutic equivalent to adalimumab pour</p> <p>Axial spondyloarthritis</p> <p>Therapeutic equivalent to adalimumab pour</p> <p>Crohn disease site</p> <p>Therapeutic equivalent to adalimumab pour</p> <p>Rheumatoid arthritis</p>
<p>Medicines for juvenile joint diseases</p>	<p>Therapeutic equivalent to adalimumab pour</p> <p>Juvenile idiopathic arthritis</p>

Ethambutol

Section	Indications
Antituberculosis medicines	Other specified tuberculosis
Parenteral > General injections > IV: 1000 mg per 10 mL; 2000 mg per 20 mL	Tuberculosis
Oral > Solid > dispersible tablet: 100 mg	
Oral > Solid > tablet: 100 mg (hydrochloride); 400 mg (hydrochloride)	

Ethambutol + isoniazid

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 400 mg + 150 mg tablet	

Ethambutol + isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet	

Ethambutol + isoniazid + rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid: 275 mg + 75 mg + 150 mg tablet	

Ethanol

Section	Indications
Antiseptics	Denatured alcohol
Local > Topical > Solution: 70% (denatured)	

Ether

Section	Indications
General anaesthetics and oxygen > Inhalational medicines	Anaesthetics and therapeutic gases
Respiratory > Inhalation > solution:	

Ethinylestradiol + etonogestrel

Section	Indications
Intravaginal contraceptives	Contact with health services for contraceptive management
Local > Vaginal > vaginal ring: 2.7 mg + 11.7 mg	

Ethinylestradiol + levonorgestrel

Section	Indications
Oral hormonal contraceptives Oral > Solid: 50 µg + 250 µg [4]	Contact with health services for postcoital contraception
Oral > Solid: 30 µg + 150 µg	Contact with health services for contraceptive management

Ethinylestradiol + norethisterone

Section	Indications
Oral hormonal contraceptives Oral > Solid: 35 µg + 1 mg	Contact with health services for contraceptive management

Ethionamide

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 125 mg	Multi-drug resistant Mycobacterium tuberculosis
Oral > Solid > tablet: 250 mg	Tuberculous meningitis

Ethosuximide

Section	Indications
Antiseizure medicines Oral > Liquid: 250 mg per 5 mL	Absence seizures, typical
Oral > Solid: 250 mg	

Etonogestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 68 mg single rod	Contact with health services for insertion of contraceptive device

Etoposide

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule</p> <p>Oral > Solid: 100 mg capsule</p> <p>Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate)</p> <p>Oral > Solid: 100 mg capsule; 50 mg capsule</p>	<p>Unspecified malignant neoplasms of ill-defined or unspecified sites</p> <p>Burkitt lymphoma including Burkitt leukaemia</p> <p>Ewing sarcoma of bone and articular cartilage of unspecified sites</p> <p>Lymphoid leukaemia, not elsewhere classified</p> <p>Myeloid leukaemia</p> <p>Other specified malignant neoplasms of the ovary</p> <p>Other specified malignant neoplasms of bronchus or lung</p> <p>Germ cell tumour of testis</p> <p>Malignant trophoblastic neoplasms of placenta</p> <p>Malignant neoplasms of kidney, except renal pelvis</p> <p>Osteosarcoma of bone and articular cartilage of unspecified sites</p> <p>Hodgkin lymphoma</p> <p>Retinoblastoma</p> <p>Anaplastic large cell lymphoma, ALK-positive</p> <p>Anaplastic large cell lymphoma, ALK-negative</p>

Everolimus

Section	Indications
<p>Targeted therapies</p> <p>Oral > Solid > dispersible tablet: 2 mg; 3 mg; 5 mg</p> <p>Oral > Solid > tablet: 2.5 mg; 5 mg; 7.5 mg; 10 mg</p>	Other specified gliomas of brain

Fentanyl

Section	Indications
<p>Opioid analgesics</p> <p>Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour</p> <p>Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate)</p> <p>Oral > Other: 200 to 1200 µg buccal film (as citrate)</p>	<p>Chronic cancer pain</p> <p>Other specified chronic cancer related pain</p>

Ferrous salt

Section	Indications
Antianaemia medicines	Iron deficiency anaemia
Oral > Liquid: 25 mg per mL iron (as sulfate)	
Oral > Solid: 60 mg iron	

Ferrous salt + folic acid

Section	Indications
Antianaemia medicines	Iron deficiency anaemia
Oral > Solid: 60 mg iron + 2.8 mg	Clefts of lip, alveolus or palate
Oral > Solid > tablet: 60 mg elemental iron + 400 µg tablet; 60 mg elemental iron + 2.8 mg	Iron deficiency anaemia
	Anencephaly or similar anomalies
	Spina bifida

Fexinidazole

Section	Indications
Medicines for the treatment of 1st stage African trypanosomiasis Oral > Solid: 600 mg	African trypanosomiasis
Medicines for the treatment of 2nd stage African trypanosomiasis Oral > Solid: 600 mg	African trypanosomiasis

Fexofenadine

Section	Indications
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to loratadine pour Allergic or hypersensitivity conditions of unspecified type

Fifth generation cephalosporins

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases

Filgrastim

Section	Indications
Immunomodulators Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial	Acquired neutropaenia

Fingolimod

Section	Indications
Immunomodulators for non-malignant disease Oral > Solid: 0.25 mg; 0.5 mg	Relapsing-remitting multiple sclerosis

Flomoxef

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 0.5 g in vial powder for injection; 1 g in vial powder for injection	Other specified other antibiotic resistant Enterobacteriales

Fluconazole

Section	Indications
Antifungal medicines	Cryptococcosis
Parenteral > General injections > IV: 2 mg per mL in vial	Candidosis
Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid	
Oral > Solid > capsule: 50 mg	

Flucytosine

Section	Indications
Antifungal medicines	Cryptococcosis
Parenteral > General injections > IV: 2.5 g per 250 mL infusion	
Oral > Solid: 250 mg capsule	

Fludarabine

Section	Indications
Cytotoxic medicines	Chronic lymphocytic leukaemia or small lymphocytic lymphoma
Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection	
Oral > Solid: 10 mg tablet (fludarabine phosphate)	

Fludrocortisone

Section	Indications
Adrenal hormones and synthetic substitutes Oral > Solid: 100 µg (acetate)	Adrenocortical insufficiency

Flunisolide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Fluorescein

Section	Indications
Diagnostic agents > Ophthalmic medicines Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)	Fluorescein

Fluoride

Section	Indications
Dental medicines and preparations Local > Dental > Gel: 2500 to 12500 ppm fluoride (any type) Local > Dental > Toothpaste, cream or gel: 1000 to 1500 ppm fluoride (any type) Local > Dental > Mouthrinse: 230 to 900 ppm fluoride (any type) Local > Dental > Varnish: 22500 ppm fluoride (any type)	Dental caries

Fluorouracil

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Ointment: 5% Cytotoxic medicines Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule Parenteral > General injections > IV: 50 mg per mL in vial	Plantar warts Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of nasopharynx Malignant neoplasm metastasis in large intestine Malignant neoplasms of breast Malignant neoplasms of rectum Malignant neoplasms of colon

Fluoxetine

Section	Indications
Medicines for depressive disorders Oral > Solid: 20 mg (as hydrochloride)	Depressive disorders
Medicines for other common symptoms in palliative care Oral > Solid: 20 mg (as hydrochloride)	Palliative care
Medicines for anxiety disorders Oral > Solid: 20 mg (as hydrochloride)	Social anxiety disorder Generalised anxiety disorder Panic disorder
Medicines for obsessive compulsive disorders Oral > Solid: 20 mg (as hydrochloride)	Obsessive-compulsive disorder

Fluphenazine

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 25 mg per mL in ampoule (decanoate); 25 mg per mL in ampoule (enanthate)	Schizophrenia or other primary psychotic disorders

Flutamide

Section	Indications
Hormones and antihormones	Therapeutic equivalent to bicalutamide pour Malignant neoplasms of prostate

Fluticasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Fluticasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Fluticasone furoate + vilanterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Fluvastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia
	Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Folic acid

Section	Indications
Antianaemia medicines	Folate deficiency anaemia
Oral > Solid: 1 mg tablet; 5 mg tablet	Anencephaly or similar anomalies
Oral > Solid: 400 µg	Spina bifida
	Cephalocele

Fomepizole

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base)	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fosfomycin (injection)

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases
Parenteral > General injections > IV: 2 g in vial (as sodium) powder for injection; 4 g in vial (as sodium) powder for injection	Carbapenem resistant Acinetobacter baumannii
	Carbapenem-resistant Pseudomonas aeruginosa
	Carbapenem resistant Enterobacteriales

Fourth generation cephalosporins

Section	Indications
Watch group antibiotics	Other specified bacterial diseases

Fresh-frozen plasma

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Fulvestrant

Section	Indications
Hormones and antihormones Parenteral > General injections > IV: 250 mg per 5 mL	Malignant neoplasms of breast

Furosemide

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL Oral > Solid > tablet: 40 mg; 20 mg	Heart failure
Diuretics Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL Oral > Solid > tablet: 40 mg; 20 mg	Anuria or oliguria Oedema

Gabapentin

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg	Neuropathic pain

Gallamine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 40 mg per mL in 2 mL ampoule (gallamine triethiodide)	Muscle relaxants

Gatifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis

Gefitinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib pour Other specified malignant neoplasms of bronchus or lung

Gemcitabine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection	Other specified carcinomas of ovary Other specified malignant neoplasms of bronchus or lung

Gentamicin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)	Premier choix Neonatal meningitis co-prescrite avec amoxicillin Sepsis without septic shock co-prescrite avec amoxicillin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec ampicillin Bacterial pneumonia (Community-acquired pneumonia - severe) [children] co-prescrite avec ampicillin Sepsis without septic shock co-prescrite avec ampicillin Acute malnutrition in infants, children or adolescents (complicated) [children] co-prescrite avec ampicillin Peritoneal abscess (severe) co-prescrite avec ampicillin Peritonitis (mild-moderate) co-prescrite avec ampicillin Peritonitis (severe)

	<p>co-prescrite avec ampicillin</p> <p>Peritoneal abscess (mild-moderate)</p>
	<p>co-prescrite avec benzylpenicillin</p> <p>Bacterial pneumonia (Community-acquired pneumonia - severe) [children]</p>
	<p>co-prescrite avec benzylpenicillin</p> <p>Acute malnutrition in infants, children or adolescents (complicated) [children]</p>
	<p>co-prescrite avec benzylpenicillin</p> <p>Sepsis without septic shock</p>
	<p>Second choix</p> <p>Other specified prophylactic measures</p> <p>Gonococcal infection</p>
	<p>Autres indications</p> <p>Bacterial infection of unspecified site</p>
	<p>Other specified conjunctivitis</p> <p>Infectious blepharitis</p>

Glass ionomer cement

Section	Indications
Dental medicines and preparations Local > Dental > Powder + liquid: Single-use capsules: 0.4 g powder + 0.09 mL liquid; Multi-use bottle: powder + liquid	Dental caries

Glatiramer acetate

Section	Indications
Medicines for multiple sclerosis Parenteral > General injections > SC: 20 mg per mL; 40 mg per mL Parenteral > General injections > SC: 20 mg per mL in pre-filled syringe; 40 mg per mL in pre-filled syringe	Relapsing-remitting multiple sclerosis Multiple sclerosis

Glecaprevir + pibrentasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid: 50 mg + 20 mg granules Oral > Solid > tablet: 100 mg + 40 mg	Chronic hepatitis C

Glibenclamide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 5 mg; 2.5 mg	Type 2 diabetes mellitus

Gliclazide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release	Type 2 diabetes mellitus

Glucagon

Section	Indications
Medicines for hypoglycaemia Parenteral > General injections > unspecified: 1 mg per mL	Hypoglycaemia in the context of diabetes

Glucose

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral	Other specified disorders of fluid, electrolyte or acid-base balance
Parenteral > General injections > IV: 5% isotonic; 10% hypertonic	Hypoglycaemia without associated diabetes
Parenteral > General injections > IV: 50% hypertonic	

Glucose + sodium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 4% + 0.18% (equivalent to Na+ 30 mmol/L and Cl- 30 mmol/L; 5% + 0.9% (equivalent to Na+ 150 mmol/L and Cl- 150 mmol/L); 5% + 0.45% (equivalent to Na+ 75 mmol/L and Cl- 75 mmol/L)	Other specified disorders of fluid, electrolyte or acid-base balance

Glutaral

Section	Indications
Disinfectants Local > Topical > Solution: 2%	Glutaral nonmedicinal

Glyceryl trinitrate

Section	Indications
Antianginal medicines Oral > Solid: 500 µg sublingual tablet	Angina pectoris

Glycopyrronium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Golimumab

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis
	Therapeutic equivalent to adalimumab pour Crohn disease site
	Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Goserelin

Section	Indications
Hormones and antihormones	Therapeutic equivalent to leuprorelin pour Malignant neoplasms of breast
	Therapeutic equivalent to leuprorelin pour Malignant neoplasms of prostate

Granisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Griseofulvin

Section	Indications
Antifungal medicines Oral > Liquid: 125 mg per 5 mL Oral > Solid: 125 mg; 250 mg	Dermatophytosis

Haemophilus influenzae type b vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Haloperidol

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per mL Oral > Solid: 0.5 mg; 2 mg; 5 mg Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule	Palliative care
Medicines for psychotic disorders Parenteral > General injections > IM: 5 mg per mL in ampoule Oral > Solid: 2 mg; 5 mg	Schizophrenia or other primary psychotic disorders

Haloperidol decanoate

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to fluphenazine pour Schizophrenia or other primary psychotic disorders

Halothane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines	Anaesthetics and therapeutic gases
Respiratory > Inhalation > liquid:	

Heparin sodium

Section	Indications
Medicines affecting coagulation	Acute ischaemic heart disease
Parenteral > General injections > IV: 1000 IU per mL in 1 mL ampoule; 20000 IU per mL in 1 mL ampoule; 5000 IU per mL in 1 mL ampoule	Unstable angina
	Venous thromboembolism

Hepatitis A vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against viral hepatitis

Hepatitis B vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against viral hepatitis

Homatropine

Section	Indications
Ophthalmological preparations > Mydriatics Local > Ophthalmological > Solution (eye drops): 2% (hydrobromide)	Anterior uveitis Therapeutic equivalent to atropine pour Anterior uveitis

Human papilloma virus (HPV) vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Hydralazine

Section	Indications
Antihypertensive medicines	Essential hypertension
Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)	Gestational hypertension
Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)	
Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)	

Hydrochlorothiazide

Section	Indications
Antihypertensive medicines	Essential hypertension
Oral > Liquid: 50 mg per 5 mL	
Oral > Solid: 25 mg; 12.5 mg	
Medicines used in heart failure	Heart failure
Oral > Liquid: 50 mg per 5 mL	
Oral > Solid: 25 mg	
Diuretics	Oedema
Oral > Solid: 25 mg	

Hydrocortisone

Section	Indications
Hormones and antihormones Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)	Lymphoid leukaemia, not elsewhere classified Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia
Adrenal hormones and synthetic substitutes Oral > Solid: 5 mg; 10 mg; 20 mg	Adrenocortical insufficiency
Ophthalmological preparations > Anti-inflammatory agents Local > Ophthalmological > Ointment: 1% (acetate)	Scleritis Anterior uveitis
Antiallergics and medicines used in anaphylaxis Parenteral > General injections > unspecified: 100 mg in vial powder for injection (as sodium succinate)	Anaphylaxis
Anti-inflammatory medicines Local > Rectal > Suppository: 25 mg (acetate) Local > Rectal > Retention enema: 100 mg per 60 mL	Crohn disease site Ulcerative colitis
Dermatological medicines > Anti-inflammatory and antipruritic medicines Local > Topical > Cream: 1% (acetate) Local > Topical > Ointment: 1% (acetate)	Dermatitis or eczema Lichen planus Pruritus due to skin disorder Psoriasis of unspecified type

Hydromorphone

Section	Indications
Opioid analgesics	Therapeutic equivalent to morphine pour Pain

Hydroxocobalamin

Section	Indications
Antianaemia medicines Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)	Megaloblastic anaemia due to vitamin B12 deficiency

Hydroxycarbamide (hydroxyurea)

Section	Indications
Cytotoxic medicines	Squamous cell carcinoma of oropharynx
Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g	Chronic myeloid leukaemia, not elsewhere classified
Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg	
Other medicines for haemoglobinopathies	Other specified sickle cell disorders or other haemoglobinopathies
Oral > Solid > dosage form: 200 mg; 500 mg; 1 g; 100 mg	

Hydroxychloroquine

Section	Indications
Antimalarial medicines > For chemoprevention	Therapeutic equivalent to chloroquine pour
Disease-modifying anti-rheumatic drugs (DMARDs)	Malaria due to Plasmodium falciparum
Oral > Solid: 200 mg (as sulfate)	Systemic lupus erythematosus
	Cutaneous lupus erythematosus of unspecified type

Hyoscine butylbromide

Section	Indications
Medicines for other common symptoms in palliative care	Palliative care
Parenteral > General injections > unspecified: 20 mg per mL injection	

Hyoscine hydrobromide

Section	Indications
Medicines for other common symptoms in palliative care	Palliative care
Local > Topical > Transdermal patch: 1 mg per 72 hours transdermal patch	
Parenteral > General injections > unspecified: 400 µg per mL injection; 600 µg per mL injection	

Hypochlorous acid

Section	Indications
Antiseptics	Disinfectant, not elsewhere classified
Local > Topical > Solution:	

Hypromellose

Section	Indications
Ophthalmological preparations Local > Ophthalmological > Solution (eye drops): 0.3%	Keratoconjunctivitis sicca

Ibrutinib

Section	Indications
Targeted therapies Oral > Solid: 140 mg	Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Ibuprofen

Section	Indications
Medicines administered to the neonate [c]	Patent arterial duct
PARENTERAL > General injections > IV: 5 mg per mL	
NON-OPIOIDS AND NON-STEROIDAL ANTI-INFLAMMATORY MEDICINES (NSAIMS)	Pain
Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL	
Oral > Solid > tablet: 200 mg; 400 mg; 600 mg	
ANTIMIGRAINE MEDICINES > For treatment of acute attack	Migraine
Oral > Liquid: 100 mg per 5 mL; 200 mg per 5 mL	
Oral > Solid > tablet: 200 mg; 400 mg	

Iodoxuridine

Section	Indications
OPHTHALMOLOGICAL PREPARATIONS > Anti-infective agents	
Local > Ophthalmological > Solution (eye drops): 0.1%	Herpes simplex keratitis
Local > Ophthalmological > Ointment: 0.2%	

Ifosfamide

Section	Indications
Cytotoxic medicines	Other specified malignant neoplasms of the ovary
Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection	Germ cell tumour of testis
Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection	Ewing sarcoma of bone and articular cartilage of unspecified sites
	Osteosarcoma of bone and articular cartilage of unspecified sites
	Rhabdomyosarcoma primary site
	Unspecified malignant neoplasms of ill-defined or unspecified sites
	Burkitt lymphoma including Burkitt leukaemia
	Malignant neoplasms of kidney, except renal pelvis
	Anaplastic large cell lymphoma, ALK-negative
	Anaplastic large cell lymphoma, ALK-positive

Imatinib

Section	Indications
Targeted therapies	B lymphoblastic leukaemia or lymphoma with t(9;22) (q34;q11.2); BCR-ABL1
Oral > Solid: 100 mg; 400 mg	Gastrointestinal stromal tumour of unspecified gastrointestinal sites
	Chronic myeloid leukaemia, not elsewhere classified

Imipenem + cilastatin

Section	Indications
Watch group antibiotics	<p>Second choix</p> <p>Therapeutic equivalent to meropenem pour</p> <ul style="list-style-type: none">Neutropenia (high-risk)
Parenteral > General injections > IV: 250 mg (as monohydrate) + 250 mg (as sodium salt) powder for injection; 500 mg (as monohydrate) + 500 mg (as sodium salt) powder for injection	<ul style="list-style-type: none">Therapeutic equivalent to meropenem pourPeritoneal abscess (severe)
	<ul style="list-style-type: none">Therapeutic equivalent to meropenem pourPeritonitis (severe)
Antituberculosis medicines	<p>Autres indications</p> <ul style="list-style-type: none">Pseudomonas aeruginosa resistant to other antibioticAntibiotic resistant Acinetobacter baumannii <p>Therapeutic equivalent to meropenem pour</p> <ul style="list-style-type: none">Multi-drug resistant Mycobacterium tuberculosis

Imipenem + cilastatin + relebactam

Section	Indications
Reserve group antibiotics	<ul style="list-style-type: none">Other specified other antibiotic resistant EnterobacterialesCarbapenem resistant EnterobacterialesCarbapenem-resistant Pseudomonas aeruginosa
Parenteral > General injections > IV: 500 mg + 500 mg + 250 mg powder for injection	

Indapamide

Section	Indications
Antihypertensive medicines	<p>Therapeutic equivalent to hydrochlorothiazide pour</p> <ul style="list-style-type: none">Essential hypertension
Medicines used in heart failure	<p>Therapeutic equivalent to hydrochlorothiazide pour</p> <ul style="list-style-type: none">Heart failure

Indinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 400 mg (as sulfate)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Indometacin

Section	Indications
Medicines administered to the neonate [c]	Therapeutic equivalent to ibuprofen pour Patent arterial duct

Infliximab

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis
	Therapeutic equivalent to adalimumab pour Crohn disease site
	Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Influenza vaccine (seasonal)

Section	Indications
Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against influenza

Insulin

Section	Indications
Insulins Parenteral > General injections > SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen (soluble)	Type 2 diabetes mellitus Type 1 diabetes mellitus

Insulin analogues

Section	Indications
Insulins	Type 1 diabetes mellitus
Not specified	Type 2 diabetes mellitus

Insulin degludec

Section	Indications
Insulins	Therapeutic equivalent to insulin glargine pour
Parenteral > General injections > SC: 100 units per mL	Type 1 diabetes mellitus
	Therapeutic equivalent to long-acting insulin analogues pour
	Type 1 diabetes mellitus
	Therapeutic equivalent to long-acting insulin analogues pour
	Type 2 diabetes mellitus

Insulin detemir

Section	Indications
Insulins	Therapeutic equivalent to insulin glargine pour
	Type 1 diabetes mellitus
	Therapeutic equivalent to long-acting insulin analogues pour
	Type 1 diabetes mellitus
	Therapeutic equivalent to long-acting insulin analogues pour
	Type 2 diabetes mellitus

Insulin glargine

Section	Indications
Insulins	Type 1 diabetes mellitus
Parenteral > General injections > SC: 100 units per mL	Therapeutic equivalent to long-acting insulin analogues pour
	Type 1 diabetes mellitus
	Therapeutic equivalent to long-acting insulin analogues pour
	Type 2 diabetes mellitus

Intermediate-acting insulin

Section	Indications
Insulins Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin	Type 1 diabetes mellitus Type 2 diabetes mellitus

Intraperitoneal dialysis solution

Section	Indications
Peritoneal dialysis solution parenteral solution	Care involving peritoneal dialysis

Iodine

Section	Indications
Vitamins and minerals Parenteral > General injections > IM: 0.5 mL (240 mg iodine) iodized oil in ampoule; 1 mL (480 mg iodine) iodized oil in ampoule	Iodine deficiency
Oral > Liquid: 1 mL (480 mg iodine) iodized oil in ampoule; 0.5 mL (240 mg iodine) iodized oil in ampoule; 0.57 mL (308 mg iodine) iodized oil in dispenser bottle	
Oral > Solid: 190 mg	
Antiseptics	Therapeutic equivalent to povidone iodine pour Povidone iodine

Iohexol

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule	X-ray contrast media, iodinated

Ipecacuanha

Section	Indications
Antidotes and other substances used in poisonings > Non-specific Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine	Unspecified injury, poisoning or certain other consequences of external causes

Ipratropium bromide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 20 µg per dose	Asthma Chronic obstructive pulmonary disease

Irinotecan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site

Isoflurane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > liquid:	Anaesthetics and therapeutic gases

Isoniazid

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 100 mg per mL Oral > Solid > dispersible tablet: 100 mg Oral > Solid > tablet: 100 mg; 300 mg	Other specified tuberculosis Tuberculosis

Isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 50 mg + 150 mg + 75 mg	Tuberculosis

Isoniazid + pyridoxine + sulfamethoxazole + trimethoprim

Section	Indications
Medicines for prevention of HIV-related opportunistic infections Oral > Solid: 300 mg + 25 mg + 800 mg + 160 mg tablet (scored)	Other specified prophylactic measures

Isoniazid + rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid > dispersible tablet: 50 mg + 75 mg	
Oral > Solid > tablet: 75 mg + 150 mg; 150 mg + 300 mg	

Isoniazid + rifapentine

Section	Indications
Antituberculosis medicines	Latent tuberculosis
Oral > Solid: 300 mg + 300 mg tablet (scored)	

Isoprenaline

Section	Indications
Antiarrhythmic medicines	Bradycardia
Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)	Cardiogenic shock
Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)	

Isosorbide dinitrate

Section	Indications
Antianginal medicines	Angina pectoris
Oral > Solid: 5 mg sublingual tablet	

Itraconazole

Section	Indications
Antifungal medicines	Chronic pulmonary aspergillosis
Oral > Liquid: 10 mg per mL	Chromoblastomycosis
Oral > Solid: 100 mg	Histoplasmosis
	Talaromycosis
	Sporotrichosis
	Paracoccidioidomycosis

Ivermectin

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 3 mg	Strongyloidiasis Ascariasis Trichuriasis Hookworm diseases Ancylostomiasis
Antifilarials Oral > Solid > tablet: 3 mg	Onchocerciasis Lymphatic filariasis
Medicines for ectoparasitic infections Oral > Solid > tablet: 3 mg	Scabies

Kanamycin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis
	Therapeutic equivalent to gentamicin pour Infectious blepharitis

Kanamycin (injection)

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1000 mg powder for injection (vial)	Multi-drug resistant Mycobacterium tuberculosis

Ketamine

Section	Indications
General anaesthetics and oxygen > Injectable medicines Parenteral > General injections > unspecified: 50 mg per mL in 10 mL vial (as hydrochloride)	Anaesthetics and therapeutic gases

Ketoconazole

Section	Indications
Medicines for endocrine disorders Oral > Solid > tablet: 200 mg	Cushing syndrome

Lactulose

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 3.1 to 3.7 g per 5 mL oral liquid	Palliative care

Lamivudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 50 mg per 5 mL Oral > Solid: 150 mg tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + nevirapine + zidovudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals	Therapeutic equivalent to emtricitabine + tenofovir pour Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Therapeutic equivalent to emtricitabine + tenofovir pour

Contact with or exposure to human immunodeficiency virus

Lamivudine + zidovudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamotrigine

Section	Indications
Antiseizure medicines Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet; 200 mg tablet; 2 mg tablet (chewable, dispersible); 5 mg tablet (chewable, dispersible); 25 mg tablet (chewable, dispersible); 50 mg tablet (chewable, dispersible); 100 mg tablet (chewable, dispersible); 200 mg tablet (chewable, dispersible)	Epilepsy or seizures

Latanoprost

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 50 µg per mL eye drops	Primary open-angle glaucoma

Ledipasvir + sofosbuvir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 90 mg + 400 mg tablet	Chronic hepatitis C

Lenalidomide

Section	Indications
Immunomodulators Oral > Solid: 25 mg	Plasma cell myeloma

Letrozole

Section	Indications
Ovulation inducers Oral > Solid > dosage form: 2.5 mg	Female infertility without specification whether primary or secondary

Leuprorelin

Section	Indications
Hormones and antihormones	Malignant neoplasms of breast
Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe	Malignant neoplasms of prostate

Levamisole

Section	Indications
Intestinal antihelminthics Oral > Solid > tablet: 50 mg (as hydrochloride); 150 mg (as hydrochloride)	Ascariasis

Levetiracetam

Section	Indications
Antiseizure medicines	Epilepsy or seizures
Oral > Liquid: 100 mg per mL	
Oral > Solid > tablet: 250 mg; 500 mg; 750 mg; 1000 mg	Status epilepticus
Parenteral > General injections > IV: 500 mg per 5 mL in 5 mL vial; 5 mg per mL in 100 mL bag; 10 mg per mL in 100 mL bag; 15 mg per mL in 100 mL bag	

Levodopa

Section	Indications
Antiseizure medicines	Parkinson disease
Oral > Solid: 250 mg	

Levodopa + benserazide

Section	Indications
Medicines for parkinsonism	Therapeutic equivalent to levodopa + carbidopa pour Parkinson disease

Levodopa + carbidopa

Section	Indications
Medicines for parkinsonism	Parkinson disease
Oral > Solid: 100 mg + 10 mg tablet; 250 mg + 25 mg tablet; 100 mg + 25 mg tablet	

Levofloxacin

Section	Indications
Antituberculosis medicines	Multi-drug resistant Mycobacterium tuberculosis
Oral > Solid > dispersible tablet: 100 mg	
Oral > Solid > tablet: 250 mg; 500 mg; 750 mg	

Levonorgestrel

Section	Indications
Oral hormonal contraceptives Oral > Solid: 750 µg [2]; 1.5 mg [1]	Contact with health services for postcoital contraception
Oral > Solid: 30 µg	Contact with health services for contraceptive management
Intrauterine devices Intrauterine system with reservoir containing 52 mg of levonorgestrel.	Contact with health services for insertion of contraceptive device

Levonorgestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 75 mg per rod (two-rods)	Contact with health services for insertion of contraceptive device

Levothyroxine

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 µg (sodium salt); 100 µg (sodium salt); 25 µg (sodium salt)	Hypothyroidism

Lidocaine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)	Ventricular tachyarrhythmia
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 5% in 2 mL ampoule (hydrochloride) + 7.5% glucose solution Parenteral > Locoregional injections > Other: 1% in vial (hydrochloride); 2% in vial (hydrochloride) Local > Topical > unspecified: 2 to 4% (hydrochloride)	Local anaesthetics

Lidocaine + epinephrine

Section	Indications
<p>Local anaesthetics</p> <p>Parenteral > Locoregional injections > Other: 1% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial; 2% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial</p> <p>Local > Dental > Cartridge: 2% (lidocaine hydrochloride) + 1:80 000 epinephrine</p>	Local anaesthetics

Lindane

Section	Indications
<p>Dermatological medicines > Scabicides and pediculicides</p> <p>Local > Topical > Cream: 1%</p> <p>Local > Topical > Lotion: 1%</p>	Scabies

Linezolid

Section	Indications
<p>Reserve group antibiotics</p> <p>Parenteral > General injections > IV: 2 mg per mL in 300 mL bag</p> <p>Oral > Liquid: 100 mg per 5 mL powder for oral liquid</p> <p>Oral > Solid: 400 mg tablet; 600 mg tablet</p> <p>Oral > Solid > dispersible tablet: 150 mg</p> <p>Oral > Solid > tablet: 600 mg</p>	<p>Other specified bacterial diseases</p> <p>Methicillin resistant <i>Staphylococcus aureus</i></p> <p>Vancomycin resistant <i>Staphylococcus aureus</i></p> <p>Vancomycin resistant <i>Enterococcus</i></p>
<p>Antituberculosis medicines</p> <p>Oral > Solid > dispersible tablet: 150 mg</p> <p>Oral > Solid > tablet: 600 mg</p>	Multi-drug resistant <i>Mycobacterium tuberculosis</i>

Liraglutide

Section	Indications
<p>Medicines for endocrine disorders</p> <p>Parenteral > General injections > SC: 6 mg per mL in 3 mL pre-filled pen</p>	Obesity

Lisinopril + amlodipine

Section	Indications
<p>Antihypertensive medicines</p> <p>Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg</p>	Essential hypertension

Lisinopril + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg	Essential hypertension

Lisocabtagene maraleucel

Section	Indications
Antineoplastics and supportive medicines	Therapeutic equivalent to axicabtagene ciloleucel pour Diffuse large B-cell lymphomas

Lithium carbonate

Section	Indications
Medicines for bipolar disorders Oral > Solid: 300 mg	Bipolar or related disorders

Long-acting insulin analogues

Section	Indications
Insulins Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen	Type 1 diabetes mellitus Type 2 diabetes mellitus

Loperamide

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid > dosage form: 2 mg	Palliative care

Lopinavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid > tablet: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg tablet (heat-stable) Oral > Solid > dosage form: 40 mg + 10 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Loratadine

Section	Indications
Antiallergics and medicines used in anaphylaxis Oral > Liquid: 1 mg per mL Oral > Solid: 10 mg	Allergic or hypersensitivity conditions of unspecified type

Lorazepam

Section	Indications
Antiseizure medicines Parenteral > General injections > IV: 2 mg per mL in 1 mL ampoule; 4 mg per mL in 1 mL ampoule	Status epilepticus
Medicines for anxiety disorders	Therapeutic equivalent to diazepam pour Anxiety

Losartan

Section	Indications
Antihypertensive medicines Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet	Essential hypertension
Medicines used in heart failure Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet	Heart failure

Lovastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia
	Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Lugol's solution

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Liquid: 13% w/v (about 130 mg/mL total iodine)	Thyrotoxicosis

Magnesium sulfate

Section	Indications
Antiseizure medicines Parenteral > General injections > IV: 500 mg per mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume)	Severe pre-eclampsia Eclampsia
Parenteral > General injections > IM: 500 mg per mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume)	

Mannitol

Section	Indications
Diuretics Parenteral > General injections > IV: 10% solution; 20% solution	Ocular hypertension Cerebral oedema

Measles vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against measles alone

Mebendazole

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)	Ascariasis Enterobiasis Trichuriasis Hookworm diseases
Cysticidal medicines Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)	Other specified echinococcosis

Medroxyprogesterone acetate

Section	Indications
Injectable hormonal contraceptives Parenteral > General injections > IM: 150 mg per mL in 1 mL vial Parenteral > General injections > SC: 104 mg per 0.65 mL in pre-filled syringe; 104 mg per 0.65 mL in single dose injection delivery system	Contact with health services for reasons associated with reproduction
Progestogens Oral > Solid: 5 mg	Excessive menstruation with irregular cycle

Mefloquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 250 mg tablet (hydrochloride)	Malaria due to <i>Plasmodium falciparum</i> Malaria due to <i>Plasmodium vivax</i>
Antimalarial medicines > For chemoprevention Oral > Solid: 250 mg tablet (hydrochloride)	Malaria due to <i>Plasmodium falciparum</i>

Meglumine antimoniate

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule	Visceral leishmaniasis Cutaneous leishmaniasis Mucocutaneous leishmaniasis

Meglumine iotroxate

Section	Indications
Diagnostic agents > Radiocontrast media Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL	X-ray contrast media, iodinated

Melarsoprol

Section	Indications
Medicines for the treatment of 2nd stage African trypanosomiasis Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)	African trypanosomiasis

Melphalan

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection Oral > Solid: 2 mg	Plasma cell myeloma

Meningococcal meningitis vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Mercaptopurine

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Oral > Solid: 50 mg tablet	Acute myeloid leukaemia with recurrent genetic abnormalities
Oral > Liquid: 20 mg per mL	Lymphoid leukaemia, not elsewhere classified
	Langerhans cell histiocytosis

Meropenem

Section	Indications
Watch group antibiotics	Second choix
PARENTERAL > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Neutropenia (high-risk)
	Peritoneal abscess (severe)
	Peritonitis (severe)
	Neonatal meningitis
Antituberculosis medicines	Multi-drug resistant <i>Mycobacterium tuberculosis</i>
PARENTERAL > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	

Meropenem + vaborbactam

Section	Indications
Reserve group antibiotics	Carbapenem-resistant <i>Pseudomonas aeruginosa</i>
PARENTERAL > General injections > IV: 1 g in vial + 1 g in vial powder for injection	Carbapenem resistant <i>Enterobacteriales</i>
	Carbapenem resistant <i>Acinetobacter baumannii</i>

Mesalazine

Section	Indications
Anti-inflammatory medicines	Therapeutic equivalent to sulfasalazine pour Crohn disease site
	Therapeutic equivalent to sulfasalazine pour Ulcerative colitis

Mesna

Section	Indications
<p>Supportive medicines</p> <p>Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule</p> <p>Oral > Solid: 400 mg; 600 mg</p>	<p>Osteosarcoma of bone and articular cartilage of other specified sites</p> <p>Other specified malignant neoplasms of the ovary</p> <p>Germ cell tumour of testis</p> <p>Ewing sarcoma of bone and articular cartilage of unspecified sites</p> <p>Rhabdomyosarcoma primary site</p> <p>Unspecified malignant neoplasms of ill-defined or unspecified sites</p> <p>Burkitt lymphoma including Burkitt leukaemia</p> <p>Malignant neoplasms of kidney, except renal pelvis</p>

Metformin

Section	Indications
<p>Oral hypoglycaemic agents</p> <p>Oral > Solid: 500 mg (hydrochloride)</p>	Type 2 diabetes mellitus

Methadone

Section	Indications
<p>Medicines for opioid use disorders</p> <p>Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)</p>	Opioid dependence
<p>Opioid analgesics</p> <p>Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)</p>	Chronic cancer pain
<p>Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)</p>	

Methimazole

Section	Indications
<p>Thyroid hormones and antithyroid medicines</p> <p>Oral > Solid: 5 mg; 10 mg; 20 mg</p>	Thyrotoxicosis

Methionine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg (DL)	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methotrexate

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial (as sodium salt) Oral > Solid: 2.5 mg (as sodium salt) Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection	Unspecified malignant neoplasms of ill-defined or unspecified sites Lymphoid leukaemia, not elsewhere classified Acute myeloid leukaemia with recurrent genetic abnormalities Malignant trophoblastic neoplasms of placenta Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis
Dermatological medicines > Medicines affecting skin differentiation and proliferation Oral > Solid > tablet: 2.5 mg (as sodium salt); 10 mg (as sodium salt)	Psoriasis of unspecified type
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 2.5 mg (as sodium salt)	Rheumatoid arthritis, serology unspecified
Immunomodulators for non-malignant disease Parenteral > General injections > SC: pre-filled syringe in multiple strengths	Crohn disease site Psoriatic arthritis
Medicines for juvenile joint diseases Oral > Solid > tablet: 2.5 mg (as sodium salt)	Juvenile idiopathic arthritis

Methoxy polyethylene glycol-epoetin beta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents pour Anaemia due to chronic disease

Methyldopa

Section	Indications
Antihypertensive medicines Oral > Solid: 250 mg	Essential hypertension Gestational hypertension

Methylergometrine

Section	Indications
Uterotonics	Therapeutic equivalent to ergometrine pour Postpartum haemorrhage

Methylphenidate

Section	Indications
Medicines for mental and behavioural disorders Multiple formulations and strengths.	Attention deficit hyperactivity disorder

Methylprednisolone

Section	Indications
Hormones and antihormones Parenteral > General injections > IV: 40 mg per mL in 1 mL single-dose vial (as sodium succinate); 40 mg per mL in 5 mL multi-dose vial (as sodium succinate); 80 mg per mL in 1 mL single-dose vial (as sodium succinate)	Lymphoid leukaemia, not elsewhere classified Burkitt lymphoma including Burkitt leukaemia

Methylthioninium chloride

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule	Acquired methaemoglobinuria

Metoclopramide

Section	Indications
Medicines for other common symptoms in palliative care	Palliative care
Oral > Liquid: 5 mg per 5 mL	
Oral > Solid: 10 mg (hydrochloride)	
Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	
Antiemetic medicines	Nausea or vomiting
Oral > Liquid: 5 mg per 5 mL	
Oral > Solid: 10 mg (hydrochloride)	
Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	

Metoprolol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Metronidazole

Section	Indications
<p>Access group antibiotics</p> <p>Oral > Liquid: 200 mg per 5 mL (as benzoate)</p> <p>Oral > Solid: 200 to 500 mg tablet</p> <p>Parenteral > General injections > unspecified: 500 mg per 100 mL in vial Injection</p> <p>Local > Rectal > Suppository: 500 mg; 1 g</p> <p>Parenteral > General injections > unspecified: 500 mg per 100 mL in vial</p> <p>Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg</p>	<p>Premier choix</p> <p>Intestinal infections due to <i>Clostridioides difficile</i></p> <p>Trichomoniasis</p> <p>co-prescrite avec cefazolin</p> <p>Other specified prophylactic measures</p> <p>co-prescrite avec cefotaxime</p> <p>Peritonitis (severe)</p> <p>co-prescrite avec cefotaxime</p> <p>Peritonitis (mild-moderate)</p> <p>co-prescrite avec cefotaxime</p> <p>Peritoneal abscess (mild-moderate)</p> <p>co-prescrite avec cefotaxime</p> <p>Peritoneal abscess (severe)</p> <p>co-prescrite avec ceftriaxone</p> <p>Necrotising fasciitis</p> <p>co-prescrite avec ceftriaxone</p> <p>Peritonitis (severe)</p> <p>co-prescrite avec ceftriaxone</p> <p>Peritoneal abscess (mild-moderate)</p> <p>co-prescrite avec ceftriaxone</p> <p>Peritoneal abscess (severe)</p> <p>co-prescrite avec ceftriaxone</p> <p>Peritonitis (mild-moderate)</p> <p>Second choix</p> <p>co-prescrite avec ciprofloxacin</p> <p>Peritonitis (mild-moderate)</p> <p>co-prescrite avec ciprofloxacin</p> <p>Peritoneal abscess (mild-moderate)</p> <p>Autres indications</p> <p>Bacterial infection of unspecified site</p> <p>Amoebiasis</p>
<p>Antiamoebic and antigiardiasis medicines</p> <p>Oral > Liquid: 200 mg per 5 mL (as benzoate)</p> <p>Parenteral > General injections > unspecified: 500 mg in 100 mL vial</p> <p>Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg</p>	

Micafungin

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection	Systemic or invasive candidosis

Miconazole

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Cream: 2% (nitrate) Local > Topical > Ointment: 2% (nitrate)	Fungal infection of the skin

Midazolam

Section	Indications
Antiseizure medicines Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration); 5 mg per mL in 1 mL vial (for buccal administration); 5 mg per mL in 3 mL vial (for buccal administration)	Therapeutic equivalent to lorazepam pour Status epilepticus Status epilepticus
Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)	
Preoperative medication and sedation for short-term procedures Parenteral > General injections > IV: 1 mg per mL	Sedative-hypnotic and anxiolytic drugs
Oral > Liquid: 2 mg per mL	
Oral > Solid > dosage form: 15 mg; 7.5 mg	
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per mL	Palliative care
Parenteral > General injections > unspecified: 1 mg per mL; 5 mg per mL	
Oral > Solid > dosage form: 7.5 mg; 15 mg	

Mifepristone - misoprostol

Section	Indications
Uterotonics Oral > Solid: 200 mg + 200 µg; 200 mg [1] + 200 µg [4] in co-package	Induced abortion Unspecified time of fetal death, cause not specified

Miltefosine

Section	Indications
Antileishmaniasis medicines Oral > Solid: 10 mg; 50 mg	Visceral leishmaniasis Mucocutaneous leishmaniasis Cutaneous leishmaniasis

Misoprostol

Section	Indications
Uterotonics Oral > Solid: 200 µg Local > Vaginal > tablet: 25 µg	Spontaneous abortion, incomplete, without complication Unspecified obstetric condition Postpartum haemorrhage

Mometasone

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide pour Asthma

Mometasone + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Morphine

Section	Indications
<p>Opioid analgesics</p> <p>Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate)</p> <p>Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate)</p> <p>Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate)</p> <p>Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)</p> <p>Preoperative medication and sedation for short-term procedures</p> <p>Parenteral > General injections > IV: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)</p>	<p>Pain</p> <p>Opioids or related analgesics and agents affecting opioid receptors</p>

Moxifloxacin

Section	Indications
<p>Antituberculosis medicines</p> <p>Oral > Solid > dispersible tablet: 100 mg</p> <p>Oral > Solid > tablet: 400 mg</p>	<p>Multi-drug resistant Mycobacterium tuberculosis</p> <p>Tuberculosis</p>

Multiple micronutrient powder

Section	Indications
<p>Vitamins and minerals</p> <p>Sachets containing:</p> <ul style="list-style-type: none">- iron (elemental) 12.5 mg (as coated ferrous fumarate)- zinc (elemental) 5 mg- vitamin A 300 micrograms- with or without other micronutrients at recommended daily values	<p>Iron deficiency anaemia</p>

Multiple micronutrient supplement

Section	Indications
<p>Other medicines administered to the mother Tablet containing:</p> <p>Vit A: 800 mcg retinol activity equivalent Vit C: 70 mg Vit D: 5 mcg (200 IU) Vit E: 10 mg alpha tocopherol equivalent Vit B1: 1.4 mg Vit B2: 1.4 mg Vit B3: 18 mg niacin equivalent Vit B6: 1.9 mg Folic acid: 400 mcg Vit B12: 2.6 mcg Iron: 30 mg Iodine: 150 mcg Zinc: 15 mg Selenium: 65 mcg Copper: 2 mg</p>	<p>Maternal care for other specified conditions predominantly related to pregnancy</p>

Mumps vaccine

Section	Indications
<p>Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.</p>	<p>Need for immunization against mumps alone</p>

Mupirocin

Section	Indications
<p>Dermatological medicines > Anti-infective medicines Local > Topical > Cream: 2% (as mupirocin calcium) Local > Topical > Ointment: 2%</p>	<p>Superficial bacterial folliculitis Impetigo</p>

Nadroparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin pour Acute ischaemic heart disease
	Therapeutic equivalent to enoxaparin pour Venous thromboembolism

Nalidixic acid

Section	Indications
Anti-infective medicines Oral > Solid: 250 mg; 500 mg	Shigella resistant to unspecified antibiotic

Naloxone

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)	Unintentional exposure to or harmful effects of opioids or related analgesics

Naltrexone

Section	Indications
Medicines for alcohol use disorders Parenteral > General injections > IM: 380 mg in vial (extended-release) Oral > Solid > tablet: 50 mg	Disorders due to use of alcohol

Natamycin

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Suspension: 5% eye drops	Infectious keratitis

Nelfinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 250 mg (as mesilate) Oral > Other: 50 mg per g oral powder	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Neostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Oral > Solid: 15 mg (neostigmine bromide) Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate); 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate)	Myasthenia gravis Anticholinesterase reversible

Netilmicin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis
	Therapeutic equivalent to gentamicin pour Infectious blepharitis

Nevirapine

Section	Indications
Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors Oral > Liquid: 50 mg per 5 mL Oral > Solid > dispersible tablet: 50 mg Oral > Solid > tablet: 200 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Niclosamide

Section	Indications
Intestinal anthelmintics Oral > Solid: 500 mg tablet (chewable)	Hymenolepiasis Diphyllobothriasis Taeniasis due to Taenia saginata Taeniasis due to Taenia solium

Nicotinamide

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg	Pellagra

Nicotine replacement therapy

Section	Indications
<p>Medicines for nicotine use disorders</p> <p>Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour</p> <p>Oral > Solid > lozenge: 2 mg; 4 mg lozenge</p> <p>Local > Buccal > Oral spray: 1 mg per actuation</p> <p>Local > Buccal > Chewing gum: 2 mg; 4 mg</p>	Nicotine dependence

Nifedipine

Section	Indications
<p>Antioxytocos (tocolytics)</p> <p>Oral > Solid: 10 mg (immediate-release)</p>	Preterm labour without delivery
<p>Antianginal medicines</p> <p>Oral > Solid: 10 mg</p>	Angina pectoris
<p>Antihypertensive medicines</p> <p>Oral > Solid: 10 mg tablet; sustained-release formulations</p>	Essential hypertension

Nifurtimox

Section	Indications
<p>American trypanosomiasis</p> <p>Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)</p>	Chagas disease
<p>Medicines for the treatment of 2nd stage African trypanosomiasis</p> <p>Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)</p>	African trypanosomiasis

Nilotinib

Section	Indications
<p>Targeted therapies</p> <p>Oral > Solid > capsule: 150 mg; 200 mg</p>	Chronic myeloid leukaemia, not elsewhere classified

Nilutamide

Section	Indications
<p>Hormones and antihormones</p>	<p>Therapeutic equivalent to bicalutamide pour Malignant neoplasms of prostate</p>

Nitrofurantoin

Section	Indications
Access group antibiotics Oral > Liquid: 25 mg per 5 mL oral liquid Oral > Solid > dosage form: 100 mg; 50 mg	Premier choix Infectious cystitis

Nitrous oxide

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation:	Anaesthetics and therapeutic gases

Nivolumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion	Other specified malignant neoplasms of bronchus or lung Melanoma of skin

Norethisterone

Section	Indications
Oral hormonal contraceptives Oral > Solid: 350 µg	Contact with health services for contraceptive management
Progestogens Oral > Solid: 5 mg	Excessive menstruation with irregular cycle Therapeutic equivalent to medroxyprogesterone acetate pour Excessive menstruation with irregular cycle

Norethisterone enantate

Section	Indications
Injectable hormonal contraceptives Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule oily solution	Contact with health services for reasons associated with reproduction

Normal immunoglobulin

Section	Indications
Human immunoglobulins	
Parenteral > General injections > IV: 5% protein solution; 10% protein solution	Primary immunodeficiencies
Parenteral > General injections > IM: 16% protein solution	Langerhans cell histiocytosis
Parenteral > General injections > SC: 15% protein solution; 16% protein solution	Mucocutaneous lymph node syndrome

Nystatin

Section	Indications
Antifungal medicines	
Oral > Liquid: 100000 IU per mL	Candidosis
Local > Topical > Other: 100000 IU pessary	
Oral > Solid > dosage form: 500000 IU	
Oral > Solid > lozenge: 100000 IU	

Ocrelizumab

Section	Indications
Medicines for multiple sclerosis	
Parenteral > General injections > IV: 30 mg per mL in 10 vial concentrate for solution	Multiple sclerosis
Parenteral > General injections > IV: 30 mg per mL in 10 mL vial	Relapsing-remitting multiple sclerosis

Octreotide

Section	Indications
Medicines for disorders of the pituitary hormone system	
Parenteral > General injections > IM: 20 mg in vial (modified-release, as acetate) plus diluent	Acromegaly or pituitary gigantism
Parenteral > General injections > SC: 0.05 mg per mL in 1 mL vial (immediate-release, as acetate); 0.1 mg per mL in 1 mL vial (immediate-release, as acetate); 0.5 mg per mL in 1 mL vial (immediate-release, as acetate)	

Ofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant <i>Mycobacterium tuberculosis</i>
Ear, nose and throat medicines [c]	Therapeutic equivalent to ciprofloxacin pour Infectious diseases of external ear
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.3%	Other specified conjunctivitis Infectious keratitis

Olanzapine

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 10 mg in vial powder for injection	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Omadacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 100 mg lyophilized powder for injection Oral > Solid: 300 mg	Methicillin resistant <i>Staphylococcus aureus</i>

Ombitasvir + paritaprevir + ritonavir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 12.5 mg + 75 mg + 50 mg tablet	Chronic hepatitis C

Omeprazole

Section	Indications
Antiulcer medicines Oral > Liquid: 20 mg powder for oral liquid; 40 mg powder for oral liquid Oral > Solid: 10 mg; 20 mg; 40 mg Parenteral > General injections > IV: 40 mg in vial	Peptic ulcer, site unspecified Gastro-oesophageal reflux disease Gastrointestinal bleeding

Ondansetron

Section	Indications
<p>Medicines for other common symptoms in palliative care</p> <p>Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)</p> <p>Oral > Liquid: 4 mg per 5 mL (ondansetron base)</p> <p>Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base)</p>	Palliative care
<p>Antiemetic medicines</p> <p>Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)</p> <p>Oral > Liquid: 4 mg per 5 mL (ondansetron base)</p> <p>Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base); 24 mg (ondansetron base)</p>	Nausea or vomiting

Oral rehydration salts

Section	Indications
<p>Oral rehydration</p> <p>Powder for dilution in 200 mL, 500 mL, 1 L:</p> <ul style="list-style-type: none">- glucose 75 mEq- sodium 75 mEq or mmol/L- chloride 65 mEq or mmol/L- potassium 20 mEq or mmol/L- citrate 10 mmol/L- osmolarity 245 mOsm/L- glucose 13.5 g/L- sodium chloride 2.6 g/L- potassium chloride 1.5 g/L- trisodium citrate dihydrate 2.9 g/L	Diarrhoea
<p>Solutions correcting water, electrolyte and acid-base disturbances > Oral</p> <p>Powder for dilution in 200 mL, 500 mL, 1 L:</p> <ul style="list-style-type: none">- glucose 75 mEq- sodium 75 mEq or mmol/L- chloride 65 mEq or mmol/L- potassium 20 mEq or mmol/L- citrate 10 mmol/L- osmolarity 245 mOsm/L- glucose 13.5 g/L- sodium chloride 2.6 g/L- potassium chloride 1.5 g/L- trisodium citrate dihydrate 2.9 g/L	Other specified disorders of fluid, electrolyte or acid-base balance

Oral rehydration salts - zinc sulfate

Section	Indications
<p>Medicines used in diarrhoea</p> <p>Oral > Other: ORS powder for dilution and zinc sulfate solid oral dosage form 20 mg (co-packaged)</p>	Diarrhoea

Oseltamivir

Section	Indications
<p>Other antivirals</p> <p>Oral > Solid: 30 mg capsule (as phosphate); 45 mg capsule (as phosphate); 75 mg capsule (as phosphate)</p>	Influenza due to identified seasonal influenza virus

Osimertinib

Section	Indications
Targeted therapies Oral > Solid: 40 mg (as mesylate); 80 mg (as mesylate)	Other specified malignant neoplasms of bronchus or lung

Oxaliplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial	Malignant neoplasm metastasis in large intestine Malignant neoplasms of colon

Oxamniquine

Section	Indications
Antischistosomals and other antitrematode medicines Oral > Liquid: 250 mg per 5 mL Oral > Solid > capsule: 250 mg	Schistosomiasis due to Schistosoma mansoni

Oxazolidinones

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases

Oxycodone

Section	Indications
Opioid analgesics	Therapeutic equivalent to morphine pour Pain

Oxygen

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation: (medical gas)	Anaesthetics and therapeutic gases
Medical gases Respiratory > Inhalation:	Respiratory failure as acute or chronic

Oxytetracycline

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to tetracycline pour Other specified conjunctivitis
	Therapeutic equivalent to tetracycline pour Infectious keratitis
	Therapeutic equivalent to tetracycline pour Infectious blepharitis
	Therapeutic equivalent to tetracycline pour Trachoma

Oxytocin

Section	Indications
Uterotonics Parenteral > General injections > unspecified: 10 IU per mL	Postpartum haemorrhage

P-aminosalicylate sodium

Section	Indications
Antituberculosis medicines Oral > Liquid: 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid) powder for oral solution	Multi-drug resistant Mycobacterium tuberculosis

P-aminosalicylic acid

Section	Indications
Antituberculosis medicines Oral > Solid: 4 g granules in sachet	Multi-drug resistant Mycobacterium tuberculosis

Paclitaxel

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 6 mg per mL	Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Kaposi sarcoma of unspecified primary site Other specified malignant neoplasms of breast Other specified carcinomas of ovary Malignant neoplasms of breast Malignant neoplasms of cervix uteri

Palbociclib

Section	Indications
Targeted therapies Oral > Solid: 75 mg; 100 mg; 125 mg	Other specified malignant neoplasms of breast

Paliperidone

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 25 mg in pre-filled syringe (as palmitate); 50 mg in pre-filled syringe (as palmitate); 75 mg in pre-filled syringe (as palmitate); 100 mg in pre-filled syringe (as palmitate); 150 mg in pre-filled syringe (as palmitate)	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Palonosetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Pancreatic enzymes

Section	Indications
Pancreatic enzymes Oral > Solid: Age-appropriate formulations and doses including lipase, protease and amylase.	Enzyme intestinal

Paracetamol (acetaminophen)

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 100 mg; 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg Antimigraine medicines > For treatment of acute attack Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Pain Migraine

Paromomycin

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)	Visceral leishmaniasis
Antiamoebic and antigiardiasis medicines Oral > Liquid: 125 mg per 5 mL as sulfate Oral > Solid: 250 mg as sulfate	Amoebiasis

Pegasparagase

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection	Lymphoid leukaemia, not elsewhere classified

Pegfilgrastim

Section	Indications
Immunomodulators Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe	Acquired neutropaenia

Pegylated interferon alfa (2a)

Section	Indications
Medicines for hepatitis C > Other antivirals for hepatitis C Parenteral > General injections > SC: 180 µg in vial; 180 µg in prefilled syringe	Chronic hepatitis C

Pegylated interferon alfa (2b)

Section	Indications
Medicines for hepatitis C > Other antivirals for hepatitis C Parenteral > General injections > SC: 80 µg in prefilled syringe; 100 µg in prefilled syringe	Chronic hepatitis C

Pembrolizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 50 mg powder for injection	Therapeutic equivalent to nivolumab pour Melanoma of skin Other specified malignant neoplasms of bronchus or lung

Penicillamine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 250 mg	Rheumatoid arthritis, serology unspecified

Pentamidine

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)	Visceral leishmaniasis
Medicines for the treatment of 1st stage African trypanosomiasis Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection	African trypanosomiasis
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid > tablet: 200 mg (as isethionate); 300 mg (as isethionate)	Pneumocystosis

Permethrin

Section	Indications
Dermatological medicines > Scabicides and pediculicides	Pediculosis corporis
Local > Topical > Lotion: 1%	Pediculosis capitis
Local > Topical > Cream: 5%	Scabies

Pertussis vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against pertussis alone

Pertuzumab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 240 mg per 14 mL in vial concentrated solution	Carcinoma of breast, specialised type

Phenelzine

Section	Indications
Medicines for depressive disorders Oral > Solid > dosage form: 15 mg (as sulfate)	Other specified depressive disorders

Phenobarbital

Section	Indications
Antiseizure medicines Oral > Liquid: 15 mg per 5 mL Oral > Solid: 15 to 100 mg Parenteral > General injections > IV: 200 mg per mL (phenobarbital sodium); 30 mg per mL (phenobarbital sodium); 60 mg per mL (phenobarbital sodium)	Epilepsy or seizures Status epilepticus

Phenoxymethylpenicillin

Section	Indications
Access group antibiotics	Premier choix
Oral > Liquid: 250 mg per 5 mL (as potassium salt) powder for oral liquid	Acute pharyngitis
Oral > Solid: 250 mg (as potassium salt) tablet	Periapical abscess without sinus
Oral > Solid: 250 mg (as potassium salt); 500 mg (as potassium salt)	Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)
	Autres indications
	Bacterial infection of unspecified site

Phenytoin

Section	Indications
Antiseizure medicines	Epilepsy or seizures
Oral > Liquid: 30 mg per 5 mL (phenytoin)	
Parenteral > General injections > unspecified: 50 mg per mL in vial (phenytoin sodium)	
Oral > Solid > tablet: 50 mg (phenytoin) (chewable)	
Oral > Solid > dosage form: 25 mg (phenytoin sodium); 50 mg (phenytoin sodium); 100 mg (phenytoin sodium)	

Phosphorus

Section	Indications
Medicines for endocrine disorders	Hypophosphataemic rickets
Oral > Other: 500 mg in sachet (elemental phosphorus) granules	
Oral > Solid > tablet: 250 mg (elemental phosphorus); 500 mg (elemental phosphorus); 500 mg (elemental phosphorus) (effervescent)	

Phytomenadione

Section	Indications
Medicines affecting coagulation	Haemorrhagic disorder due to other specified circulating anticoagulants
Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule	
Oral > Solid: 10 mg tablet	Haemorrhagic or haematological disorders of fetus or newborn

Pilocarpine

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 2% (hydrochloride or nitrate); 4% (hydrochloride or nitrate)	Acute angle closure with pupillary block Ocular hypertension Primary open-angle glaucoma

Piperacillin + tazobactam

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 2 g (as sodium salt) + 250 mg (as sodium salt) powder for injection; 4 g (as sodium salt) + 500 mg (as sodium salt) powder for injection	Premier choix Peritoneal abscess (severe) Other specified pneumonia (Hospital-acquired pneumonia) Neutropenia (high-risk) Peritonitis (severe) co-prescrite avec clindamycin Necrotising fasciitis

Platelets

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Plazomicin

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500 mg per 10 mL	Carbapenem resistant Acinetobacter baumannii Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacteriales

Pneumococcal vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Podophyllotoxin

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to podophyllum resin pour Anogenital warts
	Therapeutic equivalent to podophyllum resin pour Plantar warts

Podophyllum resin

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Anogenital warts
Local > Topical > Solution: 10 to 25%	Plantar warts

Poliomyelitis vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against poliomyelitis

Polygeline

Section	Indications
Plasma substitutes	Therapeutic equivalent to dextran 70 pour
Parenteral > General injections > IV: 3.5% injectable solution	Hypovolaemia
Parenteral > General injections > IV: 3.5% in solution	Hypovolaemia

Polymyxin B (injection)

Section	Indications
Reserve group antibiotics	Carbapenem resistant Acinetobacter baumannii
Parenteral > General injections > IV: 500000 IU in vial powder for injection (equivalent to 50 mg polymyxin B base)	Carbapenem-resistant Pseudomonas aeruginosa
	Carbapenem resistant Enterobacteriales

Polymyxins

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases
.	

Potassium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Oral Oral > Liquid: powder for solution	Hypokalaemia
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 11.2% in 20 mL ampoule (equivalent to K+ 1.5 mmol/mL and Cl- 1.5 mmol/mL); 7.5% solution for dilution (equivalent to K+ 1 mmol/mL and Cl- 2 mmol/mL); 15% solution for dilution (equivalent to K+ 2 mmol/mL and Cl- 2 mmol/mL)	Hypokalaemia

Potassium ferric hexacyanoferrate (Prussian blue)

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: Powder for oral administration	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium iodide

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 60 mg	Thyrotoxicosis
Antifungal medicines Local > Topical > Solution: Saturated solution	Sporotrichosis

Potassium permanganate

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Solution: aqueous solution: 1:10 000	Tropical phagedaenic ulcer
	Pemphigus
	Other specified pyogenic bacterial infection of skin and subcutaneous tissue
	Impetigo

Povidone iodine

Section	Indications
Antiseptics Local > Topical > Solution: 10% (equivalent to 1% available iodine)	Povidone iodine

Pralidoxime

Section	Indications
Antidotes and other substances used in poisonings > Specific	Exposure to or harmful effects of undetermined intent of pesticides

Pramipexole

Section	Indications
Antiseizure medicines Oral > Solid: 125 µg immediate-release; 250 mg immediate-release; 500 µg immediate-release; 750 µg immediate-release; 1 mg immediate-release; 1.5 mg immediate-release; 375 µg extended-release; 750 µg extended-release; 1.5 mg extended-release; 2.25 mg extended-release; 3 mg extended-release; 3.75 mg extended-release; 4.5 mg extended-release	Parkinson disease

Pravastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia
	Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Praziquantel

Section	Indications
Antischistosomals and other antitrematode medicines Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Paragonimiasis Clonorchiasis Opisthorchiasis Schistosomiasis
Intestinal anthelmintics Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Taeniasis due to Taenia saginata Taeniasis due to Taenia solium Hymenolepasis Diphyllobothriasis
Cysticidal medicines Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg	Cysticercosis of central nervous system

Precipitated sulfur

Section

Dermatological medicines > Scabicides and pediculicides

Local > Topical > Ointment:

Indications

Therapeutic equivalent to [benzyl benzoate](#) pour

[Pthiriasis](#)

Therapeutic equivalent to [benzyl benzoate](#) pour

[Pediculosis corporis](#)

Therapeutic equivalent to [benzyl benzoate](#) pour

[Pediculosis capitis](#)

Therapeutic equivalent to [benzyl benzoate](#) pour

[Scabies](#)

Prednisolone

Section	Indications
Hormones and antihormones Oral > Liquid: 5 mg per mL Oral > Solid: 5 mg; 25 mg	Unspecified malignant neoplasms of ill-defined or unspecified sites Plasma cell myeloma Follicular lymphoma Lymphoid leukaemia, not elsewhere classified Malignant neoplasms of prostate Chronic lymphocytic leukaemia or small lymphocytic lymphoma Hodgkin lymphoma Diffuse large B-cell lymphomas Burkitt lymphoma including Burkitt leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis
Anti-inflammatory medicines Local > Rectal > Retention enema: Local > Rectal > Retention enema: 20 mg per 100 mL (as sodium phosphate)	Therapeutic equivalent to hydrocortisone pour Crohn disease site Crohn disease site Ulcerative colitis
Antiallergics and medicines used in anaphylaxis Oral > Liquid: 5 mg per mL Oral > Solid: 5 mg; 25 mg	Allergic or hypersensitivity conditions of unspecified type
Ophthalmological preparations > Anti-inflammatory agents Local > Ophthalmological > Solution (eye drops): 0.5% (sodium phosphate)	Scleritis Anterior uveitis

Prednisone

Section	Indications
Hormones and antihormones	<p>Therapeutic equivalent to prednisolone pour</p> <p>Plasma cell myeloma</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Follicular lymphoma</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Lymphoid leukaemia, not elsewhere classified</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Malignant neoplasms of prostate</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Chronic lymphocytic leukaemia or small lymphocytic lymphoma</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Hodgkin lymphoma</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Diffuse large B-cell lymphomas</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Anaplastic large cell lymphoma, ALK-negative</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Anaplastic large cell lymphoma, ALK-positive</p> <p>Therapeutic equivalent to prednisolone pour</p> <p>Langerhans cell histiocytosis</p>
Antiallergics and medicines used in anaphylaxis	<p>Therapeutic equivalent to prednisolone pour</p> <p>Allergic or hypersensitivity conditions of unspecified type</p>

Pretomanid

Section	Indications
Antituberculosis medicines Oral > Solid > tablet: 200 mg	<p>Multi-drug resistant Mycobacterium tuberculosis</p>

Primaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)	<p>Malaria due to Plasmodium ovale</p> <p>Malaria due to Plasmodium vivax</p>

Probenecid

Section	Indications
Medicines used to treat gout Oral > Solid: 500 mg	Gout

Procainamide

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride) Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)	Cardiac arrhythmia

Procaine benzylpenicillin

Section	Indications
Access group antibiotics Parenteral > General injections > IM: 1 g in vial (=1 million IU) powder for injection; 3 g in vial (=3 million IU) powder for injection	Premier choix Congenital syphilis [children] Second choix Syphilis

Procarbazine

Section	Indications
Cytotoxic medicines Oral > Solid: 50 mg (as hydrochloride) Oral > Solid > capsule: 50 mg (as hydrochloride)	Unspecified malignant neoplasms of ill-defined or unspecified sites Hodgkin lymphoma

Progesterone vaginal ring

Section	Indications
Intravaginal contraceptives Local > Vaginal > vaginal ring: 2.074 g micronized progesterone	Contact with health services for contraceptive management

Proguanil

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 100 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum

Promethazine

Section	Indications
<p>Antiemetic medicines</p> <p>Oral > Liquid: 5 mg per 5 mL (hydrochloride)</p> <p>Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)</p> <p>Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule</p>	Nausea or vomiting

Propanol

Section	Indications
<p>Antiseptics</p>	Therapeutic equivalent to ethanol pour Denatured alcohol

Propofol

Section	Indications
<p>General anaesthetics and oxygen > Injectable medicines</p> <p>Parenteral > General injections > IV: 10 mg per mL; 20 mg per mL</p>	Anaesthetics and therapeutic gases

Propranolol

Section	Indications
<p>Antihypertensive medicines</p> <p>Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)</p>	Essential hypertension
<p>Antianginal medicines</p> <p>Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)</p>	Angina pectoris
<p>Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)</p>	
<p>Antiarrhythmic medicines</p> <p>Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)</p>	Cardiac arrhythmia
<p>Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)</p>	
<p>Antimigraine medicines > For prophylaxis</p> <p>Oral > Solid: 40 mg tablet (hydrochloride); 20 mg tablet (hydrochloride)</p>	Migraine

Propylthiouracil

Section	Indications
<p>Thyroid hormones and antithyroid medicines</p> <p>Oral > Solid: 50 mg</p>	Thyrotoxicosis

Prostaglandin E1

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 0.5 mg per mL in alcohol	Other specified congenital anomaly of great arteries including arterial duct

Prostaglandin E2

Section	Indications
Medicines administered to the neonate [c] Parenteral > General injections > IV: 1 mg per mL	Therapeutic equivalent to prostaglandin E1 pour Other specified congenital anomaly of great arteries including arterial duct Other specified congenital anomaly of great arteries including arterial duct

Protamine sulfate

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 10 mg per mL in 5 mL ampoule	Haemorrhagic disorder due to other specified circulating anticoagulants

Protonamide

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to ethionamide pour Multi-drug resistant <i>Mycobacterium tuberculosis</i>

Pyrantel

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 250 mg (chewable) (as embonate or pamoate)	Ascariasis Trichuriasis Enterobiasis Hookworm diseases

Pyrazinamide

Section	Indications
Antituberculosis medicines Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 400 mg; 500 mg	Tuberculosis

Pyridostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 1 mg in 1 mL ampoule (pyridostigmine bromide) Oral > Solid: 60 mg (pyridostigmine bromide)	Myasthenia gravis

Pyridoxine

Section	Indications
Vitamins and minerals Oral > Solid: 25 mg (hydrochloride)	Vitamin B6 deficiency Sideroblastic anaemia Other specified polyneuropathy

Pyrimethamine

Section	Indications
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid: 25 mg	Toxoplasmosis

Quetiapine

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders
Medicines for bipolar disorders Oral > Solid > tablet: 25 mg (immediate-release); 100 mg (immediate-release); 150 mg (immediate-release); 200 mg (immediate-release); 300 mg (immediate-release); 50 mg (modified-release); 150 mg (modified-release); 200 mg (modified-release); 300 mg (modified-release); 400 mg (modified-release)	Bipolar or related disorders

Quinidine

Section	Indications
Antiarrhythmic medicines Oral > Solid: 200 mg (sulfate)	Cardiac arrhythmia

Quinine

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride) Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)	Malaria due to <i>Plasmodium falciparum</i>

Rabies vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against rabies

Raltegravir

Section	Indications
Antiretrovirals > Integrase inhibitors Oral > Liquid: 100 mg granules for oral suspension Oral > Solid: 400 mg tablet; 25 mg tablet (chewable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Ranibizumab

Section	Indications
Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations	Retinal venous occlusions Diabetic macular oedema Degenerative high myopia Age related macular degeneration

Ranitidine

Section	Indications
Antiulcer medicines Oral > Liquid: 75 mg per 5 mL (as hydrochloride) Oral > Solid: 150 mg (as hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (as hydrochloride)	Gastro-oesophageal reflux disease Peptic ulcer, site unspecified

Rasburicase

Section	Indications
Supportive medicines Parenteral > General injections > IV: 1.5 mg in vial powder and solvent for solution; 7.5 mg in vial powder and solvent for solution	Tumour lysis syndrome

Ravidasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg	Chronic hepatitis C

Ready to use therapeutic food

Section	Indications
Therapeutic foods Biscuit or paste of nutritional composition as determined by the UN joint statement on the community-based management of severe acute malnutrition and Codex alimentarius guidelines.	Acute malnutrition in infants, children or adolescents

Realgar-indigo naturalis formulation

Section	Indications
Cytotoxic medicines Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)	Acute myeloid leukaemia with recurrent genetic abnormalities

Red blood cells

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Resin-based composite (high-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use capsule or multi-use syringe	Dental caries

Resin-based composite (low-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use applicator or multi-use bottle	Dental caries

Retinol

Section	Indications
<p>Vitamins and minerals</p> <p>Parenteral > General injections > IM: 100000 IU in 2 mL ampoule (as palmitate) water-miscible injection</p> <p>Oral > Liquid: 100000 IU per mL (as palmitate) oral oily solution in multidose dispenser</p> <p>Oral > Solid: 10000 IU (as palmitate) sugar-coated tablet; 200000 IU (as palmitate) capsule; 50000 IU (as palmitate) capsule; 100000 IU (as palmitate) capsule</p>	<p>Vitamin A deficiency</p>

Ribavirin

Section	Indications
<p>Other antivirals</p> <p>Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution</p> <p>Oral > Solid: 200 mg; 400 mg; 600 mg</p>	<p>Viral haemorrhagic fever, not elsewhere classified</p>
<p>Medicines for hepatitis C > Other antivirals for hepatitis C</p> <p>Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution</p> <p>Oral > Solid: 200 mg; 400 mg; 600 mg</p>	<p>Chronic hepatitis C</p>

Ribociclib

Section	Indications
<p>Targeted therapies</p>	<p>Therapeutic equivalent to palbociclib pour Other specified malignant neoplasms of breast</p>

Riboflavin

Section	Indications
<p>Vitamins and minerals</p> <p>Oral > Solid: 5 mg</p>	<p>Vitamin B2 deficiency</p>

Rifabutin

Section	Indications
<p>Antituberculosis medicines</p> <p>Oral > Solid: 150 mg</p>	<p>Tuberculosis</p>

Rifampicin

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Liquid: 20 mg per mL	Other specified tuberculosis
Oral > Solid: 150 mg; 300 mg	
Parenteral > General injections > IV: 600 mg in vial powder for injection	
Antileprosy medicines	Leprosy
Oral > Liquid: 20 mg per mL	
Oral > Solid: 150 mg; 300 mg	

Rifapentine

Section	Indications
Antituberculosis medicines	Tuberculosis
Oral > Solid > tablet: 150 mg; 300 mg	Latent tuberculosis

Risdiplam

Section	Indications
Medicines for diseases of the nervous system	Spinal muscular atrophy
Oral > Liquid: 0.75 mg per mL powder for oral liquid	

Risperidone

Section	Indications
Medicines for psychotic disorders	Schizophrenia or other primary psychotic disorders
Oral > Solid: 0.25 mg; 0.5 mg; 1 mg; 2 mg; 3 mg; 4 mg; 6 mg	
Parenteral > General injections > IM:	Therapeutic equivalent to paliperidone pour Schizophrenia or other primary psychotic disorders

Ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors	
Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Rituximab

Section	Indications
<p>Targeted therapies</p> <p>Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial</p>	<p>Follicular lymphoma</p> <p>Chronic lymphocytic leukaemia or small lymphocytic lymphoma</p> <p>Diffuse large B-cell lymphomas</p> <p>Burkitt lymphoma including Burkitt leukaemia</p>
<p>Medicines for multiple sclerosis</p> <p>Parenteral > General injections > IV: 500 mg per 50 mL in 50 mL vial</p>	Multiple sclerosis

Rivaroxaban

Section	Indications
<p>Medicines affecting coagulation</p>	<p>Therapeutic equivalent to dabigatran pour</p> <p>Venous thromboembolism</p>
	<p>Therapeutic equivalent to dabigatran pour</p> <p>Other specified atrial fibrillation</p>

Ropinirole

Section	Indications
<p>Antiseizure medicines</p> <p>Oral > Solid: 0.25 mg immediate-release; 0.5 mg immediate-release; 1 mg immediate-release; 2 immediate-release; 3 mg immediate-release; 4 mg immediate-release; 5 mg immediate-release; 2 mg extended-release; 4 mg extended-release; 6 mg extended-release; 8 mg extended-release; 12 mg extended-release</p>	Parkinson disease

Rotavirus vaccine

Section	Indications
<p>Recommendations for all immunization programmes</p> <p>All vaccines should comply with the WHO requirements for biological substances.</p>	Need for immunization against certain single viral diseases

Rubella vaccine

Section	Indications
<p>Recommendations for all immunization programmes</p> <p>All vaccines should comply with the WHO requirements for biological substances.</p>	Need for immunization against rubella alone

Salbutamol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Parenteral > General injections > IV: 50 µg per mL in 5 mL ampoule (as sulfate) Respiratory > Inhalation > aerosol: 100 µg per dose (as sulfate); 100 µg per dose (as sulfate) metered-dose inhaler Respiratory > Inhalation > solution: 5 mg per mL (as sulfate) for use in nebulizers	Chronic obstructive pulmonary disease Asthma

Salicylic acid

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation Local > Topical > Solution: 5%	Seborrhoeic dermatitis Psoriasis of unspecified type Ichthyoses

Saquinavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 200 mg; 500 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Selenium sulfide

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Suspension: 2% detergent-based	Seborrhoeic dermatitis Pityriasis versicolor

Senna

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 7.5 mg per 5 mL oral liquid Laxatives Oral > Solid: 7.5 mg (sennosides) (or traditional dosage forms)	Palliative care Constipation

Sevoflurane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines	Anaesthetics and therapeutic gases
Respiratory > Inhalation > liquid:	

Silver diamine fluoride

Section	Indications
Dental medicines and preparations	Dental caries
Local > Dental > Solution: 38% w/v	

Silver nitrate

Section	Indications
Ophthalmological preparations > Anti-infective agents	Neonatal conjunctivitis or dacryocystitis
Local > Ophthalmological > Solution (eye drops): 1%	

Silver sulfadiazine

Section	Indications
Dermatological medicines > Anti-infective medicines	Burns
Local > Topical > Cream: 1%	

Simeprevir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations	Chronic hepatitis C
Oral > Solid: 150 mg	

Simvastatin

Section	Indications
Lipid-lowering agents	Mixed hyperlipidaemia
Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg	Coronary atherosclerosis
Medicines for endocrine disorders	Polycystic ovary syndrome
Oral > Solid: 20 mg	

Smallpox vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Snake antivenom

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium calcium edetate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 0.9% isotonic (equivalent to Na+ 154 mmol/L and Cl- 154 mmol/L)	Other specified disorders of fluid, electrolyte or acid-base balance

Sodium hyaluronate

Section	Indications
Ophthalmological preparations	Therapeutic equivalent to hyromellose pour Keratoconjunctivitis sicca

Sodium hydrogen carbonate

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 1.4% isotonic (equivalent to Na+ 167 mmol/L and HCO3- 167 mmol/L); 8.4% in 10 mL ampoule (equivalent to Na+ 1000 mmol/L and HCO3- 1000 mmol/L)	Acidosis

Sodium nitrite

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitroprusside

Section	Indications
Antihypertensive medicines Parenteral > General injections > IV: 50 mg in ampoule powder for infusion	Hypertensive crisis

Sodium stibogluconate

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial	Visceral leishmaniasis Mucocutaneous leishmaniasis Cutaneous leishmaniasis

Sodium thiosulfate

Section	Indications
Dermatological medicines > Antifungal medicines Local > Topical > Solution: 15%	Pityriasis versicolor
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sofosbuvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg; 400 mg	Chronic hepatitis C

Sofosbuvir + velpatasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg + 50 mg; 400 mg + 100 mg	Chronic hepatitis C

Somatropin

Section	Indications
Medicines for endocrine disorders Parenteral > General injections > SC: 5 mg in cartridge powder for injection; 5 mg per mL in 2 mL cartridge	Hypoglycaemia without associated diabetes

Spectinomycin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 2 g in vial (as hydrochloride) powder for injection	Second choix Gonococcal infection
	Autres indications
	Bacterial infection of unspecified site

Spironolactone

Section	Indications
Medicines used in heart failure Oral > Solid > tablet: 25 mg	Heart failure
Diuretics Oral > Liquid: 5 mg per 5 mL; 10 mg per 5 mL; 25 mg per 5 mL Oral > Solid > tablet: 25 mg	Oedema Ascites

Stavudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 5 mg per 5 mL powder for oral solution Oral > Solid: 15 mg; 20 mg; 30 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Streptokinase

Section	Indications
Thrombolytic medicines Parenteral > General injections > IV: 1.5 million IU in vial powder for injection	Acute myocardial infarction

Streptomycin (injection)

Section	Indications
Antituberculosis medicines Parenteral > General injections > IV: 1 g powder for injection (as sulfate) as vial	Multi-drug resistant Mycobacterium tuberculosis

Succimer

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 100 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sulfacetamide

Section	Indications
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 10% (sodium salt)	Other specified conjunctivitis

Sulfadiazine

Section	Indications
Antipneumocystosis and antitoxoplasmosis medicines Oral > Solid: 500 mg	Toxoplasmosis
Anti-infective medicines Oral > Solid: 500 mg	Bacterial infection of unspecified site
Parenteral > General injections > unspecified: 250 mg in 4 mL ampoule (sodium salt)	

Sulfadimidine

Section	Indications
Anti-infective medicines Oral > Liquid: 500 mg per 5 mL	Bacterial infection of unspecified site
Oral > Solid: 500 mg	
Parenteral > General injections > unspecified: 1 g in 3 mL ampoule (sodium salt)	

Sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For curative treatment	Malaria due to Plasmodium falciparum
Oral > Solid: 500 mg + 25 mg tablet	Malaria due to Plasmodium vivax
Antimalarial medicines > For chemoprevention	Malaria due to Plasmodium falciparum
Oral > Solid: 500 mg + 25 mg tablet	Malaria
Oral > Solid: 250 mg + 12.5 mg tablet	

Sulfamethoxazole + trimethoprim

Section	Indications
Access group antibiotics	Premier choix
Parenteral > General injections > IV: 80 mg + 16 mg per mL in 10 mL ampoule; 80 mg + 16 mg per mL in 5 mL ampoule	Infectious cystitis
Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid	Second choix
Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet	Infectious gastroenteritis or colitis without specification of infectious agent
Oral > Solid > dispersible tablet: 100 mg + 20 mg	Autres indications
Antipneumocystosis and antitoxoplasmosis medicines	Bacterial infection of unspecified site
Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule	Pneumocystosis
Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid	
Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet	
Oral > Solid > dispersible tablet: 100 mg + 20 mg	

Sulfasalazine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs)	Rheumatoid arthritis, serology unspecified
Oral > Solid: 500 mg	
Anti-inflammatory medicines	Crohn disease site
Oral > Solid: 500 mg	Ulcerative colitis
Local > Rectal > Suppository: 500 mg	
Local > Rectal > Retention enema: 3 g per 100 mL	

Sumatriptan

Section	Indications
Antimigraine medicines > For treatment of acute attack Oral > Solid: 50 mg	Migraine

Sunscreen (broad-spectrum)

Section	Indications
Dermatological medicines Topical cream, lotion, gel, ointment, spray with sun protection factor \geq 50.	Albinism or other specified genetically-determined hypomelanotic disorders Xeroderma pigmentosum

Suramin sodium

Section	Indications
Antifilarials Parenteral > General injections > IV: 1 g in vial	Filariasis
Medicines for the treatment of 1st stage African trypanosomiasis Parenteral > General injections > IV: 1 g in vial	African trypanosomiasis

Surfactant

Section	Indications
Medicines administered to the neonate [c] Respiratory > Suspension: 80 mg per mL for intratracheal instillation; 25 mg per mL for intratracheal instillation	Respiratory distress syndrome of newborn

Suxamethonium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 50 mg per mL in 2 mL ampoule (suxamethonium chloride); powder for injection (suxamethonium chloride) in vial	Muscle relaxants

Tacalcitol

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Therapeutic equivalent to calcipotriol pour Psoriasis of unspecified type

Tacrolimus

Section	Indications
<p>Immunomodulators for non-malignant disease Parenteral > General injections > IV: 5 mg per mL in 1 mL vial Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)</p>	<p>Failure or rejection of transplanted organs or tissues</p>

Tamoxifen

Section	Indications
<p>Hormones and antihormones Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)</p>	<p>Other specified malignant neoplasms of breast Malignant neoplasms of breast</p>

Tedizolid

Section	Indications
Reserve group antibiotics	<p>Therapeutic equivalent to linezolid pour Methicillin resistant Staphylococcus aureus</p>
	<p>Therapeutic equivalent to linezolid pour Vancomycin resistant Staphylococcus aureus</p>
	<p>Therapeutic equivalent to linezolid pour Vancomycin resistant Enterococcus</p>

Telmisartan + amlodipine

Section	Indications
<p>Antihypertensive medicines Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg</p>	<p>Essential hypertension</p>

Telmisartan + hydrochlorothiazide

Section	Indications
<p>Antihypertensive medicines Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg</p>	<p>Essential hypertension</p>

Tenofovir alafenamide

Section	Indications
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors	Chronic hepatitis B
Oral > Solid: 25 mg	

Tenofovir disoproxil fumarate

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified
Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)	Contact with or exposure to human immunodeficiency virus
Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors	Chronic hepatitis B
Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)	

Terbinafine

Section	Indications
Dermatological medicines > Antifungal medicines	Fungal infection of the skin
Local > Topical > Cream: 1% terbinafine hydrochloride	
Local > Topical > Ointment: 1% terbinafine hydrochloride	

Terbutaline

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to salbutamol pour
	Chronic obstructive pulmonary disease
	Therapeutic equivalent to salbutamol pour
	Asthma

Terizidone

Section	Indications
Antituberculosis medicines	Therapeutic equivalent to cycloserine pour
	Multi-drug resistant Mycobacterium tuberculosis

Testosterone

Section	Indications
Androgens Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule (enantate)	Testicular dysfunction or testosterone-related disorders

Tetanus antitoxin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IM: 50000 IU in vial	Tetanus

Tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against tetanus alone

Tetracaine

Section	Indications
Ophthalmological preparations > Local anaesthetics Local > Ophthalmological > Solution (eye drops): 0.5% (hydrochloride)	Local anaesthetics

Tetracycline

Section	Indications
Access group antibiotics Oral > Solid: 250 mg (hydrochloride)	Bacterial infection of unspecified site
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)	Neonatal conjunctivitis or dacryocystitis
	Other specified conjunctivitis
	Infectious keratitis
	Infectious blepharitis
	Trachoma

Thalidomide

Section	Indications
Immunomodulators Oral > Solid: 50 mg	Plasma cell myeloma

Thiamine

Section	Indications
Vitamins and minerals Oral > Solid: 50 mg (hydrochloride)	Vitamin B1 deficiency

Thiopental

Section	Indications
General anaesthetics and oxygen > Injectable medicines	Therapeutic equivalent to propofol pour Anaesthetics and therapeutic gases

Ticagrelor

Section	Indications
Anti-platelet medicines Oral > Solid > tablet: 60 mg; 90 mg	Acute ischaemic heart disease Presence of coronary angioplasty implant or graft

Tick-borne encephalitis vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against arthropod-borne viral encephalitis

Tigecycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg in vial powder for injection	Other specified bacterial diseases

Timolol

Section	Indications
Ophthalmological preparations > Miotics and antiglaucoma medicines Local > Ophthalmological > Solution (eye drops): 0.25% (hydrogen maleate); 0.5% (hydrogen maleate)	Primary open-angle glaucoma Ocular hypertension

Tinidazole

Section	Indications
Antiamoebic and antigiardiasis medicines	Therapeutic equivalent to metronidazole pour Amoebiasis

Tioguanine

Section	Indications
Cytotoxic medicines Oral > Solid: 40 mg	Lymphoid leukaemia, not elsewhere classified

Tiotropium bromide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > dry powder: 18 µg capsule Respiratory > Inhalation > solution: 1.25 µg per actuation; 2.5 µg per actuation	Chronic obstructive pulmonary disease

Tisagenlecleucel

Section	Indications
Antineoplastics and supportive medicines	Therapeutic equivalent to axicabtagene ciloleucel pour Diffuse large B-cell lymphomas

Tislelizumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 100 mg per 10 mL	Malignant neoplasms of urinary tract Hodgkin lymphoma Other specified malignant neoplasms of bronchus or lung

Tobramycin

Section	Indications
Ophthalmological preparations > Anti-infective agents	Therapeutic equivalent to gentamicin pour Other specified conjunctivitis Therapeutic equivalent to gentamicin pour Infectious blepharitis

Tocilizumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	Juvenile systemic arthritis

Tolbutamide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 500 mg	Type 2 diabetes mellitus

Torasemide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure
Diuretics	Therapeutic equivalent to furosemide pour Anuria or oliguria
	Therapeutic equivalent to furosemide pour Oedema

Toripalimab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 240 mg per 6 mL in vial	Malignant neoplasms of nasopharynx Malignant neoplasms of oesophagus

Tramadol

Section	Indications
<p>Opioid analgesics</p> <p>Oral > Liquid: 100 mg per mL (hydrochloride)</p> <p>Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release</p> <p>Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)</p>	Chronic cancer pain

Trametinib

Section	Indications
<p>Targeted therapies</p> <p>Oral > Solid: 0.5 mg; 2 mg</p>	Other specified melanoma of skin

Tranexamic acid

Section	Indications
<p>Medicines affecting coagulation</p> <p>Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule</p>	Haemorrhage, not elsewhere classified
<p>Other medicines administered to the mother</p> <p>Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule</p>	Postpartum haemorrhage

Trastuzumab

Section	Indications
<p>Targeted therapies</p> <p>Parenteral > General injections > IV: 60 mg in vial powder for injection; 150 mg in vial powder for injection; 440 mg in vial powder for injection</p>	Carcinoma of breast, specialised type

Trastuzumab emtansine

Section	Indications
<p>Targeted therapies</p> <p>Parenteral > General injections > IV: 100 mg in vial powder for injection; 160 mg in vial powder for injection</p>	Carcinoma of breast, specialised type

Triamcinolone hexacetonide

Section

Medicines for juvenile joint diseases

Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

Indications

Juvenile idiopathic arthritis

Triclabendazole

Section

Antischistosomals and other antitrematode medicines

Oral > Solid > tablet: 250 mg (scored)

Indications

Paragonimiasis

Fascioliasis

Trihexyphenidyl

Section

Antiseizure medicines

Oral > Solid: 2 mg (hydrochloride); 5 mg (hydrochloride)

Medicines for parkinsonism

Indications

Parkinson disease

Therapeutic equivalent to biperiden pour

Parkinson disease

Trimethoprim

Section

Access group antibiotics

Oral > Liquid: 50 mg per mL

Oral > Solid: 100 mg; 200 mg

Indications

Premier choix

Infectious cystitis

Triptorelin

Section

Hormones and antihormones

Indications

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of breast

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of prostate

Tropicamide

Section

Diagnostic agents > Ophthalmic medicines

Local > Ophthalmological > Solution (eye drops): 0.5%

Indications

Cycloplegic drug

Tropisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Tuberculin, purified protein derivative

Section	Indications
Immunologicals > Diagnostic agents Parenteral > Locoregional injections > Intradermal:	Mycobacterial diseases

Tubocurarine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg per mL in 1.5 mL ampoule (tubocurarine chloride)	Muscle relaxants

Typhoid vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against typhoid-paratyphoid alone

Ulipristal

Section	Indications
Oral hormonal contraceptives Oral > Solid: 30 mg tablet (ulipristal acetate)	Contact with health services for postcoital contraception

Umeclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Urea

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Xerosis cutis or asteatosis
Local > Topical > Cream: 10%; 5%	
Local > Topical > Ointment: 10%; 5%	

Ustekinumab

Section	Indications
Dermatological medicines > Medicines affecting skin differentiation and proliferation	Psoriasis of unspecified type
Parenteral > General injections > SC: 45 mg per 0.5 mL in vial; 90 mg per mL in pre-filled syringe	

Valaciclovir

Section	Indications
Antiherpes medicines	Therapeutic equivalent to aciclovir pour Zoster
	Therapeutic equivalent to aciclovir pour Varicella
	Therapeutic equivalent to aciclovir pour Herpes simplex infections

Valganciclovir

Section	Indications
Other antivirals	Cytomegaloviral retinitis
Oral > Liquid: 50 mg per mL powder for oral solution	
Oral > Solid > tablet: 450 mg	

Valproic acid (sodium valproate)

Section	Indications
Antiseizure medicines	Epilepsy or seizures
Oral > Liquid: 200 mg per 5 mL	Status epilepticus
Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated); 100 mg tablet (crushable)	
Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule; 100 mg per mL in 3 mL ampoule	
Medicines for bipolar disorders	Bipolar or related disorders
Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated)	

Vancomycin

Section	Indications
Watch group antibiotics	Premier choix
Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection	Necrotising fasciitis co-prescrite avec ceftazidime
Oral > Solid > capsule: 125 mg (as hydrochloride); 250 mg (as hydrochloride)	Endophthalmitis co-prescrite avec ceftriaxone
Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection	Endophthalmitis
Parenteral > Locoregional injections > Intravitreal: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection	Second choix Neutropenia (high-risk) Intestinal infections due to Clostridioides difficile
	Autres indications Methicillin resistant Staphylococcus aureus

Varenicline

Section	Indications
Medicines for nicotine use disorders	Nicotine dependence
Oral > Solid: 0.5 mg; 1 mg	

Varicella vaccine

Section	Indications
Recommendations for immunization programmes with certain characteristics	Need for immunization against certain single viral diseases
All vaccines should comply with the WHO requirements for biological substances.	

Vecuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg in vial powder for injection (vecuronium bromide)	Muscle relaxants

Vemurafenib

Section	Indications
Targeted therapies	Therapeutic equivalent to dabrafenib pour Other specified melanoma of skin

Verapamil

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride) Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)	Supraventricular tachyarrhythmia
Antianginal medicines Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)	Angina pectoris

Vinblastine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate) Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Germ cell tumour of testis Anaplastic large cell lymphoma, ALK-positive Anaplastic large cell lymphoma, ALK-negative Langerhans cell histiocytosis

Vincristine

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate)</p> <p>Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)</p>	<p>Unspecified malignant neoplasms of ill-defined or unspecified sites</p> <p>Other specified gliomas of brain</p> <p>Malignant trophoblastic neoplasms of placenta</p> <p>Follicular lymphoma</p> <p>Kaposi sarcoma of unspecified primary site</p> <p>Rhabdomyosarcoma primary site</p> <p>Hodgkin lymphoma</p> <p>Diffuse large B-cell lymphomas</p> <p>Retinoblastoma</p> <p>Burkitt lymphoma including Burkitt leukaemia</p> <p>Malignant neoplasms of kidney, except renal pelvis</p> <p>Ewing sarcoma of bone and articular cartilage of unspecified sites</p> <p>Lymphoid leukaemia, not elsewhere classified</p> <p>Langerhans cell histiocytosis</p>

Vinorelbine

Section	Indications
<p>Cytotoxic medicines</p> <p>Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial</p> <p>Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial</p> <p>Oral > Solid > capsule: 20 mg; 30 mg; 80 mg</p>	<p>Other specified malignant neoplasms of bronchus or lung</p> <p>Other specified malignant neoplasms of breast</p> <p>Rhabdomyosarcoma primary site</p>

Voriconazole

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 200 mg in vial powder for injection</p> <p>Oral > Liquid: 40 mg per mL powder for oral liquid</p> <p>Oral > Solid: 50 mg tablet; 200 mg tablet</p>	<p>Chronic pulmonary aspergillosis</p> <p>Invasive aspergillosis</p>

Warfarin

Section	Indications
Medicines affecting coagulation Oral > Solid: 5 mg (sodium salt); 1 mg (sodium salt); 2 mg (sodium salt); 0.5 mg (sodium salt)	Other specified diseases of arteries or arterioles Cerebral ischaemic stroke due to embolic occlusion Venous thromboembolism Atrial fibrillation

Water for injection

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Miscellaneous Parenteral > General injections > unspecified: 2 mL ampoule; 5 mL ampoule; 10 mL ampoule	Unspecified appropriateness of dosing or administration without injury or harm

Whole blood

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Xylometazoline

Section	Indications
Ear, nose and throat medicines [c] Local > Nasal > Spray: 0.05% w/v equivalent to 0.5 mg per mL	Nasal congestion

Yellow fever vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Zanubrutinib

Section	Indications
Targeted therapies Oral > Solid: 80 mg	Mantle cell lymphoma Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Zidovudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Parenteral > General injections > IV: 10 mg per mL in 20 mL vial solution for IV infusion Oral > Liquid: 50 mg per 5 mL Oral > Solid > tablet: 300 mg Oral > Solid > capsule: 250 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Zinc sulfate

Section	Indications
Medicines for diarrhoea Oral > Solid: 20 mg	Diarrhoea

Zoledronic acid

Section	Indications
Supportive medicines Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle	Malignant neoplasm metastasis in bone or bone marrow
Medicines for endocrine disorders Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle	Osteogenesis imperfecta

Zuclopentixol decanoate

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to fluphenazine pour Schizophrenia or other primary psychotic disorders