

Chloroquine

Section	Indications
<p>Antimalarial medicines > For chemoprevention</p> <p>Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)</p> <p>Oral > Solid: 150 mg tablet (as phosphate or sulfate)</p>	<p>Malaria due to <i>Plasmodium falciparum</i></p> <p>Malaria due to <i>Plasmodium vivax</i></p> <p>Malaria due to <i>Plasmodium ovale</i></p> <p>Malaria due to <i>Plasmodium malariae</i></p>
<p>Antimalarial medicines > For curative treatment</p> <p>Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)</p> <p>Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)</p> <p>Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)</p>	<p>Malaria due to <i>Plasmodium falciparum</i></p> <p>Malaria due to <i>Plasmodium vivax</i></p>
<p>Disease-modifying anti-rheumatic drugs (DMARDs)</p> <p>Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)</p>	<p>Rheumatoid arthritis</p>