

**Acetylsalicylic acid**

## Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral &gt; Solid: 100 to 500 mg

Local &gt; Rectal &gt; Suppository: 50 to 150 mg

## Indications

Pain

**Amitriptyline**

## Section

Medicines for other common symptoms in palliative care

Oral &gt; Solid: 25 mg; 10 mg; 75 mg

## Indications

Palliative care

**Codeine**

## Section

Opioid analgesics

Oral &gt; Solid: 30 mg tablet (codeine phosphate)

## Indications

Pain

**Cyclizine**

## Section

Medicines for other common symptoms in palliative care

Oral &gt; Solid: 50 mg tablet

Parenteral &gt; General injections &gt; unspecified: 50 mg per mL

## Indications

Palliative care

**Dexamethasone**

## Section

Medicines for other common symptoms in palliative care

Oral &gt; Liquid: 2 mg per 5 mL

Oral &gt; Solid: 2 mg; 4 mg

Parenteral &gt; General injections &gt; unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

## Indications

Palliative care

## Diazepam

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 5 mg; 10 mg

Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system

Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube

Parenteral > General injections > unspecified: 5 mg per mL

### Indications

Palliative care

## Docusate sodium

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 50 mg per 5 mL oral liquid

Oral > Solid: 100 mg capsule

### Indications

Palliative care

## Dolasetron

### Section

Medicines for other common symptoms in palliative care

### Indications

Therapeutic equivalent to ondansetron for

Palliative care

## Fentanyl

### Section

Opioid analgesics

Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour

Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate)

Oral > Other: 200 to 1200 µg buccal film (as citrate)

### Indications

Chronic cancer pain

Other specified chronic cancer related pain

## Fluoxetine

### Section

Medicines for other common symptoms in palliative care

Oral > Solid: 20 mg (as hydrochloride)

### Indications

Palliative care

## Gabapentin

### Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg

### Indications

Neuropathic pain

## Granisetron

### Section

Medicines for other common symptoms in palliative care

### Indications

Therapeutic equivalent to **ondansetron** for

Palliative care

## Haloperidol

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per mL

Oral > Solid: 0.5 mg; 2 mg; 5 mg

Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule

### Indications

Palliative care

## Hydromorphone

### Section

Opioid analgesics

### Indications

Therapeutic equivalent to **morphine** for

Pain

## Hyoscine butylbromide

### Section

Medicines for other common symptoms in palliative care

Parenteral > General injections > unspecified: 20 mg per mL injection

### Indications

Palliative care

## Hyoscine hydrobromide

### Section

Medicines for other common symptoms in palliative care

Local > Topical > Transdermal patch: 1 mg per 72 hours transdermal patch

Parenteral > General injections > unspecified: 400 µg per mL injection; 600 µg per mL injection

### Indications

Palliative care

## Ibuprofen

### Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL

Oral > Solid > tablet: 200 mg; 400 mg; 600 mg

### Indications

Pain

## Lactulose

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 3.1 to 3.7 g per 5 mL oral liquid

### Indications

Palliative care

## Loperamide

### Section

Medicines for other common symptoms in palliative care

Oral > Solid > dosage form: 2 mg

### Indications

Palliative care

## Methadone

### Section

Opioid analgesics

Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)

### Indications

Chronic cancer pain

## Metoclopramide

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 5 mg per 5 mL

Oral > Solid: 10 mg (hydrochloride)

Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)

### Indications

Palliative care

## Midazolam

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per mL

Parenteral > General injections > unspecified: 1 mg per mL; 5 mg per mL

Oral > Solid > dosage form: 7.5 mg; 15 mg

### Indications

Palliative care

## Morphine

### Section

Opioid analgesics

Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate)

Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate)

Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate)

Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)

### Indications

Pain

## Ondansetron

### Section

Medicines for other common symptoms in palliative care

Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)

Oral > Liquid: 4 mg per 5 mL (ondansetron base)

Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base)

### Indications

Palliative care

## Oxycodone

### Section

Opioid analgesics

### Indications

Therapeutic equivalent to [morphine](#) for

Pain

## Palonosetron

### Section

Medicines for other common symptoms in palliative care

### Indications

Therapeutic equivalent to [ondansetron](#) for

Palliative care

## Paracetamol (acetaminophen)

### Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs)

Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL

Local > Rectal > Suppository: 100 mg; 250 mg

Oral > Solid > dispersible tablet: 100 mg; 250 mg

Oral > Solid > tablet: 250 mg; 325 mg; 500 mg

### Indications

Pain

## Senna

### Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 7.5 mg per 5 mL oral liquid

### Indications

Palliative care

## Tramadol

### Section

Opioid analgesics

Oral > Liquid: 100 mg per mL (hydrochloride)

Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release

Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)

### Indications

Chronic cancer pain

## Tropisetron

### Section

Medicines for other common symptoms in palliative care

### Indications

Therapeutic equivalent to [ondansetron](#) for

Palliative care