

Acetylcysteine

| Section | Indications |
|---|--|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule Oral > Liquid: 10%; 20% | Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs Acute or subacute hepatic failure |

Activated charcoal

| Section | Indications |
|--|--|
| Antidotes and other substances used in poisonings > Non-specific Oral > Other: powder | Unspecified injury, poisoning or certain other consequences of external causes |

Atropine

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate) | Exposure to or harmful effects of undetermined intent of pesticides |

Calcium gluconate

| Section | Indications |
|--|------------------|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule | Hypermagnesaemia |

Deferoxamine

| Section | Indications |
|---|--|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate) | Harmful effects of drugs, medicaments or biological substances, not elsewhere classified |

Dimercaprol

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Fomepizole

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base) | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Ipecacuanha

| Section | Indications |
|--|--|
| Antidotes and other substances used in poisonings > Non-specific Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine | Unspecified injury, poisoning or certain other consequences of external causes |

Methionine

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg (DL) | Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs |

Methylthioninium chloride

| Section | Indications |
|---|-----------------------------|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule | Acquired methaemoglobinæmia |

Naloxone

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride) | Unintentional exposure to or harmful effects of opioids or related analgesics |

Penicillamine

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Potassium ferric hexacyanoferrate (Prussian blue)

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Oral > Solid: Powder for oral administration | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Pralidoxime

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific | Exposure to or harmful effects of undetermined intent of pesticides |

Sodium calcium edetate

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Sodium nitrite

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Sodium thiosulfate

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

Succimer

| Section | Indications |
|--|---|
| Antidotes and other substances used in poisonings > Specific Oral > Solid: 100 mg | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |