

Albendazole

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 400 mg (chewable, scored)	<input type="text" value="Echinococcosis"/> <input type="text" value="Enterobiasis"/> <input type="text" value="Trichuriasis"/> <input type="text" value="Hookworm diseases"/> <input type="text" value="Cysticercosis"/> <input type="text" value="Ascariasis"/>
Cysticidal medicines Oral > Solid > tablet: 400 mg (chewable, scored); 200 mg (chewable)	<input type="text" value="Other specified echinococcosis"/> <input type="text" value="Cysticercosis of central nervous system"/>
Antifilarials Oral > Solid > tablet: 400 mg (chewable, scored)	<input type="text" value="Filariasis"/>

Diethylcarbamazine

Section	Indications
Antifilarials Oral > Solid: 50 mg tablet (dihydrogen citrate); 100 mg tablet (dihydrogen citrate)	<input type="text" value="Lymphatic filariasis"/>

Ivermectin

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 3 mg	<input type="text" value="Strongyloidiasis"/> <input type="text" value="Ascariasis"/> <input type="text" value="Trichuriasis"/> <input type="text" value="Hookworm diseases"/> <input type="text" value="Ancylostomiasis"/>
Antifilarials Oral > Solid > tablet: 3 mg	<input type="text" value="Onchocerciasis"/> <input type="text" value="Lymphatic filariasis"/>

Levamisole

Section

Intestinal anthelmintics

Oral > Solid > tablet: 50 mg (as hydrochloride); 150 mg (as hydrochloride)

Indications

Ascariasis

Mebendazole

Section

Intestinal anthelmintics

Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)

Cysticidal medicines

Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)

Indications

Ascariasis

Enterobiasis

Trichuriasis

Hookworm diseases

Other specified echinococcosis

Niclosamide

Section

Intestinal anthelmintics

Oral > Solid: 500 mg tablet (chewable)

Indications

Hymenolepiasis

Diphyllobothriasis

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Oxamniquine

Section

Antischistosomes and other antitrematode medicines

Oral > Liquid: 250 mg per 5 mL

Oral > Solid > capsule: 250 mg

Indications

Schistosomiasis due to *Schistosoma mansoni*

Praziquantel

Section

Antischistosomes and other antitrematode medicines

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Intestinal anthelmintics

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Cysticidal medicines

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Indications

Paragonimiasis

Clonorchiasis

Opisthorchiasis

Schistosomiasis

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Hymenolepiasis

Diphyllobothriasis

Cysticercosis of central nervous system

Pyrantel

Section

Intestinal anthelmintics

Oral > Solid > tablet: 250 mg (chewable) (as embonate or pamoate)

Indications

Ascariasis

Trichuriasis

Enterobiasis

Hookworm diseases

Suramin sodium

Section

Antifilarials

Parenteral > General injections > IV: 1 g in vial

Indications

Filariasis

Triclabendazole

Section

Antischistosomes and other antitrematode medicines

Oral > Solid > tablet: 250 mg (scored)

Indications

Paragonimiasis

Fascioliasis