

## Anti-D immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IV: 250 µg in single-dose vial Parenteral > General injections > IM: 250 µg in single-dose vial	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Maternal care for red cell antibodies</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rh incompatibility reaction</div>

## Anti-rabies immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 150 IU per mL in vial Parenteral > Locoregional injections > Other: 150 IU per mL in vial	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rabies</div>

## Anti-tetanus immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IM: 500 IU in vial	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Tetanus</div>

## C1 esterase inhibitor

Section	Indications
Plasma-derived medicines Parenteral > General injections > IV: 500 IU in vial powder for injection	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; background-color: #f0f0f0;">Hereditary angioedema</div>

## Coagulation factor IX

Section	Indications
Blood coagulation factors Parenteral > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection	<div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Coagulation defects, purpura or other haemorrhagic or related conditions</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Haemophilia B</div>

## Coagulation factor IX complex

Section	Indications
Blood coagulation factors	Therapeutic equivalent to coagulation factor IX for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Coagulation defects, purpura or other haemorrhagic or related conditions</div> Therapeutic equivalent to coagulation factor IX for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Haemophilia B</div>

## Coagulation factor VIII

### Section

Blood coagulation factors

Parenteral > General injections > IV: 250 IU in vial powder for injection;  
500 IU in vial powder for injection; 1000 IU in vial powder for injection

### Indications

Haemophilia A

## Cryoprecipitate (not pathogen-reduced)

### Section

Blood and blood components

### Indications

Therapeutic equivalent to cryoprecipitate (pathogen-reduced) for

Haemophilia A

Therapeutic equivalent to cryoprecipitate (pathogen-reduced) for

Coagulation defects, purpura or other haemorrhagic or related conditions

Therapeutic equivalent to cryoprecipitate (pathogen-reduced) for

Von Willebrand disease

Therapeutic equivalent to cryoprecipitate (pathogen-reduced) for

Obstetric haemorrhage

## Cryoprecipitate (pathogen-reduced)

### Section

Blood and blood components

Injection: frozen liquid in bag or lyophilized powder in vial containing:

> 50 IU Factor VIII

> 100 IU vWF

> 140 mg clottable fibrinogen per unit

### Indications

Haemophilia A

Coagulation defects, purpura or other haemorrhagic or related conditions

Von Willebrand disease

Obstetric haemorrhage

## Dextran 40

### Section

Plasma substitutes

.

### Indications

Hypovolaemia

## Dextran 70

### Section

Plasma substitutes

Parenteral > General injections > IV: 6% injectable solution

### Indications

Hypovolaemia

## Fresh-frozen plasma

### Section

Blood and blood components

Parenteral > General injections > IV:

### Indications

Blood transfusion without reported diagnosis

## Normal immunoglobulin

### Section

Human immunoglobulins

Parenteral > General injections > IV: 5% protein solution; 10% protein solution

Parenteral > General injections > IM: 16% protein solution

Parenteral > General injections > SC: 15% protein solution; 16% protein solution

### Indications

Primary immunodeficiencies

Mucocutaneous lymph node syndrome

Langerhans cell histiocytosis

## Platelets

### Section

Blood and blood components

Parenteral > General injections > IV:

### Indications

Blood transfusion without reported diagnosis

## Polygeline

### Section

Plasma substitutes

Parenteral > General injections > IV: 3.5% injectable solution

Parenteral > General injections > IV: 3.5% in solution

### Indications

Therapeutic equivalent to dextran 70 for

Hypovolaemia

Hypovolaemia

## Red blood cells

### Section

Blood and blood components

Parenteral > General injections > IV:

### Indications

Blood transfusion without reported diagnosis

## Whole blood

### Section

Blood and blood components

Parenteral > General injections > IV:

### Indications

Blood transfusion without reported diagnosis