

Anti-D immunoglobulin

Section

Human immunoglobulins
 Parenteral > General injections > IV: 250 µg in single-dose vial
 Parenteral > General injections > IM: 250 µg in single-dose vial

Indications

Maternal care for red cell antibodies

Rh incompatibility reaction

Anti-rabies immunoglobulin

Section

Human immunoglobulins
 Parenteral > General injections > IM: 150 IU per mL in vial
 Parenteral > Locoregional injections > Other: 150 IU per mL in vial

Indications

Rabies

Anti-tetanus immunoglobulin

Section

Human immunoglobulins
 Parenteral > General injections > IM: 500 IU in vial

Indications

Tetanus

C1 esterase inhibitor

Section

Plasma-derived medicines
 Parenteral > General injections > IV: 500 IU in vial powder for injection

Indications

Hereditary angioedema

Coagulation factor IX

Section

Blood coagulation factors
 Parenteral > General injections > IV: 500 IU in vial powder for injection;
 1000 IU in vial powder for injection

Indications

Coagulation defects, purpura or other haemorrhagic or related conditions

Haemophilia B

Coagulation factor IX complex

Section

Blood coagulation factors

Indications

Therapeutic equivalent to coagulation factor IX pour

Coagulation defects, purpura or other haemorrhagic or related conditions

Therapeutic equivalent to coagulation factor IX pour

Haemophilia B

Coagulation factor VIII

Section

Blood coagulation factors

Parenteral > General injections > IV: 250 IU in vial powder for injection; 500 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

Haemophilia A

Normal immunoglobulin

Section

Human immunoglobulins

Parenteral > General injections > IV: 5% protein solution; 10% protein solution

Parenteral > General injections > IM: 16% protein solution

Parenteral > General injections > SC: 15% protein solution; 16% protein solution

Indications

Primary immunodeficiencies

Mucocutaneous lymph node syndrome

Langerhans cell histiocytosis