

Anti-D immunoglobulin

| Section | Indications |
|---|---|
| Human immunoglobulins Parenteral > General injections > IV: 250 µg in single-dose vial Parenteral > General injections > IM: 250 µg in single-dose vial | <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Maternal care for red cell antibodies</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rh incompatibility reaction</div> |

Anti-rabies immunoglobulin

| Section | Indications |
|---|--|
| Human immunoglobulins Parenteral > General injections > IM: 150 IU per mL in vial Parenteral > Locoregional injections > Other: 150 IU per mL in vial | <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Rabies</div> |

Anti-tetanus immunoglobulin

| Section | Indications |
|---|---|
| Human immunoglobulins Parenteral > General injections > IM: 500 IU in vial | <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Tetanus</div> |

C1 esterase inhibitor

| Section | Indications |
|---|--|
| Plasma-derived medicines Parenteral > General injections > IV: 500 IU in vial powder for injection | <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; background-color: #f0f0f0;">Hereditary angioedema</div> |

Coagulation factor IX

| Section | Indications |
|---|--|
| Blood coagulation factors Parenteral > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection | <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Coagulation defects, purpura or other haemorrhagic or related conditions</div> <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Haemophilia B</div> |

Coagulation factor IX complex

| Section | Indications |
|---------------------------|--|
| Blood coagulation factors | Therapeutic equivalent to coagulation factor IX for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px; margin-bottom: 5px;">Coagulation defects, purpura or other haemorrhagic or related conditions</div> Therapeutic equivalent to coagulation factor IX for <div style="border: 1px solid #0070C0; border-radius: 5px; padding: 2px;">Haemophilia B</div> |

Coagulation factor VIII

Section

Blood coagulation factors

Parenteral > General injections > IV: 250 IU in vial powder for injection;
500 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

Haemophilia A

Normal immunoglobulin

Section

Human immunoglobulins

Parenteral > General injections > IV: 5% protein solution; 10% protein
solution

Parenteral > General injections > IM: 16% protein solution

Parenteral > General injections > SC: 15% protein solution; 16% protein
solution

Indications

Primary immunodeficiencies

Mucocutaneous lymph node syndrome

Langerhans cell histiocytosis