

## Amiodarone

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride)

Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)

### Indications

Cardiac arrhythmia

## Bisoprolol

### Section

Antiarrhythmic medicines

Oral > Solid: 1.25 mg; 5 mg

### Indications

Cardiac arrhythmia

## Carvedilol

### Section

Antiarrhythmic medicines

### Indications

Therapeutic equivalent to bisoprolol for

Cardiac arrhythmia

## Digoxin

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid: 62.5 µg; 250 µg

### Indications

Cardiac arrhythmia

## Epinephrine

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)

### Indications

Cardiac arrest

## Isoprenaline

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)

Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)

### Indications

Bradycardia

Cardiogenic shock

## Lidocaine

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)

### Indications

Ventricular tachyarrhythmia

## Metoprolol

### Section

#### Antiarrhythmic medicines

### Indications

Therapeutic equivalent to bisoprolol for

Cardiac arrhythmia

## Procainamide

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)

Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)

### Indications

Cardiac arrhythmia

## Propranolol

### Section

#### Antiarrhythmic medicines

Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)

Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

### Indications

Cardiac arrhythmia

## Quinidine

### Section

#### Antiarrhythmic medicines

Oral > Solid: 200 mg (sulfate)

### Indications

Cardiac arrhythmia

## Verapamil

### Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

### Indications

Supraventricular tachyarrhythmia