

**Amidotrizoate**

## Section

Diagnostic agents > Radiocontrast media  
Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)

## Indications

Amidotrizoate

**Atropine**

## Section

Diagnostic agents > Ophthalmic medicines

## Indications

Therapeutic equivalent to tropicamide for

Cycloplegic drug

**Barium sulfate**

## Section

Diagnostic agents > Radiocontrast media  
Oral > Liquid: Aqueous suspension

## Indications

Barium sulfate with suspending agents

**Cyclopentolate**

## Section

Diagnostic agents > Ophthalmic medicines

## Indications

Therapeutic equivalent to tropicamide for

Cycloplegic drug

**Fluorescein**

## Section

Diagnostic agents > Ophthalmic medicines  
Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)

## Indications

Fluorescein

**Iohexol**

## Section

Diagnostic agents > Radiocontrast media  
Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule

## Indications

X-ray contrast media, iodinated

## Meglumine iotroxate

### Section

Diagnostic agents > Radiocontrast media

Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL

### Indications

X-ray contrast media, iodinated

## Tropicamide

### Section

Diagnostic agents > Ophthalmic medicines

Local > Ophthalmological > Solution (eye drops): 0.5%

### Indications

Cycloplegic drug