

Aprepitant

Section	Indications
Antiemetic medicines Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid: 80 mg; 125 mg; 165 mg	<input type="text" value="Nausea or vomiting"/>

Bisacodyl

Section	Indications
Laxatives	Therapeutic equivalent to <i>senna</i> for <input type="text" value="Constipation"/>

Cimetidine

Section	Indications
Antiulcer medicines Oral > Solid: 200 mg Parenteral > General injections > unspecified: 200 mg in 2 mL ampoule	<input type="text" value="Peptic ulcer, site unspecified"/> <input type="text" value="Gastro-oesophageal reflux disease"/>

Codeine

Section	Indications
Medicines for diarrhoea Oral > Solid: 30 mg tablet (codeine phosphate)	<input type="text" value="Diarrhoea"/>

Dexamethasone

Section	Indications
Antiemetic medicines Oral > Liquid: 0.5 mg per 5 mL; 2 mg per 5 mL Oral > Solid: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)	<input type="text" value="Nausea or vomiting"/>

Dolasetron

Section	Indications
Antiemetic medicines	Therapeutic equivalent to <i>ondansetron</i> for <input type="text" value="Nausea or vomiting"/>

Granisetron

Section

Antiemetic medicines

Indications

Therapeutic equivalent to ondansetron for

Nausea or vomiting

Hydrocortisone

Section

Anti-inflammatory medicines

Local > Rectal > Suppository: 25 mg (acetate)

Local > Rectal > Retention enema: 100 mg per 60 mL

Indications

Crohn disease site

Ulcerative colitis

Mesalazine

Section

Anti-inflammatory medicines

Indications

Therapeutic equivalent to sulfasalazine for

Crohn disease site

Therapeutic equivalent to sulfasalazine for

Ulcerative colitis

Metoclopramide

Section

Antiemetic medicines

Oral > Liquid: 5 mg per 5 mL

Oral > Solid: 10 mg (hydrochloride)

Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)

Indications

Nausea or vomiting

Omeprazole

Section

Antiulcer medicines

Oral > Liquid: 20 mg powder for oral liquid; 40 mg powder for oral liquid

Oral > Solid: 10 mg; 20 mg; 40 mg

Parenteral > General injections > IV: 40 mg in vial

Indications

Peptic ulcer, site unspecified

Gastro-oesophageal reflux disease

Gastrointestinal bleeding

Ondansetron

Section

Antiemetic medicines

Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)

Oral > Liquid: 4 mg per 5 mL (ondansetron base)

Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base); 24 mg (ondansetron base)

Indications

Nausea or vomiting

Oral rehydration salts

Section

Oral rehydration

Powder for dilution in 200 mL, 500 mL, 1 L:

- glucose 75 mEq
- sodium 75 mEq or mmol/L
- chloride 65 mEq or mmol/L
- potassium 20 mEq or mmol/L
- citrate 10 mmol/L
- osmolarity 245 mOsm/L
- glucose 13.5 g/L
- sodium chloride 2.6 g/L
- potassium chloride 1.5 g/L
- trisodium citrate dihydrate 2.9 g/L

Indications

Diarrhoea

Oral rehydration salts - zinc sulfate

Section

Medicines used in diarrhoea

Oral > Other: ORS powder for dilution and zinc sulfate solid oral dosage form 20 mg (co-packaged)

Indications

Diarrhoea

Palonosetron

Section

Antiemetic medicines

Indications

Therapeutic equivalent to **ondansetron** for

Nausea or vomiting

Pancreatic enzymes

Section

Pancreatic enzymes

Oral > Solid: Age-appropriate formulations and doses including lipase, protease and amylase.

Indications

Enzyme intestinal

Prednisolone

Section

Anti-inflammatory medicines

Local > Rectal > Retention enema:

Local > Rectal > Retention enema: 20 mg per 100 mL (as sodium phosphate)

Indications

Therapeutic equivalent to [hydrocortisone](#) for

Crohn disease site

Crohn disease site

Ulcerative colitis

Promethazine

Section

Antiemetic medicines

Oral > Liquid: 5 mg per 5 mL (hydrochloride)

Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)

Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule

Indications

Nausea or vomiting

Ranitidine

Section

Antiulcer medicines

Oral > Liquid: 75 mg per 5 mL (as hydrochloride)

Oral > Solid: 150 mg (as hydrochloride)

Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (as hydrochloride)

Indications

Gastro-oesophageal reflux disease

Peptic ulcer, site unspecified

Senna

Section

Laxatives

Oral > Solid: 7.5 mg (sennosides) (or traditional dosage forms)

Indications

Constipation

Sulfasalazine

Section

Anti-inflammatory medicines

Oral > Solid: 500 mg

Local > Rectal > Suppository: 500 mg

Local > Rectal > Retention enema:

Indications

Crohn disease site

Ulcerative colitis

Tropisetron

Section

Antiemetic medicines

Indications

Therapeutic equivalent to ondansetron for

Nausea or vomiting

Zinc sulfate

Section

Medicines for diarrhoea

Oral > Solid: 20 mg

Indications

Diarrhoea